

3 to 5 Years Later: Long-term Effects of Prophylactic Bilateral Salpingectomy on Ovarian Function

This is the peer reviewed version of the following article:

Original:

Venturella, R., Lico, D., Borelli, M., Imbrogno, M.G., Cevenini, G., Zupi, E., et al. (2017). 3 to 5 Years Later: Long-term Effects of Prophylactic Bilateral Salpingectomy on Ovarian Function. JOURNAL OF MINIMALLY INVASIVE GYNECOLOGY, 24(1), 145-150 [10.1016/j.jmig.2016.08.833].

Availability:

This version is available <http://hdl.handle.net/11365/1011085> since 2017-07-06T18:32:42Z

Published:

DOI: <http://doi.org/10.1016/j.jmig.2016.08.833>

Terms of use:

Open Access

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.

(Article begins on next page)

Full title: Three to five years later: long-term effects of prophylactic bilateral salpingectomy on ovarian function.

Precis: The addition of bilateral salpingectomy during hysterectomy is a safe procedure and does not cause ovarian reserve impairment, even 3 to 5 years after surgery.

Authors: Roberta Venturella^a, MD, Daniela Lico^a, MD, Massimo Borelli^b, Phd, Maria G. Imbrogno^a, MD, Gabriele Cevenini^c, Phd, Errico Zupi^d, Phd, Fulvio Zullo^a, Phd, and Michele Morelli^a, Phd.

^a*Department of Obstetrics & Gynecology, "Magna Graecia" University - Catanzaro, Italy.*

^b*Department of Life Sciences, University of Trieste, Italy.*

^c*Department of Medical Biotechnologies, University of Siena, Siena, Italy.*

^d*Department of Biomedicine and Prevention, Obstetrics and Gynecological Clinic, University of Rome "Tor Vergata," Rome, Italy.*

Corresponding author: Roberta Venturella, MD. mail: rovefa@libero.it Phone and fax number: +390961883234. Postal address: Department of Obstetrics & Gynaecology – "Magna Graecia" University - viale Europa, loc. Germaneto, 88100 Catanzaro.

Keywords: cancer prevention; long-term follow-up; ovarian cancer; ovarian reserve; prophylactic bilateral salpingectomy; salpingectomy.

The authors have nothing to disclose.

Structured abstract

Study Objective: Preliminary data on the effects of prophylactic bilateral salpingectomy (PBS) show that postoperative ovarian function is preserved up to 3 months after surgery. The confirmation of PBS safety on ovarian function even many years after surgery is essential to reassure the medical community that this new strategy, recently proposed for the prevention of ovarian cancer, is at least able to avoid the risk of premature surgical menopause. We investigated whether the addition of PBS during total laparoscopic hysterectomy (TLH) causes long-term effects on ovarian function.

Design: Observational study

Design classification: Canadian Task Force II-3

Setting: Department of Obstetrics & Gynecology, "Magna Graecia" University - Catanzaro, Italy.

Patients: Seventy-nine patients who underwent TLH plus salpingectomy between September 2010 and September 2012 at our Institution have been recalled to be submitted to ovarian reserve evaluation in February 2015. Eight of 79 women refused to participate in this follow-up study.

Interventions: The ovarian age of PBS patients has been determined through OvAge®, a statistical model that combines Anti-Müllerian-Hormone (AMH), Follicle-Stimulating-Hormone (FSH), 3D Antral Follicle Count (AFC), Vascular-Index (VI), Flow-Index (FI) and Vascular-Flow-Index (VFI) values. The control group consisted of a large population of 652 healthy women (with intact uterus and adnexa) previously enrolled to build the OvAge® model. Comparisons between ovarian ages

of PBS patients and the control group have been assessed by (ANCOVA) linear statistical modeling.

Measurements and Main Results: The main outcome measurement was the differences in the behavior within OvAge/age relation between PBS and control women. Descriptive statistics of those 71 enrolled PBS patients are the following: age 49.61 ± 2.15 years; OvAge 49.22 ± 2.57 years; FSH 43.02 ± 19.92 mU/mL; AMH 0.12 ± 0.20 ng/mL; 3D AFC 1.91 ± 1.28 ; VI $2.80 \pm 5.32\%$; FI 19.37 ± 5.88 ; and VFI 0.56 ± 1.12 . ANCOVA analysis disclosed that PBS and control women do not exhibit different behavior ($p = 0.900$) within OvAge/age relation.

Conclusion: According to our model, the addition of PBS to TLH in the late reproductive years does not modify the ovarian age of treated women up to 3 to 5 years after surgery.

Main text**Introduction**

Recent literature suggests that many high-grade serous carcinomas develop from the epithelium of the distal fallopian tube, and that serous tubal intraepithelial carcinoma (STIC) represents the putative precursor of these neoplasms (1). According to the new guidelines of the American Cancer Society and the American College of Obstetricians and Gynecologists (2,3), and considering the new theory on the pathogenesis and origin of these aggressive gynecologic cancers, prophylactic bilateral salpingectomy (PBS) has been suggested as a new preventive strategy for average-risk women, not carrying BRCA mutations, who completed their reproductive desire. The rationale of this approach is that, while eliminating the primary source of cancer, PBS in place of standard bilateral salpingo-oophorectomy could also reduce the risk of premature death due to cardiovascular disease seen in women subjected to salpingo-oophorectomy before the onset of natural menopause (4).

On the other hand, two recent publications about the effect of salpingectomy tubal surgery for hydrosalpinx prior to in vitro fertilization (IVF) (5,6) recommended laparoscopic salpingectomy or proximal tubal occlusion in cases of surgically irreparable hydrosalpinges to improve IVF pregnancy rates. Although meta-analytic data clearly demonstrate that salpingectomy increases pregnancy rate in women undergoing IVF (RR 2.24, 95% CI 1.27-3.95) (7), contradictory results are available on the comparison in terms of ovarian response to hyperstimulation during IVF between patients who did and did not undergo salpingectomy (8).

Preliminary data on the safety of PBS showed that postoperative ovarian function is preserved at least 3 months after surgery (9-11), but to date no evidence of the long-term effects of PBS are available in the literature. The confirmation of PBS safety in ovarian function even many years after surgery is essential for reassuring the medical community that the new proposed preventive strategy is at least able to ward off the risk of premature surgical menopause. Adverse health effects of premature surgical menopause include bone resorption; psychosexual, cognitive, and cardiovascular dysfunction; and increased incidence of fatal and non-fatal heart disease (4).

We previously evaluated the short-term effect of PBS on a population of 79 women subjected to total laparoscopic hysterectomy (TLH) with PBS between 2010 and 2012 (9). The goal of the current study was to evaluate the ovarian function of these women up to 5 years after the primary surgery to evaluate, for the first time in the literature, the long-term effect of PBS on ovarian function. For this purpose, we used OvAge®, a validated generalized linear model (GzLM) that combines a patient's biochemical and 3D ultrasonographic values and generates a number that is an estimate of the woman's ovarian age (12).

Materials and methods

This was an observational study conducted at the University "Magna Graecia" of Catanzaro, in the Department of Obstetrics and Gynecology, between February and September 2015.

The procedures used in the study were in accordance with the guidelines of the Helsinki Declaration. The study protocol was approved by the Ethical Committee

of the Department of Gynecology and Obstetrics, University "Magna Graecia" of Catanzaro.

All patients who underwent TLH with PBS for abnormal uterine bleeding related to benign pathology in our Department between September 2010 and September 2012, and who had given their consent to participate in our retrospective analysis in 2013 (9), were identified and contacted by two investigators (MGI and DL).

All these women had undergone TLH and complete bilateral excision of the fallopian tubes. Salpingectomy also had been performed at that time, according to the standard technique, thus sparing the mesosalpinx. The tubes had been coagulated and sectioned, beginning from the very distal fimbrial end, carefully preserving the ovarian vascularization, and proceeding toward the uterine cornu.

Patients who experienced any of the following after the TLH plus PBS were excluded from the study: acute or chronic pelvic inflammatory disorders; malignant neoplasms; chemotherapy or radiotherapy; autoimmune diseases; chronic, metabolic, endocrine, and systemic disorders; had ovarian surgeries; or received estrogen-progestin therapy or metformin in the 2 months prior to enrollment (13). Women with and without menopausal symptoms were analyzed.

Anti-Müllerian-Hormone (AMH), Follicle-Stimulating-Hormone (FSH), Estradiol (E2), 3D Antral Follicle Count (AFC), Vascular-Index (VI), Flow-Index (FI) and Vascular-Flow-Index (VFI) were measured in all women. In ovulating women, ovarian reserve had been evaluated when early follicular phase was confirmed by the absence at ultrasound of a dominant follicle >10 mm in any of the ovaries in conjunction with the presence of serum E2 level <60 pg/mL and P<1 ng/mL. Given

the poor reliability of the FSH values in the presence of E2 levels >60 pg/mL, data from women with basal E2 levels greater than this cut-off were excluded from analysis, and both patients and women with follicles >10 mm in any of the ovaries or with the evidence of corpus luteum were asked to come back 10 to 30 days later, according to their previous menstrual histories or to the dimension of the pre-ovulatory follicle.

A single experienced investigator (DL) performed all of the ultrasound scans using a Voluson-i (GE Healthcare Ultrasound, Zipf, Austria) and a 5–9-MHz transvaginal volume transducer, which has 3D ultrasound scanning modes. AFC and VI were measured using a 3D ultrasound dataset, with a sonography-based automated volume count and virtual organ computer-aided analysis imaging program (VOCAL™) (SonoAVC™ and VOCAL™, GE Healthcare Ultrasound, Zipf, Austria), as previously described (12).

The intra-observer reliability was expressed as the mean intra-class correlation coefficient (ICC) with a 95% CI. The mean ICCs (95% CI) for the 3D scanning of the VI, FI, and VFI were 0.9792 (0.9654, 0.9869), 0.8871 (0.7139, 0.9719), and 0.9929 (0.9748, 0.9967), respectively. The mean ICCs for data acquisition of VI, FI, and VFI were 0.9823 (0.9412, 0.9934), 0.9869 (0.9619, 0.9934), and 0.9825 (0.9513, 0.9977), respectively.

On the same day of the ultrasonography, blood samples obtained by venipuncture were centrifuged within 30 minutes of collection for 10 min at 3500 rpm and 4°C. Aliquots of each serum sample were frozen at -80°C and stored for subsequent assays of AMH, FSH, and E2. To measure serum AMH levels, an AMH-Gen II ELISA assay kit (Beckman Coulter) was used. The lowest detection limit of

AMH is 0.08 ng/mL, and the intra- and inter-assay coefficients of variation were <3.4% and 4.0%, respectively. An ECLIA method was used to measure the levels of serum FSH and E2 by a COBAS e411 auto-analyzer (Roche Diagnostics). The lowest detection limit for FSH was 0.1 IU/L, and the intra- and inter-assay coefficients of variation were <2.6% and 3.5%, respectively. The lowest detection limit for E2 was 18.4 pmol/L, with intra- and inter-assay coefficients of variation of 2% and 3%, respectively.

To determine the long-term effect of surgery on each patient's ovarian reserve, we used OvAge®, a mathematical formula that takes a patient's biochemical and ultrasonographic values as input and generates an easy-to-interpret number, also called OvAge, which is an estimate of a woman's ovarian age, according to the linear relation as follows:
$$\text{OvAge} = 48.05 - 3.14 \cdot \text{AHM} + 0.07 \cdot \text{FSH} - 0.77 \cdot \text{AFC} - 0.11 \cdot \text{FI} + 0.25 \cdot \text{VI} + 0.1 \cdot \text{AMH} \cdot \text{AFC} + 0.02 \cdot \text{FSH} \cdot \text{AFC}$$
 (12). For obtaining this formula, we previously recruited 652 healthy women, aged 18 to 55, with a history of spontaneous conception(s); intact ovaries, fallopian tubes, and uterus; and regular menses with a mean interval of 21 to 35 days to serve as control subjects. Exclusion criteria for the selection of these subjects, enrolled as training population for the OvAge model, had been: estrogen or progestin use or breastfeeding in the 2 months before enrollment; pregnancy; history of female infertility; endometriosis; presence of ovarian follicles measuring more than 10 mm at study entry ultrasonography and other cystic masses of the ovary; history of autoimmune disease; polycystic ovary syndrome (PCOS); ovarian surgery; gynecologic malignancy; previous radiation or chemotherapy; known chronic, endocrine, systemic, and metabolic disease including diabetes mellitus; hyperandrogenism; hyperprolactinemia and thyroid diseases;

hypogonadotropic hypogonadism; and a history of use of a drug that can cause menstrual irregularity.

In the original OvAge study, in women enrolled as healthy fertile controls, the model showed a high level of fit between chronological age and predicted OvAge, meaning that in the absence of risk factors known to be detrimental for ovarian function, chronological age and predicted OvAge are equivalent. Conversely, in patients with known causes of impaired ovarian function, a significant difference between these two parameters was shown, indicating that the formula produced was able to recognize pathological deviation from physiologic gonadal activity (12).

For the purpose of the current study we assumed that, if the addition of PBS to TLH did not have any detrimental effect on ovarian function, considering our previous and current exclusion criteria, the predicted OvAge of our 79 patients would be similar to their chronological age. Specifically, if the addition of PBS to TLH does not cause detrimental effects to ovarian function worse than those reported 3 months after surgery (9), differences in the behavior within OvAge/age relation between PBS and control women would not be significant.

To summarize data, means and standard deviations or absolute frequencies and proportion were addressed to continuous or count covariates, accordingly. To assess differences between OvAge® and age within PBS patients versus the healthy control group, ANCOVA linear modeling was used (14). In all instances, a 0.05 significance level was assumed, and the calculations have been performed by R (15).

Results

From February to September 2015, 79 women were called and asked to participate in this long-term follow-up study. Eight of 79 (10.1%) women refused to participate, leaving 71 patients for the study group. The patients were given the OvAge test, and the results were analyzed. Descriptive statistics of those 71 enrolled women are shown in Table 1.

To confirm the hypothesis that the addition of PBS to TLH did not have any detrimental effects on ovarian function, so that the predicted OvAge of our patients would be similar to their chronological age, we proceeded in the following way. We considered the age of a woman to be a continuous response with respect to OvAge and group (i.e., PBS treated or control) covariates. We addressed several statistical models, as explained in Crawley (14); in particular, we focused on three models: an ANCOVA model with interaction between covariates, an ANCOVA model without interaction, and a simpler linear model (i.e., a regression line) over OvAge without group information. Proceeding by a top-down strategy in model selection according to a deviance analysis (14), we disclosed that all models are equivalent in a statistical sense. This finding allows us to retain, as a minimal adequate model, the simpler model as shown in Table 2.

The simpler model, the minimally adequate one, exhibits the significant role of the OvAge in estimating women ages and, implicitly, neglects any role of the PBS treatment/control covariate, with a very similar residual standard error ($\sigma = 4.208$ on 729 degrees of freedom), but with an excellent multiple R-squared (0.9885). A deviance analysis confirms the equivalence ($p = 0.986$) of two models, confirming the hypothesis that the addition of PBS to TLH did not have any detrimental effect on ovarian function (Fig. 1).

Discussion

By means of OvAge linear model, in the current study we demonstrated that in our population the addition of PBS to TLH in the late reproductive years did not have negative effects on ovarian function, not only a few months after surgery, but also 3 to 5 years later. In our patients, ovarian age (or OvAge) was found to be similar to their chronological age, since linear models implemented for the statistical analysis showed that there is no difference between treated and untreated groups. Although limited to a well-selected population of women undergoing PBS with TLH in their late reproductive years, this is the first published study reporting information about the long-term effect of salpingectomy performed for cancer prophylaxis.

This finding can be supported by the observation that ovarian blood supply is guaranteed both by infundibolopelvic vessels and by the ovarian branch of the uterine artery, which anastomose with each other at the tubal level (16). At the time of bilateral salpingectomy, the whole infundibolopelvic blood volume, which was previously distributed between tubes and ovaries, becomes fully available to the ovaries, ensuring adequate gonadal vascularization and maintained ovarian steroid hormone synthesis (16).

In the current study, for the first time in the literature, a long-term evaluation of the effect of salpingectomy on ovarian age has been carried out on a cohort of patients who underwent standard TLH with PBS 3 to 5 years earlier. This was possible by using both a new algorithm that combines hormonal and three-dimensional ultrasonographic parameters and a very large population of healthy women as the control group (12). This sample of 652 healthy fertile women, previously enrolled to generate the OvAge algorithm, is to date the larger dataset

available for comparing treated and untreated women in terms of ovarian function. It represents the nomogram of reference for the variable “ovarian age,” having been built on a population of women deeply screened for all those factors currently known to be able to affect ovarian function. The assumption on which this study is based is that, if PBS does not cause significant detrimental effects on hormonal and ultrasonographic ovarian parameters, the ovarian age of the treated patients should not deviate significantly (i.e., as to raise at least one OvAge unit/year) from that of women who have never been submitted to this kind of surgery. This assumption was verified by our results; despite a mean 47% decrease in AMH levels from baseline through 3 to 5 years for the 71 PBS women, their OvAge was comparable with that of the control population, meaning that a similar change probably occurs in untreated women in course of their life.

Up to now, the effect of PBS on ovarian function was evaluated at only 3 months postoperatively, and the obtained trend was encouraging in all the studies. In 2007, even before the diffusion of the new theory about the tubal origin of most high-grade serous cancers, Sezik et al.(17) measured hormonal markers of ovarian function (FSH,LH, and estradiol) to evaluate ovarian reserve of women treated by hysterectomy, and they found no difference among women who underwent salpingectomy versus those who did not. In 2013, we published a retrospective analysis in which the ovarian reserve was not reduced in patients in whom PBS was added to TLH performed for benign uterine pathologies (9). In our study, ovarian reserve was evaluated by AMH, FSH, AFC, mean ovarian diameters, and peak systolic velocity. In the same year, Findley et al. (10), in a pilot randomized controlled trial, used AMH to measure ovarian reserve and concluded that salpingectomy at the

time of laparoscopic hysterectomy with ovarian preservation had no short-term deleterious effects on ovarian function.

In 2015, at our institution, women undergoing laparoscopic surgery for uterine myoma or tubal surgical sterilization were randomly subjected to standard salpingectomy or wide mesosalpinx excision (11). Ovarian function was evaluated by the measurement prior to and 3 months after surgery of AMH, FSH, three-dimensional indexes, and OvAge for each patient. We found no significant difference between groups, and we speculated that even when the surgical excision includes the removal of the entire mesosalpinx, salpingectomy does not damage the ovarian reserve (11).

Recently, a systematic review and meta-analysis of studies comparing the pregnancy outcomes of patients diagnosed with hydrosalpinx treated with salpingectomy versus those treated with proximal tubal occlusion prior to IVF showed comparable responses to controlled ovarian hyperstimulation and pregnancy outcomes between the groups (18). This is an additional demonstration that salpingectomy does not worsen the reproductive prognosis of patients who choose excisional surgery. On the other hand, results on ovarian response to medical induction of superovulation in patients treated by salpingectomy are conflicting (19-21), but authors agree that bilateral salpingectomy has been proved to be a safe procedure at least in terms of surgical outcome and short-term postoperative complications (22). Moreover, three different meta-analyses also reported increased ongoing and clinical pregnancy rates with salpingectomy versus no intervention and no difference in the clinical pregnancy rates between salpingectomy and tubal occlusion (7,23,24).

The strength of the current study is the long-term evaluation, for the first time in the literature, of a well-screened population of women subjected to PBS in the absence of tubal pathologies, even when in their late reproductive years. All the currently known confounding factors that can interfere on ovarian reserve were listed among the exclusion criteria for both the first (9) and the current study. Reliable and definitive information on the safety of PBS is critically important for the entire medical community, considering that the idea that salpingectomy should be proposed as a preventive strategy for low-risk women who undergo gynecologic surgery is gaining increasing popularity. Moreover, although risk-reducing salpingo-oophorectomy (RRSO) at around age 40 years is currently recommended to women who carry the BRCA1/2 mutation, an alternative preventive strategy has been put forward for high-risk women: early PBS and delayed oophorectomy (RRO). Although preventive RRSO decreases the ovarian cancer risk by 80% to 96%, based on its short-term and long-term morbidity, which potentially affect quality of life, and on recent insights into the fallopian tube as the possible site of origin of serous ovarian carcinomas, early PBS and delayed RRO could be the best solution.

Recently, a multi-center non-randomized trial has started enrollment, and participants will choose between standard RRSO at age 35 to 40 years (BRCA1) or 40 to 45 years (BRCA2) and the alternative strategy of PBS on completion of childbearing and RRO at age 40 to 45 years (BRCA1) or 45 to 50 years (BRCA2). The aim of the study is to measure menopause-related quality of life, but also ovarian/breast cancer incidence, surgery-related morbidity, histopathology, cardiovascular risk factors and diseases, and cost-effectiveness (25).

The increasing interest in PBS as a preventive strategy for both low- and high-risk women and the related crucial importance of long-term follow-up data are

confirmed by many web-based surveys of health professionals' acceptability/attitude toward the new proposal (26-29).

A limitation of the study is the relatively small sample size of women evaluated. However, given the homogeneous distribution of results among this population, and considering that this is currently the largest sample of women studied in the literature, the results presented deserve attention. Another limitation of the current study is that the control population is not the same used in the original study (9), that being composed of women treated by standard TLH (with adnexal preservation) between September 2008 and September 2010, matched for uterine weight. Unfortunately, none of these women had undergone an OvAge test at that time, and the gap of average age of at least 2 years makes it impossible to accurately compare that group with the PBS population. Age at menopause of these women will be an interesting outcome for forthcoming analysis, despite the need of consensus for a correct definition for women without a uterus. In this context, further research on the validation of the OvAge model to accurately predict the time to menopause is already in progress.

Two other possible limits of our study are: first, the range of time postoperatively at which the subjects have being evaluated creates a variable in comparison of the data. However, according to our previous experience (30), when there is evidence of impaired ovarian reserve caused by vascular damage, the modification of biochemical and ultrasonographic parameters is already evident at 3 months follow-up, and it remains constant at 1 and 2 years. In the same way, reassuring parameters at 3 months do not worsen at the next re-evaluation, allowing us to speculate that the results of the current study are poorly affected by the range of evaluation times.

Furthermore, at surgery and at the time of follow-up, our patients already had evidence of diminished ovarian reserve, given their mean age of 45.97 ± 2.36 and 49.61 ± 2.15 , respectively. Although it would be better to assess the impact of surgery on younger women, in which ovaries have greater potential for functional loss, unfortunately any assessment of the impact of prophylactic salpingectomy on ovarian reserve is not ethically conducive in patients who have not yet fulfilled their reproductive desire. That is why, to date, in all studies conducted in patients undergoing salpingectomy for cancer prophylaxis, in the absence of tubal pathology and not for reproductive intent, the average age of the population has always been more than 37 years (10,11,17). Ideally, the same rigorous approach used for women subjected to PBS should be applied on all future studies aimed at evaluating the effect of this kind of surgery in women undergoing IVF, to extend the assessable population and to definitively conclude the mid- and long-term safety of laparoscopic salpingectomy.

In conclusion, according to our results, the addition of PBH during TLH in the late reproductive years does not cause ovarian function impairment up to 3 to 5 years after surgery.

Table legends

Table 1. **Descriptive statistics of enrolled women**

Table 2. **Linear models applied.** Two models are summarized. In model 1, an ANCOVA with interaction (maximal) model is applied. A simpler model, the minimally adequate one, neglects any role of the treatment/control covariate, with a very similar residual standard error, confirming the hypothesis that the addition of PBS to TLH had no detrimental effect on ovarian function.

Figure Legend

Figure 1: Two-dimensional plot for regression analysis. The OvAge control group (gray and white bullets) and the TLH plus PBS treated group (black and dark gray diamonds). In the panel, three regression lines have been drawn: a 'two-parameters' thin dashed line, with intercept and slope specific for the control group; a 'two-parameters' thin continuous line, with intercept and slope specific for the treated group; and, a 'one-parameter' solid dashed line, with common slope (and null intercept) for both groups. All three regression lines do not differ in a statistical sense; consequently, the 'one-parameter common line' is the minimal adequate model to retain as a valid description of both groups.

References

1. Crum CP, Drapkin R, Miron A, Ince TA, Muto M, Kindelberger DW et al. The distal fallopian tube: a new model for pelvic serous carcinogenesis. *Curr Opin Obstet Gynecol* 2007;19:3-9.
2. ACOG Committee opinion no. 620. Salpingectomy for ovarian cancer prevention. *Obstet Gynecol* 2015;125:279-81.
3. Walker JL, Powell CB, Chen LM, Carter J, Bae Jump VL, Parker LP et al. Society of Gynecologic Oncology recommendations for the prevention of ovarian cancer. *Cancer* 2015. DOI: 10.1002/cncr.29321. [Epub ahead of print]
4. Parker WH, Broder MS, Chang E, Feskanich D, Farquhar C, Liu Z et al. Ovarian conservation at the time of hysterectomy and long-term health outcomes in the nurses' health study. *Obstet Gynecol* 2009;113:1027–37.
5. Practice Committee of American Society for Reproductive Medicine in collaboration with Society of Reproductive Surgeons. Salpingectomy for hydrosalpinx prior to in vitro fertilization. *Fertil Steril* 2008;90(5 Suppl):S66-8.
6. Practice Committee of the American Society for Reproductive Medicine. Role of tubal surgery in the era of assisted reproductive technology: a committee opinion. *Fertil Steril* 2015;103:37-43.
7. Tsiami A, Chaimani A, Mavridis D, Siskou M, Assimakopoulos E, Sotiriadis A. Surgical treatment for hydrosalpinx prior to IVF-ET: a network meta-analysis. *Ultrasound Obstet Gynecol* 2016. DOI: 10.1002/uog.15900. [Epub ahead of print]

8. Fan M, Ma L. Effect of salpingectomy on ovarian response to hyperstimulation during in vitro fertilization: a meta-analysis. *Fertil Steril* 2016. DOI: 10.1016/j.fertnstert.2016.03.053. [Epub ahead of print]
9. Morelli M, Venturella R, Mocciaro R, Di Cello A, Rania E, Lico D et al. Prophylactic salpingectomy in premenopausal low-risk women for ovarian cancer: primum non nocere. *Gynecol Oncol* 2013;129:448-51.
10. Findley AD, Siedhoff MT, Hobbs KA, Steege JF, Carey ET, McCall CA et al. Short-term effects of salpingectomy during laparoscopic hysterectomy on ovarian reserve: a pilot randomized controlled trial. *Fertil Steril* 2013;100:1704-8.
11. Venturella R, Morelli M, Lico D, Di Cello A, Rocca M, Sacchinelli A et al. Wide excision of soft tissues adjacent to the ovary and fallopian tube does not impair the ovarian reserve in women undergoing prophylactic bilateral salpingectomy: results from a randomized, controlled trial. *Fertil Steril* 2015;104:1332-9.
12. Venturella R, Lico D, Sarica A, Falbo MP, Gulletta E, Morelli M et al. OvAge: a new methodology to quantify ovarian reserve combining clinical, biochemical and 3D-ultrasonographic parameters. *J Ovarian Res* 2015;8:8-21.
13. Falbo A, Orio F, Venturella R, Rania E, Materazzo C, Tolino A et al. Does metformin affect ovarian morphology in patients with polycystic ovary syndrome? A retrospective cross-sectional preliminary analysis. *J Ovarian Res* 2009;2:5.
14. Crawley, Michael J, *Statistics: an introduction using R*, 2005, John Wiley & Sons;2005.

15. R Core Team. *R: A Language and Environment for Statistical Computing*.
R Foundation for Statistical Computing, Vienna, Austria, 2014.
16. Dietl J and Wischhusen J. The postreproductive salpingectomy. *Fertil Steril* 2014;101.
17. Sezik M, Ozkaya O, Demir F, Sezik HT, Kaya H. Total salpingectomy during abdominal hysterectomy: effects on ovarian reserve and ovarian stromal blood flow. *J Obstet Gynaecol Res* 2007;33:863-9.
18. Zhang Y, Sun Y, Guo Y, Li TC, Duan H. Salpingectomy and Proximal Tubal Occlusion for Hydrosalpinx Prior to In Vitro Fertilization: A Meta-analysis of Randomized Controlled Trials. *Obstet Gynecol Surv* 2015;70:33-38.
19. Strandell A, Lindhard A, Waldenström U, Thorburn J. Prophylactic salpingectomy does not impair the ovarian response in IVF treatment. *Hum Reprod* 2001;16:1135-9.
20. Chan CC, Ng EH, Li CF, Ho PC. Impaired ovarian blood flow and reduced antral follicle count following laparoscopic salpingectomy for ectopic pregnancy. *Hum Reprod* 2003;18:2175-80.
21. Ye XP, Yang YZ, Sun XX. A retrospective analysis of the effect of salpingectomy on serum antiMüllerian hormone level and ovarian reserve. *Am J Obstet Gynecol* 2015;212:53.e1-10.
22. Oliver Perez MR, Magriñá J, García AT, Jiménez Lopez JS. Prophylactic salpingectomy and prophylactic salpingoophorectomy for adnexal high-grade serous epithelial carcinoma: A reappraisal. *Surg Oncol* 2015;24:335-44.

23. Johnson N, van Voorst S, Sowter MC, Strandell A, Mol BW. Surgical treatment for tubal disease in women due to undergo in vitro fertilisation. *Cochrane Database Syst Rev* 2010(1):CD002125
24. Zhang Y, Sun Y, Guo Y, Li TC, Duan H. Salpingectomy and proximal tubal occlusion for hydrosalpinx prior to in vitro fertilization: a meta-analysis of randomized controlled trials. *Obstet Gynecol Surv* 2015;70:33-8.
25. Harmsen MG, Arts-de Jong M, Hoogerbrugge N, Maas AH, Prins JB, Bulten J et al. Early salpingectomy (TUbectomy) with delayed oophorectomy to improve quality of life as alternative for risk-reducing salpingo-oophorectomy in BRCA1/2 mutation carriers (TUBA study): a prospective non-randomised multicentre study. *BMC Cancer* 2015;15:593.
26. Chandrasekaran D, Menon U, Evans G, Crawford R, Saridogan E, Jacobs C et al. Risk reducing salpingectomy and delayed oophorectomy in high risk women: views of cancer geneticists, genetic counsellors and gynaecological oncologists in the UK. *Fam Cancer* 2015;14:521-30.
27. Venturella R, Rocca M, Lico D, Trapasso S, Di Cello A, Gizzo S et al. Prophylactic bilateral salpingectomy for the prevention of ovarian cancers: what is happening in Italy? *Eur J Cance Prev* 2015 [Epub ahead of print]
28. Reade CJ, Finlayson S, McAlpine J, Tone AA, Fung-Kee-Fung M, Ferguson SE. Risk-reducing salpingectomy in Canada: a survey of obstetrician-Gynaecologists. *J Obstet Gynaecol Can* 2013;35:627-34.
29. Gill SE, Mills BB. Physician opinions regarding elective bilateral salpingectomy with hysterectomy and for sterilization. *J Minim Invasive Gynecol* 2013;20:517-521.


- 479 30. Morelli M, Mocciaro R, Venturella R, Imperatore A, Lico D, Zullo F. Mesial
480 side ovarian incision for laparoscopic dermoid cystectomy: a safe and
481 ovarian tissue-preserving technique. Fertil Steril 2012;98:1336-40.

Table 1. Descriptive statistics of enrolled women

Parameters	Mean values \pm SD
Age at surgery (years)	45.85 \pm 2.40
Age at follow-up (years)	49.61 \pm 2.15
OvAge at follow-up (years)	49.34 \pm 2.12
FSH at follow-up (mU/mL)	43.02 \pm 19.92
AMH at follow-up (ng/mL)	0.12 \pm 0.20
3D AFC at follow-up (n.)	1.91 \pm 1.28
VI at follow-up (%)	2.80 \pm 5.32
FI at follow-up (1-100)	19.37 \pm 5.88
VFI at follow-up (1-100)	0.56 \pm 1.12

Table 2. **Linear models applied.** Two models are summarized. In model 1, an Ancova with interaction (maximal) model is approached. A simpler model, the minimal adequate one, neglect any role of the Treatment/control covariate, with a very similar residual standard error, confirming the hypothesis that the addition of PBS to TLH did not have any detrimental effect on ovarian function.

Maximal model			
	Estimate	Standard Error	p-value
Intercept	0.1005	0.8474	.906
OvAge	1.0028	0.0225	< .001
Treatment	1.5632	11.2572	.890
OvAge : Treatment	0.0286	0.2282	.900
Minimal adequate model			
	Estimate	Standard Error	p-value
OvAge	1.0008	0.0040	< .001



Precis:

The addition of bilateral salpingectomy during hysterectomy is a safe procedure and does not cause ovarian reserve impairment, even three to five years after surgery.