



3 to 5 Years Later: Long-term Effects of Prophylactic Bilateral Salpingectomy on Ovarian Function

This is the peer reviewed version of the following article:

Original:

Venturella, R., Lico, D., Borelli, M., Imbrogno, M.G., Cevenini, G., Zupi, E., et al. (2017). 3 to 5 Years Later: Long-term Effects of Prophylactic Bilateral Salpingectomy on Ovarian Function. JOURNAL OF MINIMALLY INVASIVE GYNECOLOGY, 24(1), 145-150 [10.1016/j.jmig.2016.08.833].

Availability:

This version is available <http://hdl.handle.net/11365/1011085> since 2017-07-06T18:32:42Z

Published:

DOI:10.1016/j.jmig.2016.08.833

Terms of use:

Open Access

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.

(Article begins on next page)

1 **Full title: Three to five years later: long-term effects of prophylactic bilateral**
2 **salpingectomy on ovarian function.**

3 **Precis:** The addition of bilateral salpingectomy during hysterectomy is a safe
4 procedure and does not cause ovarian reserve impairment, even 3 to 5 years after
5 surgery.

6 **Authors:** Roberta Venturella^a, MD, Daniela Lico^a, MD, Massimo Borelli^b, Phd, Maria
7 G. Imbrogno^a, MD, Gabriele Cevenini^c, Phd, Errico Zupi^d, Phd, Fulvio Zullo^a, Phd,
8 and Michele Morelli^a, Phd.

9 ^a*Department of Obstetrics & Gynecology, "Magna Graecia" University - Catanzaro,*
10 *Italy.*

11 ^b*Department of Life Sciences, University of Trieste, Italy.*

12 ^c*Department of Medical Biotechnologies, University of Siena, Siena, Italy.*

13 ^d*Department of Biomedicine and Prevention, Obstetrics and Gynecological Clinic,*
14 *University of Rome "Tor Vergata," Rome, Italy.*

15 **Corresponding author:** Roberta Venturella, MD. mail: rovefa@libero.it Phone and
16 fax number: +390961883234. Postal address: Department of Obstetrics &
17 Gynaecology – "Magna Graecia" University - viale Europa, loc. Germaneto, 88100
18 Catanzaro.

19 **Keywords:** cancer prevention; long-term follow-up; ovarian cancer; ovarian reserve;
20 prophylactic bilateral salpingectomy; salpingectomy.

21 **The authors have nothing to disclose.**

22

23 Structured abstract

24 **Study Objective:** Preliminary data on the effects of prophylactic bilateral
25 salpingectomy (PBS) show that postoperative ovarian function is preserved up to 3
26 months after surgery. The confirmation of PBS safety on ovarian function even many
27 years after surgery is essential to reassure the medical community that this new
28 strategy, recently proposed for the prevention of ovarian cancer, is at least able to
29 avoid the risk of premature surgical menopause. We investigated whether the
30 addition of PBS during total laparoscopic hysterectomy (TLH) causes long-term
31 effects on ovarian function.

32 **Design:** Observational study

33 **Design classification:** Canadian Task Force II-3

34 **Setting:** Department of Obstetrics & Gynecology, "Magna Graecia" University -
35 Catanzaro, Italy.

36 **Patients:** Seventy-nine patients who underwent TLH plus salpingectomy between
37 September 2010 and September 2012 at our Institution have been recalled to be
38 submitted to ovarian reserve evaluation in February 2015. Eight of 79 women
39 refused to participate in this follow-up study.

40 **Interventions:** The ovarian age of PBS patients has been determined through
41 OvAge®, a statistical model that combines Anti-Müllerian-Hormone (AMH), Follicle-
42 Stimulating-Hormone (FSH), 3D Antral Follicle Count (AFC), Vascular-Index (VI),
43 Flow-Index (FI) and Vascular-Flow-Index (VFI) values. The control group consisted
44 of a large population of 652 healthy women (with intact uterus and adnexa)
45 previously enrolled to build the OvAge® model. Comparisons between ovarian ages

46 of PBS patients and the control group have been assessed by (ANCOVA) linear
47 statistical modeling.

48 **Measurements and Main Results:** The main outcome measurement was the
49 differences in the behavior within OvAge/age relation between PBS and control
50 women. Descriptive statistics of those 71 enrolled PBS patients are the following:
51 age 49.61 ± 2.15 years; OvAge 49.22 ± 2.57 years; FSH 43.02 ± 19.92 mU/mL; AMH
52 0.12 ± 0.20 ng/mL; 3D AFC 1.91 ± 1.28 ; VI $2.80 \pm 5.32\%$; FI 19.37 ± 5.88 ; and VFI
53 0.56 ± 1.12 . ANCOVA analysis disclosed that PBS and control women do not exhibit
54 different behavior ($p = 0.900$) within OvAge/age relation.

55 **Conclusion:** According to our model, the addition of PBS to TLH in the late
56 reproductive years does not modify the ovarian age of treated women up to 3 to 5
57 years after surgery.

58

59 **Main text**60 **Introduction**

61 Recent literature suggests that many high-grade serous carcinomas develop from
62 the epithelium of the distal fallopian tube, and that serous tubal intraepithelial
63 carcinoma (STIC) represents the putative precursor of these neoplasms (1).

64 According to the new guidelines of the American Cancer Society and the American
65 College of Obstetricians and Gynecologists (2,3), and considering the new theory on
66 the pathogenesis and origin of these aggressive gynecologic cancers, prophylactic
67 bilateral salpingectomy (PBS) has been suggested as a new preventive strategy for
68 average-risk women, not carrying BRCA mutations, who completed their
69 reproductive desire. The rationale of this approach is that, while eliminating the
70 primary source of cancer, PBS in place of standard bilateral salpingo-oophorectomy
71 could also reduce the risk of premature death due to cardiovascular disease seen in
72 women subjected to salpingo-oophorectomy before the onset of natural menopause
73 (4).

74 On the other hand, two recent publications about the effect of salpingectomy
75 tubal surgery for hydrosalpinx prior to in vitro fertilization (IVF) (5,6) recommended
76 laparoscopic salpingectomy or proximal tubal occlusion in cases of surgically
77 irreparable hydrosalpinges to improve IVF pregnancy rates. Although meta-analytic
78 data clearly demonstrate that salpingectomy increases pregnancy rate in women
79 undergoing IVF (RR 2.24, 95% CI 1.27-3.95) (7), contradictory results are available
80 on the comparison in terms of ovarian response to hyperstimulation during IVF
81 between patients who did and did not undergo salpingectomy (8).

82 Preliminary data on the safety of PBS showed that postoperative ovarian
83 function is preserved at least 3 months after surgery (9-11), but to date no evidence
84 of the long-term effects of PBS are available in the literature. The confirmation of
85 PBS safety in ovarian function even many years after surgery is essential for
86 reassuring the medical community that the new proposed preventive strategy is at
87 least able to ward off the risk of premature surgical menopause. Adverse health
88 effects of premature surgical menopause include bone resorption; psychosexual,
89 cognitive, and cardiovascular dysfunction; and increased incidence of fatal and non-
90 fatal heart disease (4).

91 We previously evaluated the short-term effect of PBS on a population of 79
92 women subjected to total laparoscopic hysterectomy (TLH) with PBS between 2010
93 and 2012 (9). The goal of the current study was to evaluate the ovarian function of
94 these women up to 5 years after the primary surgery to evaluate, for the first time in
95 the literature, the long-term effect of PBS on ovarian function. For this purpose, we
96 used OvAge®, a validated generalized linear model (GzLM) that combines a
97 patient's biochemical and 3D ultrasonographic values and generates a number that
98 is an estimate of the woman's ovarian age (12).

99 **Materials and methods**

100 This was an observational study conducted at the University "Magna Graecia" of
101 Catanzaro, in the Department of Obstetrics and Gynecology, between February and
102 September 2015.

103 The procedures used in the study were in accordance with the guidelines of
104 the Helsinki Declaration. The study protocol was approved by the Ethical Committee

105 of the Department of Gynecology and Obstetrics, University "Magna Graecia" of
106 Catanzaro.

107 All patients who underwent TLH with PBS for abnormal uterine bleeding
108 related to benign pathology in our Department between September 2010 and
109 September 2012, and who had given their consent to participate in our retrospective
110 analysis in 2013 (9), were identified and contacted by two investigators (MGI and
111 DL).

112 All these women had undergone TLH and complete bilateral excision of the
113 fallopian tubes. Salpingectomy also had been performed at that time, according to
114 the standard technique, thus sparing the mesosalpinx. The tubes had been
115 coagulated and sectioned, beginning from the very distal fimbrial end, carefully
116 preserving the ovarian vascularization, and proceeding toward the uterine cornu.

117 Patients who experienced any of the following after the TLH plus PBS were
118 excluded from the study: acute or chronic pelvic inflammatory disorders; malignant
119 neoplasms; chemotherapy or radiotherapy; autoimmune diseases; chronic,
120 metabolic, endocrine, and systemic disorders; had ovarian surgeries; or received
121 estrogen-progestin therapy or metformin in the 2 months prior to enrollment (13).
122 Women with and without menopausal symptoms were analyzed.

123 Anti-Müllerian-Hormone (AMH), Follicle-Stimulating-Hormone (FSH), Estradiol
124 (E2), 3D Antral Follicle Count (AFC), Vascular-Index (VI), Flow-Index (FI) and
125 Vascular-Flow-Index (VFI) were measured in all women. In ovulating women, ovarian
126 reserve had been evaluated when early follicular phase was confirmed by the
127 absence at ultrasound of a dominant follicle >10 mm in any of the ovaries in
128 conjunction with the presence of serum E2 level <60 pg/mL and P<1 ng/mL. Given

129 the poor reliability of the FSH values in the presence of E2 levels >60 pg/mL, data
130 from women with basal E2 levels greater than this cut-off were excluded from
131 analysis, and both patients and women with follicles >10 mm in any of the ovaries or
132 with the evidence of corpus luteum were asked to come back 10 to 30 days later,
133 according to their previous menstrual histories or to the dimension of the pre-
134 ovulatory follicle.

135 A single experienced investigator (DL) performed all of the ultrasound scans
136 using a Voluson-i (GE Healthcare Ultrasound, Zipf, Austria) and a 5–9-MHz
137 transvaginal volume transducer, which has 3D ultrasound scanning modes. AFC and
138 VI were measured using a 3D ultrasound dataset, with a sonography-based
139 automated volume count and virtual organ computer-aided analysis imaging program
140 (VOCAL™) (SonoAVC™ and VOCAL™, GE Healthcare Ultrasound, Zipf, Austria),
141 as previously described (12).

142 The intra-observer reliability was expressed as the mean intra-class
143 correlation coefficient (ICC) with a 95% CI. The mean ICCs (95% CI) for the 3D
144 scanning of the VI, FI, and VFI were 0.9792 (0.9654, 0.9869), 0.8871 (0.7139,
145 0.9719), and 0.9929 (0.9748, 0.9967), respectively. The mean ICCs for data
146 acquisition of VI, FI, and VFI were 0.9823 (0.9412, 0.9934), 0.9869 (0.9619, 0.9934),
147 and 0.9825 (0.9513, 0.9977), respectively.

148 On the same day of the ultrasonography, blood samples obtained by
149 venipuncture were centrifuged within 30 minutes of collection for 10 min at 3500 rpm
150 and 4°C. Aliquots of each serum sample were frozen at -80°C and stored for
151 subsequent assays of AMH, FSH, and E2. To measure serum AMH levels, an AMH-
152 Gen II ELISA assay kit (Beckman Coulter) was used. The lowest detection limit of

153 AMH is 0.08 ng/mL, and the intra- and inter-assay coefficients of variation were
154 <3.4% and 4.0%, respectively. An ECLIA method was used to measure the levels of
155 serum FSH and E2 by a COBAS e411 auto-analyzer (Roche Diagnostics). The
156 lowest detection limit for FSH was 0.1 IU/L, and the intra- and inter-assay
157 coefficients of variation were <2.6% and 3.5%, respectively. The lowest detection
158 limit for E2 was 18.4 pmol/L, with intra- and inter-assay coefficients of variation of 2%
159 and 3%, respectively.

160 To determine the long-term effect of surgery on each patient's ovarian reserve,
161 we used OvAge®, a mathematical formula that takes a patient's biochemical and
162 ultrasonographic values as input and generates an easy-to-interpret number, also
163 called OvAge, which is an estimate of a woman's ovarian age, according to the linear
164 relation as follows: $OvAge = 48.05 - 3.14*AMH + 0.07*FSH - 0.77*AFC - 0.11*FI +$
165 $0.25*VI + 0.1*AMH*AFC + 0.02*FSH*AFC$ (12). For obtaining this formula, we
166 previously recruited 652 healthy women, aged 18 to 55, with a history of
167 spontaneous conception(s); intact ovaries, fallopian tubes, and uterus; and regular
168 menses with a mean interval of 21 to 35 days to serve as control subjects. Exclusion
169 criteria for the selection of these subjects, enrolled as training population for the
170 OvAge model, had been: estrogen or progestin use or breastfeeding in the 2 months
171 before enrollment; pregnancy; history of female infertility; endometriosis; presence of
172 ovarian follicles measuring more than 10 mm at study entry ultrasonography and
173 other cystic masses of the ovary; history of autoimmune disease; polycystic ovary
174 syndrome (PCOS); ovarian surgery; gynecologic malignancy; previous radiation or
175 chemotherapy; known chronic, endocrine, systemic, and metabolic disease including
176 diabetes mellitus; hyperandrogenism; hyperprolactinemia and thyroid diseases;

177 hypogonadotropic hypogonadism; and a history of use of a drug that can cause
178 menstrual irregularity.

179 In the original OvAge study, in women enrolled as healthy fertile controls, the
180 model showed a high level of fit between chronological age and predicted OvAge,
181 meaning that in the absence of risk factors known to be detrimental for ovarian
182 function, chronological age and predicted OvAge are equivalent. Conversely, in
183 patients with known causes of impaired ovarian function, a significant difference
184 between these two parameters was shown, indicating that the formula produced was
185 able to recognize pathological deviation from physiologic gonadal activity (12).

186 For the purpose of the current study we assumed that, if the addition of PBS
187 to TLH did not have any detrimental effect on ovarian function, considering our
188 previous and current exclusion criteria, the predicted OvAge of our 79 patients would
189 be similar to their chronological age. Specifically, if the addition of PBS to TLH does
190 not cause detrimental effects to ovarian function worse than those reported 3 months
191 after surgery (9), differences in the behavior within OvAge/age relation between PBS
192 and control women would not be significant.

193 To summarize data, means and standard deviations or absolute frequencies
194 and proportion were addressed to continuous or count covariates, accordingly. To
195 assess differences between OvAge® and age within PBS patients versus the healthy
196 control group, ANCOVA linear modeling was used (14). In all instances, a 0.05
197 significance level was assumed, and the calculations have been performed by R
198 (15).

199 **Results**

200 From February to September 2015, 79 women were called and asked to participate
201 in this long-term follow-up study. Eight of 79 (10.1%) women refused to participate,
202 leaving 71 patients for the study group. The patients were given the OvAge test, and
203 the results were analyzed. Descriptive statistics of those 71 enrolled women are
204 shown in Table 1.

205 To confirm the hypothesis that the addition of PBS to TLH did not have any
206 detrimental effects on ovarian function, so that the predicted OvAge of our patients
207 would be similar to their chronological age, we proceeded in the following way. We
208 considered the age of a woman to be a continuous response with respect to OvAge
209 and group (i.e., PBS treated or control) covariates. We addressed several statistical
210 models, as explained in Crawley (14); in particular, we focused on three models: an
211 ANCOVA model with interaction between covariates, an ANCOVA model without
212 interaction, and a simpler linear model (i.e., a regression line) over OvAge without
213 group information. Proceeding by a top-down strategy in model selection according
214 to a deviance analysis (14), we disclosed that all models are equivalent in a
215 statistical sense. This finding allows us to retain, as a minimal adequate model, the
216 simpler model as shown in Table 2.

217 The simpler model, the minimally adequate one, exhibits the significant role of
218 the OvAge in estimating women ages and, implicitly, neglects any role of the PBS
219 treatment/control covariate, with a very similar residual standard error ($\sigma = 4.208$
220 on 729 degrees of freedom), but with an excellent multiple R-squared (0.9885). A
221 deviance analysis confirms the equivalence ($p = 0.986$) of two models, confirming
222 the hypothesis that the addition of PBS to TLH did not have any detrimental effect on
223 ovarian function (Fig. 1).

224 Discussion

225 By means of OvAge linear model, in the current study we demonstrated that in our
226 population the addition of PBS to TLH in the late reproductive years did not have
227 negative effects on ovarian function, not only a few months after surgery, but also 3
228 to 5 years later. In our patients, ovarian age (or OvAge) was found to be similar to
229 their chronological age, since linear models implemented for the statistical analysis
230 showed that there is no difference between treated and untreated groups. Although
231 limited to a well-selected population of women undergoing PBS with TLH in their late
232 reproductive years, this is the first published study reporting information about the
233 long-term effect of salpingectomy performed for cancer prophylaxis.

234 This finding can be supported by the observation that ovarian blood supply is
235 guaranteed both by infundibolopelvic vessels and by the ovarian branch of the
236 uterine artery, which anastomose with each other at the tubal level (16). At the time
237 of bilateral salpingectomy, the whole infundibolopelvic blood volume, which was
238 previously distributed between tubes and ovaries, becomes fully available to the
239 ovaries, ensuring adequate gonadal vascularization and maintained ovarian steroid
240 hormone synthesis (16).

241 In the current study, for the first time in the literature, a long-term evaluation of
242 the effect of salpingectomy on ovarian age has been carried out on a cohort of
243 patients who underwent standard TLH with PBS 3 to 5 years earlier. This was
244 possible by using both a new algorithm that combines hormonal and three-
245 dimensional ultrasonographic parameters and a very large population of healthy
246 women as the control group (12). This sample of 652 healthy fertile women,
247 previously enrolled to generate the OvAge algorithm, is to date the larger dataset

248 available for comparing treated and untreated women in terms of ovarian function. It
249 represents the nomogram of reference for the variable “ovarian age,” having been
250 built on a population of women deeply screened for all those factors currently known
251 to be able to affect ovarian function. The assumption on which this study is based is
252 that, if PBS does not cause significant detrimental effects on hormonal and
253 ultrasonographic ovarian parameters, the ovarian age of the treated patients should
254 not deviate significantly (i.e., as to raise at least one OvAge unit/year) from that of
255 women who have never been submitted to this kind of surgery. This assumption was
256 verified by our results; despite a mean 47% decrease in AMH levels from baseline
257 through 3 to 5 years for the 71 PBS women, their OvAge was comparable with that
258 of the control population, meaning that a similar change probably occurs in untreated
259 women in course of their life.

260 Up to now, the effect of PBS on ovarian function was evaluated at only 3
261 months postoperatively, and the obtained trend was encouraging in all the studies. In
262 2007, even before the diffusion of the new theory about the tubal origin of most high-
263 grade serous cancers, Sezik et al.(17) measured hormonal markers of ovarian
264 function (FSH,LH, and estradiol) to evaluate ovarian reserve of women treated by
265 hysterectomy, and they found no difference among women who underwent
266 salpingectomy versus those who did not. In 2013, we published a retrospective
267 analysis in which the ovarian reserve was not reduced in patients in whom PBS was
268 added to TLH performed for benign uterine pathologies (9). In our study, ovarian
269 reserve was evaluated by AMH, FSH, AFC, mean ovarian diameters, and peak
270 systolic velocity. In the same year, Findley et al. (10), in a pilot randomized controlled
271 trial, used AMH to measure ovarian reserve and concluded that salpingectomy at the

272 time of laparoscopic hysterectomy with ovarian preservation had no short-term
273 deleterious effects on ovarian function.

274 In 2015, at our institution, women undergoing laparoscopic surgery for uterine
275 myoma or tubal surgical sterilization were randomly subjected to standard
276 salpingectomy or wide mesosalpinx excision (11). Ovarian function was evaluated by
277 the measurement prior to and 3 months after surgery of AMH, FSH, three-
278 dimensional indexes, and OvAge for each patient. We found no significant difference
279 between groups, and we speculated that even when the surgical excision includes
280 the removal of the entire mesosalpinx, salpingectomy does not damage the ovarian
281 reserve (11).

282 Recently, a systematic review and meta-analysis of studies comparing the
283 pregnancy outcomes of patients diagnosed with hydrosalpinx treated with
284 salpingectomy versus those treated with proximal tubal occlusion prior to IVF
285 showed comparable responses to controlled ovarian hyperstimulation and pregnancy
286 outcomes between the groups (18). This is an additional demonstration that
287 salpingectomy does not worsen the reproductive prognosis of patients who choose
288 excisional surgery. On the other hand, results on ovarian response to medical
289 induction of superovulation in patients treated by salpingectomy are conflicting (19-
290 21), but authors agree that bilateral salpingectomy has been proved to be a safe
291 procedure at least in terms of surgical outcome and short-term postoperative
292 complications (22). Moreover, three different meta-analyses also reported increased
293 ongoing and clinical pregnancy rates with salpingectomy versus no intervention and
294 no difference in the clinical pregnancy rates between salpingectomy and tubal
295 occlusion (7,23,24).

296 The strength of the current study is the long-term evaluation, for the first time
297 in the literature, of a well-screened population of women subjected to PBS in the
298 absence of tubal pathologies, even when in their late reproductive years. All the
299 currently known confounding factors that can interfere on ovarian reserve were listed
300 among the exclusion criteria for both the first (9) and the current study. Reliable and
301 definitive information on the safety of PBS is critically important for the entire medical
302 community, considering that the idea that salpingectomy should be proposed as a
303 preventive strategy for low-risk women who undergo gynecologic surgery is gaining
304 increasing popularity. Moreover, although risk-reducing salpingo-oophorectomy
305 (RRSO) at around age 40 years is currently recommended to women who carry the
306 BRCA1/2 mutation, an alternative preventive strategy has been put forward for high-
307 risk women: early PBS and delayed oophorectomy (RRO). Although preventive
308 RRSO decreases the ovarian cancer risk by 80% to 96%, based on its short-term
309 and long-term morbidity, which potentially affect quality of life, and on recent insights
310 into the fallopian tube as the possible site of origin of serous ovarian carcinomas,
311 early PBS and delayed RRO could be the best solution.

312 Recently, a multi-center non-randomized trial has started enrollment, and
313 participants will choose between standard RRSO at age 35 to 40 years (BRCA1) or
314 40 to 45 years (BRCA2) and the alternative strategy of PBS on completion of
315 childbearing and RRO at age 40 to 45 years (BRCA1) or 45 to 50 years (BRCA2).
316 The aim of the study is to measure menopause-related quality of life, but also
317 ovarian/breast cancer incidence, surgery-related morbidity, histopathology,
318 cardiovascular risk factors and diseases, and cost-effectiveness (25).

319 The increasing interest in PBS as a preventive strategy for both low- and high-
320 risk women and the related crucial importance of long-term follow-up data are

321 confirmed by many web-based surveys of health professionals' acceptability/attitude
322 toward the new proposal (26-29).

323 A limitation of the study is the relatively small sample size of women
324 evaluated. However, given the homogeneous distribution of results among this
325 population, and considering that this is currently the largest sample of women
326 studied in the literature, the results presented deserve attention. Another limitation of
327 the current study is that the control population is not the same used in the original
328 study (9), that being composed of women treated by standard TLH (with adnexal
329 preservation) between September 2008 and September 2010, matched for uterine
330 weight. Unfortunately, none of these women had undergone an OvAge test at that
331 time, and the gap of average age of at least 2 years makes it impossible to
332 accurately compare that group with the PBS population. Age at menopause of these
333 women will be an interesting outcome for forthcoming analysis, despite the need of
334 consensus for a correct definition for women without a uterus. In this context, further
335 research on the validation of the OvAge model to accurately predict the time to
336 menopause is already in progress.

337 Two other possible limits of our study are: first, the range of time
338 postoperatively at which the subjects have being evaluated creates a variable in
339 comparison of the data. However, according to our previous experience (30), when
340 there is evidence of impaired ovarian reserve caused by vascular damage, the
341 modification of biochemical and ultrasonographic parameters is already evident at 3
342 months follow-up, and it remains constant at 1 and 2 years. In the same way,
343 reassuring parameters at 3 months do not worsen at the next re-evaluation, allowing
344 us to speculate that the results of the current study are poorly affected by the range
345 of evaluation times.

346 Furthermore, at surgery and at the time of follow-up, our patients already had
347 evidence of diminished ovarian reserve, given their mean age of 45.97 ± 2.36 and
348 49.61 ± 2.15 , respectively. Although it would be better to assess the impact of surgery
349 on younger women, in which ovaries have greater potential for functional loss,
350 unfortunately any assessment of the impact of prophylactic salpingectomy on
351 ovarian reserve is not ethically conducive in patients who have not yet fulfilled their
352 reproductive desire. That is why, to date, in all studies conducted in patients
353 undergoing salpingectomy for cancer prophylaxis, in the absence of tubal pathology
354 and not for reproductive intent, the average age of the population has always been
355 more than 37 years (10,11,17). Ideally, the same rigorous approach used for women
356 subjected to PBS should be applied on all future studies aimed at evaluating the
357 effect of this kind of surgery in women undergoing IVF, to extend the assessable
358 population and to definitively conclude the mid- and long-term safety of laparoscopic
359 salpingectomy.

360 In conclusion, according to our results, the addition of PBH during TLH in the
361 late reproductive years does not cause ovarian function impairment up to 3 to 5
362 years after surgery.

363

364 **Table legends**365 Table 1. **Descriptive statistics of enrolled women**

366 Table 2. **Linear models applied.** Two models are summarized. In model 1, an
367 ANCOVA with interaction (maximal) model is applied. A simpler model, the minimally
368 adequate one, neglects any role of the treatment/control covariate, with a very
369 similar residual standard error, confirming the hypothesis that the addition of PBS to
370 TLH had no detrimental effect on ovarian function.

371

372 **Figure Legend**

373 **Figure 1: Two-dimensional plot for regression analysis.** The OvAge control
374 group (gray and white bullets) and the TLH plus PBS treated group (black and dark
375 gray diamonds). In the panel, three regression lines have been drawn: a 'two-
376 parameters' thin dashed line, with intercept and slope specific for the control group; a
377 'two-parameters' thin continuous line, with intercept and slope specific for the treated
378 group; and, a 'one-parameter' solid dashed line, with common slope (and null
379 intercept) for both groups. All three regression lines do not differ in a statistical
380 sense; consequently, the 'one-parameter common line' is the minimal adequate
381 model to retain as a valid description of both groups.

382 **References**

- 383 1. Crum CP, Drapkin R, Miron A, Ince TA, Muto M, Kindelberger DW et al.
384 The distal fallopian tube: a new model for pelvic serous carcinogenesis.
385 *Curr Opin Obstet Gynecol* 2007;19:3-9.
- 386 2. ACOG Committee opinion no. 620. Salpingectomy for ovarian cancer
387 prevention. *Obstet Gynecol* 2015;125:279-81.
- 388 3. Walker JL, Powell CB, Chen LM, Carter J, Bae Jump VL, Parker LP et al.
389 Society of Gynecologic Oncology recommendations for the prevention of
390 ovarian cancer. *Cancer* 2015. DOI: 10.1002/cncr.29321. [Epub ahead of
391 print]
- 392 4. Parker WH, Broder MS, Chang E, Feskanich D, Farquhar C, Liu Z et al.
393 Ovarian conservation at the time of hysterectomy and long-term health
394 outcomes in the nurses' health study. *Obstet Gynecol* 2009;113:1027–37.
- 395 5. Practice Committee of American Society for Reproductive Medicine in
396 collaboration with Society of Reproductive Surgeons. Salpingectomy for
397 hydrosalpinx prior to in vitro fertilization. *Fertil Steril* 2008;90(5 Suppl):S66-
398 8.
- 399 6. Practice Committee of the American Society for Reproductive Medicine.
400 Role of tubal surgery in the era of assisted reproductive technology: a
401 committee opinion. *Fertil Steril* 2015;103:37-43.
- 402 7. Tsiami A, Chaimani A, Mavridis D, Siskou M, Assimakopoulos E, Sotiriadis
403 A. Surgical treatment for hydrosalpinx prior to IVF-ET: a network meta-
404 analysis. *Ultrasound Obstet Gynecol* 2016. DOI: 10.1002/uog.15900.
405 [Epub ahead of print]

- 406 8. Fan M, Ma L. Effect of salpingectomy on ovarian response to
407 hyperstimulation during in vitro fertilization: a meta-analysis. *Fertil Steril*
408 2016. DOI: 10.1016/j.fertnstert.2016.03.053. [Epub ahead of print]
- 409 9. Morelli M, Venturella R, Mocciaro R, Di Cello A, Rania E, Lico D et al.
410 Prophylactic salpingectomy in premenopausal low-risk women
411 for ovarian cancer: primum non nocere. *Gynecol Oncol* 2013;129:448-51.
- 412 10. Findley AD, Siedhoff MT, Hobbs KA, Steege JF, Carey ET, McCall CA et
413 al. Short-term effects of salpingectomy during laparoscopic hysterectomy
414 on ovarian reserve: a pilot randomized controlled trial. *Fertil Steril*
415 2013;100:1704-8.
- 416 11. Venturella R, Morelli M, Lico D, Di Cello A, Rocca M, Sacchinelli A et al.
417 Wide excision of soft tissues adjacent to the ovary and fallopian tube does
418 not impair the ovarian reserve in women undergoing prophylactic bilateral
419 salpingectomy: results from a randomized, controlled trial. *Fertil*
420 *Steril* 2015;104:1332-9.
- 421 12. Venturella R, Lico D, Sarica A, Falbo MP, Gulletta E, Morelli M et al.
422 OvAge: a new methodology to quantify ovarian reserve combining clinical,
423 biochemical and 3D-ultrasonographic parameters. *J Ovarian*
424 *Res* 2015;8:8-21.
- 425 13. Falbo A, Orio F, Venturella R, Rania E, Materazzo C, Tolino A et al. Does
426 metformin affect ovarian morphology in patients with polycystic ovary
427 syndrome? A retrospective cross-sectional preliminary analysis. *J Ovarian*
428 *Res* 2009;2:5.
- 429 14. Crawley, Michael J, *Statistics: an introduction using R*, 2005, John Wiley &
430 Sons;2005.

- 431 15. R Core Team. *R: A Language and Environment for Statistical Computing*.
432 R Foundation for Statistical Computing, Vienna, Austria, 2014.
- 433 16. Dietl J and Wischhusen J. The postreproductive salpingectomy. *Fertil*
434 *Steril* 2014;101.
- 435 17. Sezik M, Ozkaya O, Demir F, Sezik HT, Kaya H. Total salpingectomy
436 during abdominal hysterectomy: effects on ovarian reserve and ovarian
437 stromal blood flow. *J Obstet Gynaecol Res* 2007;33:863-9.
- 438 18. Zhang Y, Sun Y, Guo Y, Li TC, Duan H. Salpingectomy and Proximal
439 Tubal Occlusion for Hydrosalpinx Prior to In Vitro Fertilization: A Meta-
440 analysis of Randomized Controlled Trials. *Obstet Gynecol Surv*
441 2015;70:33-38.
- 442 19. Strandell A, Lindhard A, Waldenström U, Thorburn J. Prophylactic
443 salpingectomy does not impair the ovarian response in IVF treatment.
444 *Hum Reprod* 2001;16:1135-9.
- 445 20. Chan CC, Ng EH, Li CF, Ho PC. Impaired ovarian blood flow and reduced
446 antral follicle count following laparoscopic salpingectomy for ectopic
447 pregnancy. *Hum Reprod* 2003;18:2175-80.
- 448 21. Ye XP, Yang YZ, Sun XX. A retrospective analysis of the effect of
449 salpingectomy on serum antiMüllerian hormone level and ovarian reserve.
450 *Am J Obstet Gynecol* 2015;212:53.e1-10.
- 451 22. Oliver Perez MR, Magriñá J, García AT, Jiménez Lopez JS. Prophylactic
452 salpingectomy and prophylactic salpingoophorectomy for adnexal high-
453 grade serous epithelial carcinoma: A reappraisal. *Surg Oncol* 2015;24:335-
454 44.

- 455 23. Johnson N, van Voorst S, Sowter MC, Strandell A, Mol BW. Surgical
456 treatment for tubal disease in women due to undergo in vitro fertilisation.
457 *Cochrane Database Syst Rev* 2010(1):CD002125
- 458 24. Zhang Y, Sun Y, Guo Y, Li TC, Duan H. Salpingectomy and proximal tubal
459 occlusion for hydrosalpinx prior to in vitro fertilization: a meta-analysis of
460 randomized controlled trials. *Obstet Gynecol Surv* 2015;70:33-8.
- 461 25. Harmsen MG, Arts-de Jong M, Hoogerbrugge N, Maas AH, Prins JB,
462 Bulten J et al. Early salpingectomy (TUbectomy) with delayed
463 oophorectomy to improve quality of life as alternative for risk-reducing
464 salpingo-oophorectomy in BRCA1/2 mutation carriers (TUBA study): a
465 prospective non-randomised multicentre study. *BMC Cancer* 2015;15:593.
- 466 26. Chandrasekaran D, Menon U, Evans G, Crawford R, Saridogan E, Jacobs
467 C et al. Risk reducing salpingectomy and delayed oophorectomy in high
468 risk women: views of cancer geneticists, genetic counsellors and
469 gynaecological oncologists in the UK. *Fam Cancer* 2015;14:521-30.
- 470 27. Venturella R, Rocca M, Lico D, Trapasso S, Di Cello A, Gizzo S et al.
471 Prophylactic bilateral salpingectomy for the prevention of ovarian cancers:
472 what is happening in Italy? *Eur J Canc Prev* 2015 [Epub ahead of print]
- 473 28. Reade CJ, Finlayson S, McAlpine J, Tone AA, Fung-Kee-Fung M,
474 Ferguson SE. Risk-reducing salpingectomy in Canada: a survey of
475 obstetrician-Gynaecologists. *J Obstet Gynaecol Can* 2013;35:627-34.
- 476 29. Gill SE, Mills BB. Physician opinions regarding elective bilateral
477 salpingectomy with hysterectomy and for sterilization. *J Minim Invasive*
478 *Gynecol* 2013;20:517-521.

- 479 30. Morelli M, Mocciano R, Venturella R, Imperatore A, Lico D, Zullo F. Mesial
480 side ovarian incision for laparoscopic dermoid cystectomy: a safe and
481 ovarian tissue-preserving technique. Fertil Steril 2012;98:1336-40.


ACCEPTED MANUSCRIPT

Table 1. Descriptive statistics of enrolled women

Parameters	Mean values \pm SD
Age at surgery (years)	45.85 \pm 2.40
Age at follow-up (years)	49.61 \pm 2.15
OvAge at follow-up (years)	49.34 \pm 2.12
FSH at follow-up (mU/mL)	43.02 \pm 19.92
AMH at follow-up (ng/mL)	0.12 \pm 0.20
3D AFC at follow-up (n.)	1.91 \pm 1.28
VI at follow-up (%)	2.80 \pm 5.32
FI at follow-up (1-100)	19.37 \pm 5.88
VFI at follow-up (1-100)	0.56 \pm 1.12

Table 2. **Linear models applied.** Two models are summarized. In model 1, an Ancova with interaction (maximal) model is approached. A simpler model, the minimal adequate one, neglect any role of the Treatment/control covariate, with a very similar residual standard error, confirming the hypothesis that the addition of PBS to TLH did not have any detrimental effect on ovarian function.

Maximal model			
	Estimate	Standard Error	p-value
Intercept	0.1005	0.8474	.906
OvAge	1.0028	0.0225	< .001
Treatment	1.5632	11.2572	.890
OvAge : Treatment	0.0286	0.2282	.900
Minimal adequate model			
	Estimate	Standard Error	p-value
OvAge	1.0008	0.0040	< .001



Precis:

The addition of bilateral salpingectomy during hysterectomy is a safe procedure and does not cause ovarian reserve impairment, even three to five years after surgery.

ACCEPTED MANUSCRIPT