



Long-term toxicity of surface-charged polystyrene nanoplastics to marine planktonic species *Dunaliella tertiolecta* and *Artemia franciscana*

This is a pre print version of the following article:

Original:

Bergami, E., Pugnali, S., Vannuccini, M.L., Manfra, L., Faleri, C., Savorelli, F., et al. (2017). Long-term toxicity of surface-charged polystyrene nanoplastics to marine planktonic species *Dunaliella tertiolecta* and *Artemia franciscana*. *AQUATIC TOXICOLOGY*, 189, 159-169 [10.1016/j.aquatox.2017.06.008].

Availability:

This version is available <http://hdl.handle.net/11365/1031321> since 2018-02-06T11:09:59Z

Published:

DOI:10.1016/j.aquatox.2017.06.008

Terms of use:

Open Access

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.

(Article begins on next page)

1 **Long-term toxicity of surface charged polystyrene nanoplastics to marine**
2 **planktonic species *Dunaliella tertiolecta* and *Artemia franciscana***

3
4 Bergami E.^{1*}, Pugnolini S.¹, Vannuccini M. L.¹, Manfra L.^{2,3}, Faleri C.⁴, [Savorelli F.⁵](#), Dawson K.
5 A.⁵⁶, Corsi I.¹

6 ¹Department of Physical, Earth and Environmental Sciences, University of Siena, Italy;

7 ²Institute for Environmental Protection and Research (ISPRA), Rome, Italy;

8 ³Department of Biology and Evolution of Marine Organisms, Anton Dohrn Zoological Station,
9 80121 Naples, Italy;

10 ⁴Department of Life Sciences, University of Siena, 53100 Siena, Italy;

11 ⁵[Regional Agency for Prevention, Environment and Energy of Emilia-Romagna \(Arpae ER\),](#)
12 [Ferrara, Italy](#)

13 ⁶Centre for BioNano Interactions, School of Chemistry and Chemical Biology, University College
14 Dublin, Ireland;

15

16

17

18

19

20

21 ***Corresponding author:** Elisa Bergami; bergami@student.unisi.it; Department of Physical, Earth
22 and Environmental Sciences, University of Siena, Italy.

1 Abstract

2 Plastic pollution has been globally recognized as a critical issue for marine ecosystems and
3 nanoplastics constitute one the last ~~frontier-unexplored areas~~ to understand the magnitude of this
4 threat. However, current difficulties in sampling and identifying nano-sized debris make hard to
5 assess their occurrence in marine environment. Polystyrene nanoparticles (PS NPs) ~~have-arebeen~~
6 ~~recently-adopted~~ largely used as ~~model-for~~ nanoplastics in ecotoxicological studies and ~~despite~~
7 although acute exposures have been already investigated, long-term toxicity on marine organisms is
8 lacking. Our study aims at evaluating the effects of 40 nm PS anionic carboxylated (PS-COOH) and
9 50 nm cationic amino-modified (PS-NH₂) NPs in two planktonic species, the green microalga
10 *Dunaliella tertiolecta* and the brine shrimp *Artemia franciscana*, respectively prey and predator. PS
11 NP behaviour in exposure media was determined through DLS, while their toxicity to microalgae
12 and brine shrimps s evaluated through 72 h growth inhibition test and 14 ~~days~~ long-term toxicity test
13 respectively. Moreover, the expression of target genes (i.e. *clap* and *cstb*), having a role in brine
14 shrimp larval growth and molting, was measured in 48 h brine shrimp larvae. A d~~D~~ifferent
15 behaviour ~~-of the two PS NPs~~ in exposure media as well as diverse toxicity ~~was-observed-toin~~ the
16 two planktonic species ~~-was-observedupon-PS-NP-exposure~~. PS-COOH formed micro-scale
17 ~~aggregatesobjects~~ (Z-Average > 1 µm) and did not affect the growth of microalgae up to 50 µg/ml
18 as well as brine shrimps up to 10 µg/ml. However, these negatively charged NPs were adsorbed on
19 microalgae and accumulated (and excreted) in brine shrimps, suggesting a potential trophic transfer
20 from prey to predator. On the opposite, PS-NH₂ formed nano-scale aggregates (Z-Average < 200
21 nm), ~~and~~ caused inhibition of algal growth (EC₅₀ = 12.97 µg/ml) and mortality in brine shrimps at
22 14 d (LC₅₀ = 0.83 µg/ml). Moreover, 1 µg/ml PS-NH₂ ~~(at-1 µg/ml)~~ significantly induced *clap* and
23 *cstb* genes, explaining the physiological alterations (e.g. increase in molting) previously observed in
24 48 h larvae, but also suggesting an apoptotic pathway triggered by cathepsin L-like protease in brine
25 shrimps upon PS-NH₂ exposure. These findings provide a first insight into long-term toxicity of

1 nanoplastics to marine plankton, underlining the role of the surface chemistry in determining the
2 behaviour and effects of PS NPs, in terms of adsorption, growth inhibition, accumulation, gene
3 modulation and mortality. The use of long-term end-point has been identified as ~~more~~-valuable tool
4 for assessing the impact of nanoplastics on marine planktonic species, being more predictable of
5 real exposure scenarios for risk assessment purposes.

6

7

8

9

10

11

12

13

14

15

16

17 **Keywords:** ~~nanoplastics~~, surface charge~~s~~, polystyrene nanoparticles, ecotoxicity, marine plankton,
18 growth inhibition*Artemia franciscana*, *Dunaliella* cathepsin L-like protease*tertiolecta*.

19

1. Introduction

Plastic debris have been globally recognized as a menace for marine ecosystems (Andrady, 2011), being the major portion (between 60 and 80% and up to 96.87%) of the marine litter found during monitoring surveys (Gregory and Ryan, 1997; Ruiz-Orejón et al., 2016).

The evaluation of the impacts of the smallest ~~invisible~~ fraction of plastic debris, defined as nanoplastics (< 1 µm) (Hartmann et al., 2015), constitutes one of the last unexplored areas~~the last frontier in order~~ to fully understand the importance of this emerging threat for the marine environment.

The amount of nanoplastics spread in the oceans is currently unexplored, since ~~convectional~~ conventional sampling methods (i.e. neuston nets having mesh size > 300 µm) as well as analytical techniques available for the identification of plastic polymers still prevent to isolate and quantify the nano-fraction (Cózar et al., 2014; Koelmans et al., 2015).

Notwithstanding, Zhang and co-authors (2012) were able to measure polystyrene nanoparticles (PS NPs) in the range of 22 – 220 nm; from the thermal cutting of polystyrene foam and later Lambert and Wagner (2016) found PS NPs in water suspensions obtained from weathered bulk polystyrene material under controlled laboratory conditions. This latter study demonstrated the occurrence of nanofragmentation of plastic debris in the aquatic environment as a consequence of several processes: UV-radiation, thermo-oxidation, hydrolysis, mechanical abrasion and not least biological degradation (Andrady, 2011). Such weathering processes may confer to the nano-sized fragments ~~new peculiar~~ novel properties (Andrady, 2017), as for engineered nanomaterials (Klaine et al., 2012). For example, weathered plastic debris can be characterised by an increase in crystallinity due to oxidation (Rouillon et al., 2016) and acquire carbonyl functionalities as well as negative surface charges (Fotopoulou and Karapanagioti, 2012).

1 Nano-sized plastics can ~~thus~~ be easily ingested by organisms (Cole et al., 2013), pass biological
2 barriers (Moore, 2006), penetrate tissues (Kashiwada, 2006) and even bioaccumulate in organs and
3 tissues (von Moos et al., 2012). ~~Moreover,~~ T, their high surface area can lead to exceptionally strong
4 sorption of toxic compounds (e.g. PCBs, PAH and DDTs) (Rios et al., 2007; Rochman et al., 2013;
5 Velzeboer et al., 2014), with potential added chemical toxicity, once NPs have passed the cell
6 membranes (Bexiga et al., 2011; Salvati et al., 2011). Any adverse effect of nanoplastics will be
7 thus related to their nano-scale properties, to polymer-associated chemicals (i.e. additives) or both,
8 and will mainly depend on particle size, polymer type and aging (Besseling et al., 2014). Therefore,
9 it is necessary to understand how nanoplastics behave in the marine environment depending on the
10 chemical nature of the polymer and its surface modifications. Since the marine environment could
11 be seen as the last reservoir of plastic debris, phyto- and zooplanktonic organisms can be seriously
12 affected by the presence ~~are among the primary biological targets~~ of nanoplastics, being exposed to
13 a variety of low-density polymeric ~~beads-particles~~ along the ~~whole~~ water column (Manzo et al.,
14 2013; Matranga and Corsi, 2012; Moore, 2006).

15 In order to evaluate the impact of nanoplastics on the wildlife, polymeric nano-scale particles, such
16 as PS NPs having a polystyrene core and variable surface functional groups, can be used. These
17 NPs are synthesized for a wide range of applications (Nowak and Bucheli, 2007), including
18 biosensors (Velev and Kaler, 1999), photonics (Rogach et al., 2000), nanocomposites (Merinska
19 and Dujkova, 2012) and drug delivery tools (Popielarski et al., 2005; Yap and Zhang, 2007).
20 Common PS NPs include anionic carboxylated (-COOH) and cationic amino (-NH₂) surface
21 modifications (Loos et al., 2014), which allow them to pass more easily through the cell membrane,
22 as they share a similar molecular structure to proteins (Rossi et al., 2014). Several studies on cell
23 lines confirmed their cellular uptake (Johnston et al., 2010; Salvati et al., 2011; Wang et al., 2012;
24 Loos et al., 2014) as well as toxicity (Bexiga et al., 2011; Frölich et al., 2012). In particular, positive
25 PS-NH₂ NPs were shown to trigger specific in vitro toxicity mechanisms ~~in vitro~~ including

1 lysosomal damage and ROS generation followed by apoptotic pathways through the induction of
2 cathepsins and caspases (Wang et al., 2013).

3 Ecotoxicity studies about nanoplastics impact on aquatic organisms have exponentially increased in
4 the last seven years and the majority adopted both PS-COOH and PS-NH₂ as model NPs
5 (Bhattacharya et al., 2010, Wegner et al., 2012; Casado et al., 2013; Besseling et al., 2014, Della
6 Torre et al., 2014, Cole and Galloway, 2015; Bergami et al., 2016; Canesi et al., 2015, Canesi et al.,
7 2016; Pinsino et al., 2017).

8 Our previous findings on zooplankton (Della Torre et al., 2014; Bergami et al., 2016; Pinsino et al.,
9 2017) showed diverse effects on sea urchin embryonic development (*Paracentrotus lividus*) and
10 brine shrimp larval growth (*Artemia franciscana*) depending on the surface charge of PS NPs.
11 Indeed, potential toxicity associated to the PS core could be neglected, since it does not degrade
12 under environmental conditions even in long-term studies ([Besseling et al., 2014](#); Loos et al., 2014).

13 After short-term exposures (48 h), [negatively charged](#) ~~negative~~ PS-COOH aggregates were strongly
14 retained in the digestive tract of sea urchin embryos and brine shrimp larvae, while positively
15 charged NPs seriously affected the development and growth of these species. In sea urchin
16 embryos, PS-NH₂ were able to elicit developmental defects through the induction of genes related
17 to stress (i.e. *hsp70*) but also apoptosis (*cas8*), similarly to previous findings in human cell lines
18 (Wang et al., 2013).

19 In brine shrimp larvae, PS-NH₂ were mostly stuck to the external appendices, clearly impairing
20 their swimming and increasing the molting of about 50% respect to the control (Bergami et al.,
21 2016). However, whereas EC₅₀ values were determined for sea urchin embryos exposed to PS-NH₂
22 (Della Torre et al., 2014), for brine shrimp larvae it was not possible to discriminate the toxicity of
23 PS NPs through the standard acute test (Artoxkit, 2014) and the mechanism behind the
24 physiological alterations observed was not fully elucidated.

1 In ecotoxicology, the acute toxicity is determined at first and mainly in terms of mortality, with
2 model organisms exposed to high concentrations of contaminants over a short time period.
3 However, any result based only on short-term and high exposure concentrations hamper
4 extrapolation of data to a more realistic scenario (Rand, 1995). On the contrary, long-term toxicity
5 tests follow as needed, usually lasting for 10% of the organism's lifespan (Newman, 2010) and
6 focus on both mortality and sub-lethal effects (e.g. growth and reproduction) (Cattaneo et al., 2009).
7 Such potential alterations, as a result of a long-term exposure, constitute a the-ultimate more reliable
8 endpoint (Rand, 1995) and represent the most appropriate tool for studying emerging contaminants
9 (Comfort et al., 2014) including nanoplastics. For example, Besseling and co-authors (2014)
10 observed reduced body size and alteration in the reproduction of *Daphnia magna* exposed for 21
11 days to 70 nm PS-COOH.

12 Considering the brine shrimp of the genus *Artemia* as model marine organisms, Rotini et al. (2015)
13 recently compared the EC₅₀ values obtained from different ~~ecotoxicity tests~~ assays, showing the
14 reliability of the long-term toxicity test compared to hatching and acute tests.

15 Our study aims at evaluating the effects caused by two different surface charged nanoplastics (40
16 nm ~~PS anionic carboxylated~~ (PS-COOH) and 50 nm ~~cationic amino-modified~~ (PS-NH₂) NPs on two
17 planktonic species, the green microalga *Dunaliella tertiolecta* and the brine shrimp *Artemia*
18 *franciscana*, through 72 h growth inhibition test and 14 d long-term toxicity test respectively. PS
19 NP disposition as well as their consequences to marine plankton (i.e. surface adsorption,
20 accumulation, growth inhibition and mortality) were assessed. Moreover, the expression of *clap* and
21 *cstb* genes related to larval growth and molting was determined in 48 h brine shrimp larvae exposed
22 to PS-NH₂.

23 These model organisms represent the first levels at the bottom of the marine trophic web,
24 respectively as prey and predator, thus any negative effect on them leads to serious repercussions on
25 the health of the marine ecosystem. Both negatively (-COOH) and positively (-NH₂) surface

1 charged PS NPs were used in order to correlate the different functionalization (i.e. surface
2 chemistry) to the observed toxicity.

3

4 **2. Materials and Methods**

5 **2.1. PS NP behaviour in exposure media**

6 40 nm yellow-green fluorescently labelled PS-COOH (505 nm excitation, 515 nm emission) were
7 purchased from Invitrogen, whereas 50 nm unlabelled PS-NH₂ NPs from Bangs Laboratories Inc.
8 Functionalized PS NPs (anionic PS-COOH and cationic PS-NH₂) are the most common
9 nanoplastics used in ecotoxicological studies on aquatic organisms published in the last 7 years
10 (Bhattacharya et al., 2010, Besseling et al., 2014, Della Torre et al., 2014, Cole and Galloway,
11 2015, Canesi et al., 2015, Canesi et al., 2016, Pinsino et al., 2017). In this study, the use of the same
12 batches for PS NPs allowed us to compare the obtained results with our previous findings on brine
13 shrimp ~~as model organism~~ (Bergami et al., 2016).

14 PS NP stock solutions were supplied in deionised milli-Q water without any surfactants or
15 preservatives (mQW) used for ecotoxicity tests and contained 50 and 100 mg/ml of PS-COOH and
16 PS-NH₂ respectively. Concerning the fluorescent labelling of PS-COOH, the dye was contained
17 inside the polymer matrix instead of being attached on the surface. Therefore, any potential toxicity
18 related to the dye was considered negligible.

19 Primary characterization of PS NPs was performed as reported in Bergami et al. (2016). Secondary
20 characterization of PS NPs in milli-Q water (mQW), microalgae medium (prepared in 0.45 µm
21 filtered NSW, T = 20°C, salinity 30‰, pH 8.3) and natural sea water (NSW) (0.45 µm filtered, T =
22 25°C, salinity 38‰, pH 8.3) was made using Dynamic Light Scattering (DLS, Malvern
23 instruments), combined with the Zetasizer Nano Series software, version 7.02 (Particular Sciences,

1 UK). Z-average (nm), Polydispersity Index (PDI, dimensionless) and Zeta (ζ -) potential (mV) were
2 measured as key parameters describing NP behaviour in complex environmental media (SCENIHR,
3 2007; Stone et al., 2010). Measurements were carried out in triplicate, each containing 11 runs of 10
4 second for size parameters, 20 runs for ζ -potential.

5 Ecotoxicity tests were performed in NSW-based media collected from the Tuscany archipelago area
6 (Tyrrhenian Sea) (Bergami et al., 2016). Physico-chemical parameters of NSW included: TOC
7 1.3%, total oxygen 6.6 mg/l, total PAH 0.12 mg/Kg, Cr < 1 μ g/l, As 1 μ g/l, Cd 0.09 μ g/l, Hg 0.02
8 μ g/l, Pb < 1 μ g/l (data available at: SIRA RSS www.sira.arpat.toscana.it/). PS NP final suspensions
9 in NSW were prepared from the stock solutions and quickly vortexed without sonication prior to
10 use, since no important changes in PS NP size measurements in NSW with and without sonication
11 have been previously observed (see Della Torre et al., 2014 SI). This method of dispersion can be
12 seen as a more realistic representation of natural conditions occurring for nanoplastics in the marine
13 environment (Bhattacharya et al., 2010).

14 PS NP concentrations used are reported as μ g/ml (for particle numbers, refer to Table S1 in
15 Supporting Information).

16

17 **2.2. Ecotoxicity**

18 In this study, the toxicity of PS nanoplastics was evaluated on two planktonic species, ~~as~~ the
19 unicellular green microalga *D. tertiolecta*, supplied by the Regional Agency for Environmental
20 Protection (ARPA Emilia-Romagna), and the microcrustacean brine shrimp *A. franciscana* ~~larvae~~,
21 purchased from MicroBioTests (Ghent, Belgium). Growth inhibition test (72 h) and long-term
22 lethal test (14 d) were conducted, respectively, on the green microalga and the microcrustacean by
23 testing both PS NPs following standardized procedures, as described below. Microalgae were

1 further used to feed brine shrimp larvae during long-term lethal toxicity test, ~~using a prey to~~
2 ~~predator approach.~~

3

4 *2.2.1. Growth Inhibition of the green microalga*

5 The growth inhibition test (72 h) using green microalgae was performed according to a standard
6 guideline (ISO, 2006) with some modifications as follow. Sterile microalgae medium was prepared
7 in NSW (salinity adjusted to 30‰) by the addition of f/2 Guillard and then autoclaved. Experiments
8 were performed in 24-well plates, with average initial cell density of $7 \cdot 10^5$ cells/ml. Tested PS NPs
9 were 0.5 - 1 - 5 - 10 - 25 - 50 $\mu\text{g/ml}$ and microalgae medium was used as control. For each group,
10 three replicates were run and each experiment repeated at least three times for each PS NP. During
11 the test, microalgae were incubated under controlled static conditions ($20 \pm 1^\circ\text{C}$, photoperiod of 16:8
12 h light:darkness, without shaking). After 72 h, microalgae were fixed in 1:1 lugol:ethanol solution
13 and cell density estimated by counting under optical microscope Olympus BX51 (40X), equipped
14 with a Neubauer Improved chamber. The number of cells/ml, growth rate (μ) and inhibition of
15 growth rate (I_{μ}) were determined. EC_{50} were calculated by plotting the I_{μ} against the logarithm of
16 the concentration of PS NP tested.

17

18 *2.2.2. Short- and long-term toxicity on the brine shrimp*

19 In order to provide new insight into the expression to genes related to physiological alterations
20 specifically provoked by PS-NH₂ in brine shrimp larvae after 48 h (Bergami et al., 2016), Hatching
21 ~~of brine shrimp cysts and~~ short-term toxicity test ~~were was~~ performed ~~as previously described in~~
22 Bergami et al. (2016) accordingly, with some modifications as reported. Briefly, about 200 newly
23 hatched nauplii (Instar I nauplius stage) were exposed to PS-NH₂ suspensions (at 0, 0.1 and 1
24 $\mu\text{g/ml}$) prepared in NSW in glass beakers (final volume of 100 ml) and incubated at $25 \pm 1^\circ\text{C}$ ~~under~~

1 dark static conditions, without providing food. At 48 h, all swimming larvae were collected and
2 stored at -80°C for gene expression analysis. The experiment was performed in triplicates and
3 repeated three times.

4 ~~The long-term toxicity of both PS NPs (PS-COOH and PS-NH₂) test~~ was then carried out
5 according to Savorelli et al. (2007) and Manfra et al. (2012), with some changes concerning the
6 preparation of the newly hatched larvae obtained according the protocol available from Artoxkit
7 (2014), since we decided to maintain the same conditions as Bergami et al. (2016), for hatching and
8 start the long-term toxicity test using brine shrimp larvae at Instar I nauplius stage.

9 Preliminarily, PS NP concentrations tested for evaluating the long-term toxicity were 1 and 10
10 µg/ml. The former one, corresponding to about 1.5 – 2.8 · 10¹⁰ NPs/ml (Table S1), ~~could be~~
11 ~~representative of a putative environmental concentration of nanoplastics in the marine environment~~
12 ~~(Lambert and Wagner, 2016)~~, while the second ~~concentration one~~ was in the same order of
13 magnitude of the EC₅₀ calculated for PS-NH₂ in the growth inhibition test (see paragraph 3.2. in the
14 results section). However, due to the high mortality observed for brine shrimps exposed to PS-NH₂
15 in the first attempts, a further concentration series for this positively charged nanoplastic was set to
16 0.5 - 1 - 1.5 - 2.5 - 5 µg/ml.

17 PS NP test solutions were prepared in 50 ml flasks (final volume of 30 ml) containing NSW, to
18 which a microalgae inoculum (to reach a final density of 10⁵ cells/ml) and 10 newly hatched brine
19 shrimp nauplii were added in sequence. For the whole duration of the test, the flasks were
20 maintained under controlled conditions (25±1°C, photoperiod of 16:8 h light:darkness) and test
21 solutions renewed every 2-3 days (at 2, 5, 7, 9 and 12 d). Three replicates were set for each PS NP
22 concentration and the control group (containing only NSW and microalgae) and the test was
23 repeated at least three times. As end-point, the number of dead larvae after 14 days of exposure was
24 recorded on a worksheet and the percentage of mortality at each concentration calculated. Brine
25 shrimps were considered dead if they did not display any movement for 10 seconds (Gambardella et

1 al., 2014). The test was considered acceptable if the control group displayed an average mortality \leq
2 20 %. LC₅₀ value and 95% confidence limits were calculated by comparison of the mortality at each
3 concentration to the control group, using EPA Probit Analysis Program (version 1.5).

4 In order to further investigate the effect of PS-NH₂ on larval development, images of brine shrimps
5 from the 14 days toxicity test were taken through optical microscopy (Olympus BX51) and
6 analysed using ImageJ software (Version 1.49, Wayne Rasband, National Institutes of Health,
7 USA). For each organism, the length from the cephalic region (i.e. nauplius eye) to the telson was
8 measured. Antennule and caudal rami were not considered in the measurements, since these parts
9 were usually present at different focus distances with respect to the main body (i.e. thorax and
10 abdomen) of the animal.

11

12 **2.3. Disposition of PS nanoplastics in planktonic organisms**

13 Disposition of fluorescent PS-COOH (at 5 $\mu\text{g/ml}$) was assessed in 72 h exposed green microalgae,
14 washed twice in NSW and immobilized on slides pre-treated with 0.05% protamine sulfate and
15 nuclei labelled with DAPI. As negative control, microalgae ~~in~~-suspensions without NPs were also
16 considered. Cells were observed under optical fluorescent microscope AXIO IMAGER Z1 using
17 Apotome system (Zeiss), with filter FITC 470/525 for PS-COOH and filter DAPI 365/445 for
18 nuclei staining. Images were taken with AxioCam MRm camera at 63X using Axio Vision
19 Software.

20 Disposition of unlabelled PS-NH₂ (at 5 $\mu\text{g/ml}$) was determined on microalgae surface using
21 scanning electron microscopy (SEM), according to the method described by Miller et al. (2012)
22 with some modifications. Briefly, cells were centrifuged (20°C, 1600 rpm, 3') and pellets fixed in
23 5% glutaraldehyde for 1 h, dehydrated in a graded series of ethanol up to 70% and maintained at
24 4°C until further analysis. As negative control, microalgae ~~in~~-suspensions without NPs were

1 considered. For the observation at SEM, samples were placed on a glass coverslip coated with a thin
2 layer of poly-L-lysine dehydrated in absolute ethanol and subsequently subjected to the critical point
3 drying. At last, samples were metallized and observed under SEM (Philips XL20), operating at 100
4 KW accelerating voltage.

5 Concerning the brine shrimp *A. franciscana*, the disposition of fluorescent PS-COOH (at 10 µg/ml)
6 was determined under optical fluorescent microscope (Olympus BX51) at 14 days of long-term
7 toxicity tests.

8

9 **2.4. RNA extraction, cDNA synthesis and Real Time q-PCR**

10 Gene expression analysis was performed following the procedure reported in Vannuccini et al.
11 (2015). RNA was extracted using reagent RNA extraction buffer (Sigma, UK) in 2 mL centrifuge
12 tubes using a Tissue Lyser (Qiagen, Hilden, Germany) and precipitated by adding 0.5 vol. of
13 isopropyl alcohol and 0.5 vol. of RNA precipitation solution (1.2 M sodium chloride, 0.8 M
14 disodium citrate). Three independent RNA isolations were performed for each sample. RNA
15 concentrations were measured using a traycell spectrophotometer (Eppendorf) at 260 nm λ and
16 RNA quality confirmed on 1% agarose gel, showing discrete 18 S and 28 S ribosomal RNA bands.
17 Total RNA (200 ng) was transcribed to cDNA using qScript™ XLT One-Step RT-qPCR
18 ToughMix® (Quanta Biosciences) according to manufacturer's protocol.

19 The expression of selected genes related to molting in brine shrimp larvae, as *cathepsin L*
20 *associated protein (clap)* and *cystatin B inhibitor (cstb)* was investigated through quantitative Real
21 Time PCR (RT q-PCR), using *glyceraldehyde 3-phosphatase dehydrogenase (gapdh)* as
22 housekeeping gene (Chen et al., 2009). Primers of these genes are listed in Table S24,
23 Supplementary Information. RT q-PCR was performed using a Stratagene Mx 3000P thermal
24 cycler. Each amplification reaction contained PCR for each gene was performed in triplicate in a
25 total volume of 20 µL containing 1 µL cDNA, 100 nM of each primer and 10 µL of SsoAdvanced™

1 Universal SYBR® Green Supermix 2x (BIORAD, Biorad Laboratories, USA). The cycling
2 conditions were: 95°C for 2 min for polymerase activation, followed by 40 PCR cycles 10 s at
3 95°C; 20 s at 50-60°C, 5 s at 72°C. Cycle threshold (Ct) values corresponded to the number of
4 cycles at which the fluorescence emission monitored in real time exceeded the threshold limit.
5 Melting curve analysis and gel electrophoresis of selected samples were performed to confirm the
6 production of a single amplification in these reaction. Real-time efficiency of primer set (E) was
7 determined for each gene from the slopes given by MxPro™ QPCR software (Stratagen,USA),
8 apply the equation $E = 10^{(-1/\text{slope})}$. The calculated relative expression ratio of each gene was based on
9 the PCR efficiency (E) and Ct of sample compared with control, and expressed in comparison to the
10 reference genes.

11

12 **2.5. Data analysis**

13 All statistical analysis were performed using Graphpad Prism 5. For the ecotoxicity results,
14 normality was verified using Shapiro-Wilk test and homogeneity of variances by Bartlett's test, data
15 were conformed to the assumptions. ~~For ecotoxicity results,~~ The one-way variance analysis
16 (ANOVA) was performed to compare the various treatments and $p < 0.05$ was taken as significant
17 cut-off. Results of long-term toxicity tests are mean of at least three independent experiments. LC₅₀
18 values were calculated by fitting the percentage of alive larvae to a classical sigmoidal dose-
19 response model according to the equation: $y = b + (a - b) / 1 + 10^{(\text{Log LC}_{50} - x)}$ where y is response, b
20 response minimum, a response maximum, x the logarithm of effect concentration and LC₅₀ the
21 concentration of effect giving 50% of maximum effect. Each experiment has been performed 3 to 5
22 times.

23 Data obtained from gene expression analysis are expressed as mean \pm standard deviation (s.d.). To
24 compare the various treatments, ANOVA analysis was applied, using Bonferroni post-hoc and
25 taking $p < 0.05$ (*) and $p < 0.001$ (***) as significant cut-off.

26

1 **3. Results and Discussion**

2 The aim of the present study was to investigate the effects caused by short- and long-term exposure
3 to nanoplastics in two planktonic species at the base of the marine trophic web, the green microalga
4 *D. tertiolecta* and the brine shrimp *A. franciscana*, using a prey-to-predator approach. Both
5 negatively (-COOH) and positively (-NH₂) surface charged PS NPs were used as model for
6 nanoplastics and the different functionalization (i.e. surface charges) vs the observed toxicity was
7 discussed.

8 9 **3.1. Behaviour of PS NPs in exposure media**

10 Nominal sizes of 40 nm for PS-COOH and 50 nm for PS-NH₂ in mQW was confirmed by TEM
11 imaging as reported in our previous study (Bergami et al., 2016). DLS results (shown in Table 1)
12 confirmed an optimal dispersion and stability in mQW for both PS NPs, with a Z-Average of 54.47
13 ± 0.82 nm for PS-COOH and 56.48 ± 0.60 nm for PS-NH₂ as well as low PDI values (0.116 and
14 0.161 respectively). ζ -potential values in mQW confirmed the negative surface charge (-66 ± 1.1
15 mV) for PS-COOH and the positive one ($+53.1 \pm 1.21$ mV) for PS-NH₂, characteristic of PS NP
16 functionalization.

17 Secondary characterization in the two exposure media (i.e. microalgae medium and NSW)
18 confirmed the different behaviour of the two PS NPs according to their different surface
19 functionalization. Intensity-based distributions obtained by DLS show high aggregation occurring
20 for PS-COOH in both exposure media (Figure 1A), opposed to a better dispersion of PS-NH₂
21 having only few aggregates of large size (Figure 1B). Table 1 reports a comparison among PS NP
22 parameters in the two exposure media compared to mQW, adopted as reference medium for the
23 dispersion. Z-Average values indicate that PS-COOH originated micro-scale aggregates both in
24 microalgae medium and NSW (as 1237.5 ± 107 nm and 1064 ± 101 nm respectively), whereas PS-

1 NH₂ resulted far less aggregated, with an hydrodynamic size of 127 ± 5 nm in microalgae medium
2 and 196 ± 7 nm in NSW. Such acquired dimensions are congruent with PDI values (> 0.220) and
3 lower absolute values of ζ -potential indicating a broader size distribution of PS NPs and instability
4 in such exposure media, as compared to the ones in mQW.

5 These results confirmed our previous findings on the same batch of PS NPs dispersed in NSW
6 (Della Torre et al., 2014; Bergami et al., 2016). NP surface charge represents one of main property
7 driving NP behaviour in aquatic environments, in terms of stability, aggregation but also fate as
8 mobility or deposition (Quigg et al., 2013), therefore secondary characterization is needed in
9 ecotoxicological studies, as also recommended by SCENIHR (2007).

10 In general, the low stability observed in the exposure media can be due mainly to the high content
11 of ionic salts (salinity from 30‰ in microalgae medium to 38‰ in NSW), but also to natural
12 organic matter (NOM) and other compounds such as proteins and exopolymeric substances (EPS).
13 naturally present in NSW. These factors may specifically trigger the transformation due to eco-
14 interactions of the NPs in complex aquatic environments (Quik et al., 2014; Corsi et al., 2014). For
15 instance, Kach and Ward (2008) reported that the strong aggregation of PS NPs (up to 0.5 μ m) in
16 high ionic strength media significantly reduces their bioavailability and thus limiting toxicity to
17 aquatic organisms.

18 From the overall results obtained from the secondary characterization, both microalgae *D.*
19 *tertiolecta* and brine shrimp *A. franciscana* in NSW were exposed to micro-aggregates (> 1000 nm)
20 of the ~~negativ~~negatively charged PS-COOH and nano-scale aggregates of positively charged PS-
21 NH₂ (< 200 nm). These acquired new dimensions must be taken into consideration when looking at
22 the observed toxicity.

23

24 **3.2. Effects on the green microalga**

1 The unicellular green microalga *D. tertiolecta* ~~has been~~is recommended as bioindicator for standard
2 ecotoxicity tests (IRSA, 1978; US EPA, 1974; APHA-AWWA-WEF, 1999). In the last years, many
3 studies regarding the effects of engineered NPs (e.g. Ag-~~NPs~~, TiO₂ ~~NPs~~ and ZnO NPs) on the
4 growth of this model organism have been reported (Oukarrom et al., 2012; Miller et al., 2012;
5 Hazani et al., 2013; Manzo et al., 2013). However, there is still a paucity of information on the
6 impact of nanoplastics on marine phytoplankton, which plays a fundamental role in marine
7 ecosystem's net primary productivity, accounting for more than an half of it. Moreover, it
8 influences the global carbon cycle and ultimate climate by dominating the ocean and freshwater
9 planktonic community (Field et al., 1998).

10 The two tested PS NPs were able to affect microalga *D. tertiolecta* growth in a different way
11 (Figure 2A). PS-COOH was found to not significantly alter the growth of the microalgae up to 50
12 $\mu\text{g/ml}$ ($F(6, 48) = 0.7219, n.s.$), with a maximum average inhibition ~~in average of the~~ -specific
13 growth rate ($I_{\mu i}$) of 25.37% after 72 h. Similar negligible effects on the same species were reported
14 by Sjollem et al. (2016) in term of growth inhibition and decrease in photosynthetic efficiency up
15 to 250 $\mu\text{g/ml}$ PS-COOH. Besseling et al. (2014) also reported limited effect to freshwater microalga
16 *S. obliquus* growth by 70 nm PS-COOH, with an inhibition of 2.5% after exposure to 1 g/l PS-
17 COOH.

18 The observed strong aggregation pattern of PS-COOH (Z-average > 1 μm) in the microalgae
19 medium (Figure 1A) could be related to reduced bioavailability of nano-scale PS-COOH and ~~thus~~
20 explain its ~~peculiar~~ lack of toxicity, as previously reported for other aquatic species (Handy et al.,
21 2008; Kach and Ward, 2008; Della Torre et al., 2014).

22 On the opposite, exposure to PS-NH₂, which were still present as NP aggregates of around 127 nm
23 in the medium, caused a strong inhibition of the algal growth ($F(6, 48) = 14.48, p < 0.0001$), with
24 an $EC_{50(0-72\text{ h})}$ of $12.97 \pm 0.57 \mu\text{g/ml}$ and an $I_{\mu i}$ of 17.71 ~~61.02~~ % at 0.50 ~~0.50~~ $\mu\text{g/ml}$. Alike, the inhibition
25 in freshwater microalga *P. subcapitata* growth was observed by Casado et al. (2013) after exposure

1 to 55 and 110 nm positive surface charged PEI-modified PS NPs, with EC₅₀ (0-72 h) values of 0.58
2 and 0.54 µg/ml respectively, suggesting that the amino groups on NP surface play an important role
3 in determining the toxicity.

4 Considering the results obtained from the growth inhibition test, in-depth imaging analyses (i.e.
5 fluorescent and SEM) performed on microalgae exposed to 5 µg/ml PS NPs revealed the peculiar
6 specific disposition of these charged nanoplastics. Both PS NPs were clearly adsorbed on
7 microalgae cell surface, as indicated by the yellow-green fluorescence of PS-COOH (Figure 3) and
8 the rounded aggregates of PS-NH₂ observed by SEM (Figure 4C,D). SEM images of the control
9 group (Figure 4A,B) show few clusters around microalgae cells withal, probably due to NOM
10 and/or EPS present in the medium. EPS are polysaccharides that can be actively produced and
11 released by algae concomitantly to an extracellular stress as defence mechanism (Mishra and Jha,
12 2009; Adeleye and Keller, 2014) to limit the penetration of foreign substances through the cell
13 surface (Kumar et al., 2007). Since the unicellular *D. tertiolecta* lacks rigid cell wall (Oren, 2005),
14 EPS surrounding the plasma membrane could have a major protective role against nanoplastics and
15 particularly to PS-NH₂ (EC₅₀ of 12.97 µg/ml). Therefore, it is likely that the synthesis of these
16 biopolymers is enhanced under PS-NH₂ exposure, driving the heteroaggregation observed in Figure
17 4C,-D.

18 In addition, the observed physical adsorption of PS NPs on microalgae surface has been associated
19 to a decrease in algal photosynthesis and ROS production in freshwater algae (Bhattacharya et al.,
20 2010). In our study, the decrease in growth rate observed under PS NP exposure suggest that
21 photosynthesis might be impaired and ROS production triggered in *D. tertiolecta* species. However,
22 this phenomenon appears to be negligible for negatively charged PS-COOH (EC₅₀ > 50 µg/ml),
23 probably due to the electrostatic repulsion exerted by its carboxyl groups on microalgae cell
24 membrane (Bhattacharya et al., 2010).

1 On the contrary, the adsorption of PS-NH₂ to microalgae has been reported to be dose-dependent
2 (Bhattacharya et al., 2010) and to alter the plasma membrane by generating holes (Leroueil et al.,
3 2008). Other studies support the “proton sponge” hypothesis, suggesting that positively ly-charged
4 NPs such as PS-NH₂ bind with high affinity to lipid bilayers on the cell membrane in favour of
5 cellular uptake via endocytosis (Nel et al., 2009; Van Lehn and Alexander-Katz, 2011; Lin and
6 Alexander-Katz, 2013) and thus generating toxicity (Bexiga et al., 2011; Salvati et al., 2011; Wang
7 et al., 2013). Further studies on the green microalga *D. tertiolecta* should be addressed to stress-
8 related responses in terms of ROS production but also EPS released upon exposure to nanoplastics.

9

10 **3.3. Effects on the brine shrimp**

11 As phytoplankton, zooplankton plays a pivotal role in marine trophic web and can be ~~considered~~
12 ~~seriously affected~~ ~~target of~~ ~~by~~ nanoplastic pollution, being continuously exposed to the floating
13 fraction of small plastic debris present in ~~the~~ surface waters (Moore, 2006; Matranga and Corsi,
14 2012; Frias et al., 2014).

15 In our previous study (Bergami et al., 2016), we emphasized how short-term exposure of *A.*
16 *franciscana* larvae to PS-NH₂ (in the range 5 – 100 µg/ml) ~~was~~ ~~could~~ hampering larvae motility and
17 inducing multiple molting and the latter effect was suggested as potential defence mechanism
18 against these positively ly charged nanoplastics. In this study, we decided to have an in-depth look at
19 the molecular level behind this physiological process in order to better understand any effect over a
20 prolonged exposure. Therefore, the expression of two genes (*clap* and *cstb*), which are known to be
21 involved in the molting of *A. franciscana* embryos and larvae (Warner et al., 1995; Warner and
22 Matheson, 1998; Liu and Warner, 2006), was investigated.

23 Figure 5 shows a significant up-regulation of *clap* ($p < 0.001$) and *cstb* ($p < 0.05$) genes in 48 h
24 larvae exposed to 1 µg/ml PS-NH₂ with respect to the control, whereas no differences ($p > 0.05$)

1 were observed at the lowest concentration ~~considered~~ (0.1 µg/ml). In brine shrimp *A. franciscana*,
2 *clap* gene codes for the non-catalytic sub-unit of the cathepsin L-like protease (Chen et al., 2009),
3 the major cysteine peptidase involved in processes related to growth, including molting,
4 organogenesis and tissue remodelling in early larvae (up to 44 h after hatching) (Warner et al.,
5 1995; Warner and Matheson, 1998). Through ecdysteroid stimuli, cathepsin L-like protease is
6 responsible for the release of serine proteases in the degradation of the cuticle detached from the
7 epidermis (Warner and Matheson, 1998). Therefore, the strong induction of *clap* (having a
8 normalized relative expression of 14.3) (Figure 5A) might have driven the physiological alterations
9 previously observed in 48 h larvae (Bergami et al., 2016), suggesting a specific mechanism
10 provoked by PS-NH₂ affecting larval molting.

11 The up-regulation of *cstb* gene (Figure 5B) by PS-NH₂ at 1 µg/ml could be explained as a tentative
12 of the endogenous cystatin B inhibitor in regulating the cathepsin L-like protease activity and thus
13 limit multiple molting, which is high energy consuming (Warner et al., 2004; Turk et al., 2012).
14 However, the low normalized relative expression of *cstb* compared to *clap* suggests that a long-term
15 exposure to PS-NH₂ could seriously affect brine shrimp growth, physiology and survival, as
16 previously hypothesized (Bergami et al., 2016).

17 In order to verify this hypothesis, brine shrimp larvae of *A. franciscana* were exposed to
18 ~~negativenegatively~~ -PS-COOH and positively charged PS-NH₂ for 14 d and fed with microalgae *D.*
19 *tertiolecta*.

20 The results of long-term toxicity test are reported in Figure 2B as % of mortality to increasing
21 concentrations of PS NPs. The average mortality observed in the control group was 10 ± 3.25% and
22 14 ± 6.81% for PS-COOH and PS-NH₂ experiments respectively (mean ± s.d.). Even in 14 d
23 exposure scenarios, micro-scale aggregates of PS-COOH did not significantly affect brine shrimps
24 up to 10 µg/ml (F (2, 17) = 1.112, n.s.) confirming previous findings on short-term negligible
25 effects. On the contrary, PS-NH₂ resulted better dispersed in the medium and caused a dose-

1 dependent toxicity (LC₅₀ of 0.83 µg/ml, IC 95%: 0.07 – 0.94) after 14 d. A significant difference
2 respect to the control group was found for treatments above 1 µg/ml PS-NH₂ (F (6, 62) = 84.51, p <
3 0.0001) and all ~~with the exposed organisms were~~ 100% of found dead ~~organisms~~ at concentrations
4 above 5 µg/ml. The strong effect observed could be related to PS-NH₂ low aggregation in NSW,
5 having an hydrodynamic size of 196 nm (Figure 1B) and therefore being dispersed and bioavailable
6 to planktonic species (Kach and Ward, 2008; Della Torre et al., 2014).

7 These toxicity data are more informative with respect to those from the short-term toxicity test (48
8 h), where no significant mortality up to 100 µg/ml was found (Bergami et al., 2016). Therefore, for
9 this species the long-term exposure was crucial to discriminate the different toxicity between
10 negatively and positively surface charged PS nanoplastics.

11 The higher sensitivity of brine shrimp larvae to toxicants in long-term exposure conditions
12 compared to acute and hatching tests s has been recently reported by Manfra et al. (2015) and Rotini
13 et al. (2015), as a consequence of integration of responses at different brine shrimp larval stages
14 over an extended period of time. Moreover, from an ecotoxicological point of view, in order to
15 assess the hazard posed by nanoplastics, the long-term test mimics more likely environmental
16 realistic scenarios (Lenz et al., 2016) and at the same time represents two trophic levels.

17 In terms of particles disposition, optical fluorescent microscopy showed yellow-green fluorescent
18 PS-COOH aggregates of an average size of 1 µm (Table 1) both in brine shrimp digestive tract
19 (Figure 6A,B) as well as faecal pellets (Figure 6C,D). Such disposition resembles the one observed
20 in acute exposure (Bergami et al., 2016). ~~and is~~ in agreement with studies performed on other
21 zooplanktonic species (Cole et al., 2013; Lee et al., 2013; Della Torre et al., 2014). Although it has
22 been hypothesized that plastic particles in the digestive tract could hamper the feeding, inhibit the
23 digestion and reduce physiological functions after prolonged period (Besseling et al., 2014; Cole et
24 al., 2013; Lee et al., 2013). Nevertheless, s in this study the accumulation of PS-COOH seemed not
25 affecting the survival of brine shrimps up to 10 µg/ml. Nonetheless, processes of trophic web

1 transfer (biomagnification) cannot be excluded upon the evidence of nanoplastic ingestion by
2 marine zooplankton, as already hypothesized by Cerdevall et al. (2012) and Mattson et al. (2015).
3 In addition, considering the ability of PS NPs to adsorb hydrophobic contaminants (Rios et al.,
4 2007; Rochman et al., 2013, Velzeboer et al., 2014), their uptake and sequestration in the gut of
5 exposed organisms may indicate dramatic consequences for ~~the~~ marine organisms.

6 Furthermore, the dose-dependent mortality observed validates ~~our~~ the hypothesis we formulated in
7 our previous study (Bergami et al., 2016) regarding the long-term toxicity of PS-NH₂(Bergami et
8 al., 2016),— in line with the results from gene expression showing a significant *clap* and *cstb*
9 modulation in 48 h larvae at 1 µg/ml PS-NH₂ (Figure 5). The long-term toxicity test is a semi-static
10 assay where solutions are renewed every 2-3 days (Savorelli et al., 2007; Manfra et al., 2012), thus
11 the strong induction of *clap* gene ~~after in~~ 48 h ~~in early~~ larvae indicates that PS-NH₂ ~~over a~~
12 ~~prolonged exposure, is able to~~ disrupt the physiology and the energy flow in brine shrimp
13 developing larvae over a prolonged exposure.

14 Optical microscopy images of 9 d brine shrimps exposed to high concentrations (5 and 10 µg/ml) of
15 PS-NH₂ (Figure 7B) support these findings, showing that the exposed organisms were not able to
16 reach the same larval stage of the control group (Figure 7A). However, no measurement of larvae
17 length was made at this stage (9 d) and, ~~since~~ all the ~~exposed~~ organisms exposed at these
18 concentrations were found dead ~~at these concentrations~~ after 14 d. Likewise, at lower
19 concentrations (~~0.2.5~~ and 1 µg/ml), no significant difference in total length of PS-NH₂ exposed
20 organisms versus control was observed (p > 0.05) (Figure S1). Accordingly, when *Artemia* long-
21 term toxicity test was proposed (Manfra et al., 2012), the authors underlined that growth (~~carapace~~
22 i.e. length after 14 d) as sub-lethal endpoint was far less sensitive compared to mortality.

23 In the last decade, cysteine cathepsins have been ~~also~~ associated to apoptotic signalling as a
24 consequence of lysosomal destabilization and leakage caused by ROS (Chwieralski and Bühling,
25 2006). The release of cathepsins into the cytoplasm can be related to a cascade of intracellular

1 degradative events that can lead to a direct process of apoptosis or promoting mitochondria ROS
2 generation and amplifying the activation of caspases (Chwieralski et al., 2006; Turk and Stoka,
3 2007; Turk et al., 2012).

4 In our study, the strong induction of *clap* gene encoding for cathepsin L-like protease in response to
5 PS-NH₂ ~~in brine shrimp larvae~~ confirms the apoptotic pathway already observed in our previous
6 studies in sea urchin embryos and mussel's hemocytes (Della Torre et al., 2014; Canesi et al., 2015;
7 Pinsino et al., 2017) and in studies with human cell lines (Wang et al., 2013) for the same
8 particle upon PS-NH₂ exposure.

9 Regarding the mechanism of toxicity behind PS-NH₂, different hypotheses can be formulated: (I) a
10 direct toxicity caused by the prolonged exposure up to 14 d with strong up-regulation of *clap* and
11 *cstb* genes and consequently disruption of larval molting and energetic metabolism; (II) indirect
12 toxicity, driven by the strong physical adsorption of positively charged nanoplastics on microalgae,
13 leading to uptake and toxicity to ~~the~~ brine shrimps once they feed on microalgae; (III) a
14 combination of direct and indirect toxicity as integration of prey-to-predator responses.

15 Overall, by comparing LC₅₀ values from 14 d ~~exposure of~~ brine shrimps with E(L)C₅₀ from other
16 marine model organisms exposed to the same PS NPs (Table 2), it can be noted a similar pattern of
17 toxicity among different species, likely related to the surface charge of the nanoplastics, with
18 absence of toxicity for PS-COOH opposed to strong toxic effects provoked by PS-NH₂. The long-
19 term toxicity test can be seen as the integration of prey to predator different responses and, appears
20 to be a sensitive means to determine nanoplastics toxicity. ~~Therefore, t~~The use of long-term end-
21 point should be included in the best practices for determining the impact of emerging contaminants
22 in the marine environment, such as nanoplastics and engineered nanomaterials.

23

24 **4. Conclusion**

1 Nanoplastic exposure and toxicity constitute one of the main unexplored areas of plastic pollution
2 ~~last frontier~~ to fully understand the magnitude and consequences of this threat plastic pollution and
3 ~~its consequences for to~~ the marine environment. This study investigates the effects of ~~model~~
4 negatively (-COOH) and positively (-NH₂) surface charged PS NPs to marine planktonic species at
5 low tested concentrations and in a prolonged exposure scenario. Green microalga *D. tertiolecta* and
6 brine shrimp *A. franciscana* were chosen as model organisms representing two levels at the base of
7 the marine trophic web, as prey and predator respectively. PS-COOH formed micro-scale
8 aggregates in the media and did not affect the growth of microalgae up to 50 µg/ml as well as brine
9 shrimps up to 10 µg/ml. However, these ~~negativenegatively charged~~ NPs were found adsorbed on
10 microalgae as well as accumulated and excreted in brine shrimps, suggesting a potential trophic
11 transfer along marine trophic webs. On the opposite, PS-NH₂ was found as nanometric aggregates
12 in both media, causing inhibition of algal growth (EC₅₀ = 12.97 µg/ml) and mortality in brine
13 shrimps at 14 d (LC₅₀ = 0.83 µg/ml). Moreover, 1 µg/ml PS-NH₂ significantly induced *clap* and
14 *cstb* genes, explaining the physiological alterations previously observed in 48 h larvae, but also
15 suggesting an apoptotic pathway triggered by cathepsin L-like protease in brine shrimps
16 continuously exposed to positively charged nanoplastics. Eventually, the mortality at 14 d was
17 critical to discriminate PS NP toxicity to brine shrimp, being also representative of the interactions
18 occurring between the nanoplastics and the organisms during the exposure two trophic levels.
19 Overall, our findings indicate that PS NP surface chemistry is a key parameter responsible for the
20 behaviour, eco-interactions and impact on marine phyto- and zooplankton in terms of adsorption,
21 accumulation and toxicity. The use of long-term end-point has been identified as valuable tool for
22 determining the impact of nanoplastics on brine shrimp.

23

24

25

1 **Acknowledgements**

2 This study was performed in the framework of the PhD project entitled “Polystyrene nanoparticles
3 and their impact on marine ecosystems: accumulation, disposal and toxicity in marine species from
4 Antarctic and Mediterranean Seas”. PhD student Elisa Bergami, PhD School in Geological,
5 Environmental and Polar sciences and technologies, Department of Physical, Earth and
6 Environmental Sciences, University of Siena (Italy). Funded by the University of Siena and the
7 Italian National Antarctic Museum “Felice Ippolito”. ~~The authors gratefully acknowledge Federica
8 Savorelli for having provided the culture of microalgae *D. tertiolecta* and for her precious advices
9 in the ecotoxicity tests.~~ PS NP characterization by DLS analysis was conducted at facilities
10 available at the Department of Biotechnologies, Chemistry and Pharmacy of the University of Siena
11 (Italy).

12

13

14

15

16

17

18

19

20

21

22

23

1 **References**

- 2 Adeleye, A. S., Keller, A. A., 2014. Long-term colloidal stability and metal leaching of single wall
3 carbon nanotubes: Effect of temperature and extracellular polymeric substances. *Water Res.* 49,
4 236–250.
- 5
- 6 Andrady, A. L., 2011. Microplastics in the marine environment. *Mar. Pollut. Bull.* 62, 1596–1605.
- 7
- 8 [Andrady, A. L. 2017. The plastic in microplastics: A review. *Mar. Pollut. Bull.* DOI:
9 <http://dx.doi.org/10.1016/j.marpolbul.2017.01.082>.](http://dx.doi.org/10.1016/j.marpolbul.2017.01.082)
- 10
- 11 APHA-AWWA-WEF, 1999. Standard Methods for the Examination of Water and Wastewater,
12 Greenberg A., American Public Health Association - American Water Works Association - Water
13 Environment Federation, 20th ed. Pp. 1220.
- 14
- 15 Artoxkit, M., 2014. *Artemia* toxicity screening test for estuarine and marine waters. Standard
16 Operational Procedure. Mariakerke-Gent: Microbiotests.
- 17
- 18 Bergami, E., Bocci, E., Vannuccini, M. L., Monopoli, M., Salvati, A., Dawson, K. A., Corsi, I.,
19 2016. Nano-sized polystyrene affects feeding, behavior and physiology of brine shrimps *Artemia*
20 *franciscana* larvae. *Ecotoxicol. Env. Saf.* 123, 18–25.
- 21
- 22 Besseling, E., Wang, B., Lurling, M., Koelmans, A. A., 2014. Nanoplastic affects growth of *S.*
23 *obliquus* and reproduction of *D. magna*. *Environ. Sci. Technol.* 48, 12336–12343.
- 24
- 25 Bexiga, M. G., Varela, J. A., Wang, F., Fenaroli, F., Salvati, A., Lynch, I., Simpson, J. C., Dawson,
26 K. A., 2011. Cationic nanoparticles induce caspase 3-, 7- and 9-mediated cytotoxicity in a human
27 astrocytoma cell line. *Nanotoxicology.* 5, 557-67.
- 28
- 29 Bhattacharya, P., Lin, S., Turner, J. P., Ke, P. C., 2010. Physical adsorption of charged plastic
30 nanoparticles affects algal photosynthesis. *J. Phys. Chem. C* 114, 16556–16561.
- 31
- 32 Canesi, L., Ciacci, C., Bergami, E., Monopoli, M. P., Dawson, K. A., Papa, S., Canonico, B., Corsi,
33 I., 2015. Evidence for immunomodulation and apoptotic processes induced by cationic polystyrene
34 nanoparticles in the hemocytes of the marine bivalve *Mytilus*. *Mar. Environ. Res.* 111, 34–40.
- 35
- 36 Canesi, L., Ciacci, C., Fabbri, R., Balbi, T., Salis, A., Damonte, G., Cortese, K., Caratto, V.,
37 Monopoli, M. P., Dawson, K. A., Bergami, E., Corsi, I., 2016. Interactions of cationic polystyrene
38 nanoparticles with marine bivalve hemocytes in a physiological environment: Role of soluble
39 hemolymph proteins. *Environ. Res.* 150, 73–81.
- 40
- 41 Casado, M. P., Macken, A., Byrne, H. J., 2013. Ecotoxicological assessment of silica and
42 polystyrene nanoparticles assessed by a multitrophic test battery. *Environ. Int.* 51, 97–105.
- 43
- 44 Cattaneo, A. G., Gornati, R., Chiriva-Internati, M., Bernardini, G., 2009. Ecotoxicology of
45 nanomaterials: the role of invertebrate testing. *Invertebrate Survival Journal* 6, 78–97.
- 46
- 47 Cedervall, T., Hansson, L-A., Lard, M., Frohm, B., Linse, S., 2012. Food Chain Transport of
48 Nanoparticles Affects Behaviour and Fat Metabolism in Fish. *PLoS ONE* 7 (2), e32254. DOI
49 10.1371/journal.pone.0032254.
- 50

- 1 Chen, W-H., Ge, X., Wang, W., Yu, J., Hu, S., 2009. A gene catalogue for post-diapause
2 development of an anhydrobiotic arthropod *Artemia franciscana*. BMC Genomics 10, 52. DOI
3 10.1186/1471-2164-10-52.
4
- 5 Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., Galloway, T. S., 2013.
6 Microplastic ingestion by zooplankton. Environ. Sci. Technol. 47, 6646–55.
7
- 8 Cole, M., Galloway T. S., 2015. Ingestion of Nanoplastics and Microplastics by Pacific Oyster
9 Larvae. Environ. Sci. Technol. 49, 14625–14632.
10
- 11 Comfort, K. K., Braydich-Stolle, L. K., Maurer, E. I., Hussain S. M., 2014. Less Is More: Long-
12 Term in Vitro Exposure to Low Levels of Silver Nanoparticles Provides New Insights for
13 Nanomaterial Evaluation. ACS Nano 8 (4), 3260–3271.
14
- 15 Corsi, I., Cherr, G. N., Lenihan, H. S., Labille, J., Hasselov, M., Canesi, L., Dondero, F., Frenzilli,
16 G., Hristozov, D., Pundes, V., Della Torre, C., Pinsino, A., Libralato G., Marcomini, A., Sabbioni,
17 E., Matranga, V., 2014. Common Strategies and Technologies for the Ecosafety Assessment and
18 Design of Nanomaterials Entering the Marine Environment. ACS Nano 8 (10), 9694-709.
19
- 20 Cózar, A., Echevarría, F., González-Gordillo, I. J., Irigoien, X., Úbeda, B., Hernández-Leónd, S.,
21 Palmae, A. T., Navarrof, S., García-de-Lomas, J., Ruizg, A., Fernández-de-Puellesh, M. L., Duartei
22 C. M., 2014. Plastic debris in the open ocean. P. Natl. Acad. Sci. 111 (28), 10239–10244.
23
- 24 Della Torre, C., Bergami, E., Salvati, A., Faleri, C., Cirino, P., Dawson, K. A., Corsi I., 2014.
25 Accumulation and embryotoxicity of polystyrene nanoparticles at early stage of development of sea
26 urchin Embryos *Paracentrotus lividus*. Environ. Sci. Technol. 48, 12302–12311.
27
- 28 Field, C. B., Behrenfeld, M. J., Randerson, J. T., Falkowski, P., 1998. Primary production of the
29 biosphere: integrating terrestrial and oceanic components. Science, 281, 237–240.
30
- 31 [Fotopoulou, K. N., Karapanagioti, H. N., 2012. Surface properties of beached plastic pellets. Mar.
32 Environ. Res. 81, 70–77.](#)
33
- 34 Frias, J. P. G. L., Otero, V., Sobral, P., 2014. Evidence of microplastics in samples of zooplankton
35 from Portuguese coastal waters. Mar. Environ. Res. 95, 89–95.
36
- 37 Fröhlich, E., Meindl, C., Roblegg, E., Ebner, B., Absenger, M., Pieber, T. R., 2012. Action of
38 polystyrene nanoparticles of different sizes on lysosomal function and integrity. Particle and Fibre
39 Toxicology 9, 26.
40
- 41 Gambardella, C., Mesarič, T., Milivojević, T., Sepčić, K., Gallus, L., Carbone, S., Faimali, M.,
42 2014. Effects of selected metal oxide nanoparticles on *Artemia salina* larvae: evaluation of
43 mortality and behavioural and biochemical responses. Environ. Monit. Assess. 186 (7), 4249–4259.
44
- 45 Gregory, M. R., Ryan, P. G., 1997. Pelagic plastics and other seaborne persistent synthetic debris: a
46 review of Southern Hemisphere perspectives. In: Coe, J. M., Rogers, D. B. (Eds.), Marine Debris-
47 Sources, Impacts and Solutions. Springer-Verlag, New York, pp. 49–66.
48
- 49 Handy, R. D., Owen, R., Valsami-Jones, E., 2008. The ecotoxicology of nanoparticles and
50 nanomaterials: current status, knowledge gaps, challenges, and future needs. Ecotoxicology 17,
51 315–325.

1
2 Hartmann, N., Nolte, T., Sørensen, M., Jensen, P., Baun, A., 2015. Aquatic ecotoxicity testing of
3 nanoplastics. Lessons Learned From Nanoecotoxicology. DTU Environment.
4
5 Hazani, A. A., Ibrahim, M. M., Shehata, A. I., El-Gaaly, G. A., Daoud, M., Fouad, D., Rizwana, H.,
6 Moubayed, N., 2013. Ecotoxicity of Ag-nanoparticles on two microalgae, *Chlorella vulgaris* and
7 *Dunaliella tertiolecta*. Archives of Biological Sciences 65 (4), 1447–1457.
8
9 IRSA, 1978. Metodologia di saggio algale per lo studio della contaminazione delle acque marine.
10 Quaderni dell’Istituto di Ricerca sulle Acque, 39, IRSA-CNR, Milano. Pp. 116.
11
12 ISO, 2006. Water quality - Marine algal growth inhibition test with *Skeletonema costatum* and
13 *Phaeodactylum tricornutum*, ISO/TC 147/SC 5 10253, pp. 12.
14
15 Johnston, H. J., Semmler-Behnke, M., Brown D. M., Kreyling W., Tran, L., Stone, V., 2010.
16 Evaluating the uptake and intracellular fate of polystyrene nanoparticles by primary and hepatocyte
17 cell lines in vitro. Toxicol. Appl. Pharm. 242, 66–78.
18
19 Kach, D J., Ward, J. E., 2008. The role of marine aggregates in the ingestion of picoplankton-size
20 particles by suspension-feeding molluscs. Mar. Biol. 153 (5), 797–805.
21
22 Kashiwada, S., 2006. Distribution of nanoparticles in the see-through medaka (*Oryzias latipes*).
23 Environ Health Persp 114, 1697–1702.
24
25 Klaine, S. J., Koelmans, A. A, Horne, N., Carley, S., Handy, R. D., Kapustka, L., Nowack, B., von
26 der Kammer, F., 2012. Paradigms to assess the environmental impact of manufactured
27 nanomaterials. Environ. Toxicol. Chem. 31, 3–14.
28
29 Koelmans, A. A., Besseling E., Shim, W. J., 2015. Chapter 12 – Nanoplastics in the Aquatic
30 Environment. Critical Review. In: M. Bergmann et al. (eds.), Marine Anthropogenic Litter,
31 DOI 10.1007/978-3-319-16510-3_12.
32
33 Lambert, S., Wagner, M., 2016. Characterisation of nanoplastics during the degradation of
34 polystyrene. Chemosphere 145, 265–268.
35
36 Lee, K. W., Shim, W. J., Kwon, O. Y., Kang J. H., 2013. Size-dependent effects of
37 micropolystyrene particles in the marine copepod *Tigriopus japonicus*. Environ. Sci. Technol. 47,
38 11278–11283.
39
40 Lenz, R., Enders, K., Nielsen, T. G., 2016. Microplastic exposure studies should be
41 environmentally realistic. Proc. Natl. Acad. Sci. USA. DOI 10.1073/pnas.1606615113.
42
43 Leroueil, P. R., Berry, S. A., Duthie, K., Han, G., Rotello, V. M., McNerny, D. Q., Baker, J. R. Jr.,
44 Orr, B. G., Banaszak Holl, M. M., 2008. Wide Varieties of Cationic Nanoparticles Induce Defects
45 in Supported Lipid Bilayers. Nano Lett. 8 (2), 420–424. DOI 10.1021/nl0722929.
46
47 Lin, J., Alexander-Katz, A., 2013. Cell Membranes Open “Doors” for Cationic
48 Nanoparticles/Biomolecules: Insights into Uptake Kinetics. ACS Nano 7(12), 10799–10808. DOI
49 0.1021/nn4040553.
50

- 1 Loos, C., Syrovets, T., Musyanovych A., Mailänder, V., Landfester, K., Nienhaus, G. U., Simmet
2 T., 2014. Functionalized polystyrene nanoparticles as a platform for studying bio–nano interactions.
3 Beilstein J. Nanotechnol. 5, 2403–2412. DOI 10.3762/bjnano.5.250.
4
- 5 Manfra, L., Savorelli, F., Pisapia, M., Magaletti, E., Cicero, A. M., 2012. Long-term mortality lethal
6 toxicity test with the crustacean *Artemia franciscana*. J. Vis. Exp. 62, 2182–2185.
7
- 8 Manfra, L., Savorelli, F., Di Lorenzo, B., Libralato, G., Comin, S., Conti, D., Migliore, L., 2015.
9 Intercalibration of ecotoxicity testing protocols with *Artemia franciscana*. Ecological Indicators 57,
10 41–47.
11
- 12 Manzo, S., Miglietta, M. L., Rametta, G., Buono, S., Di Francia, G., 2013. Embryotoxicity and
13 spermotoxicity of nano-sized ZnO for Mediterranean sea urchin *Paracentrotus lividus*. J. Hazard.
14 Mater. 254–255, 1–9.
15
- 16 Matranga, V., Corsi, I., 2012. Toxic effects of engineered nanoparticles in the marine environment:
17 model organisms and molecular approaches. Mar. Environ. Res. 76, 32–40.
18
- 19 Mattsson, K., Ekvall, M. T., Hansson, L-A., Linse, S., Malmendal, A., Cedervall, T., 2015. Altered
20 Behavior, Physiology, and Metabolism in Fish Exposed to Polystyrene Nanoparticles. Environ. Sci.
21 Technol. 2015, 49, 553–561.
22
- 23 Merinska, D., Dujkova, Z., 2012. Polystyrene (nano)composites with possible antibacterial effect.
24 Mathematical Methods and Techniques in Engineering and Environmental Science. ISBN: 978-1-
25 61804-046-6.
26
- 27 Miller, R. J., Bennett, S., Keller, A. A., Pease, S., Lenihan, H. S., 2012. TiO₂ nanoparticles are
28 phototoxic to marine phytoplankton. PloS one 7(1), e30321.
29
- 30 Mishra, A., Jha, B., 2009. Isolation and characterization of extracellular polymeric substances from
31 micro-algae *Dunaliella salina* under salt stress. Bioresource Technology 100, 3382–3386.
32
- 33 Moore, M. N., 2006. Do nanoparticles present ecotoxicological risks for the health of the aquatic
34 environment? Environ. Int. 32, 967–76.
35
- 36 Newman, M. C., 2010. Fundamentals of Ecotoxicology. 3rd ed. Boca Raton, FL: Taylor &
37 Francis/CRC Press.
38
- 39 Nowack, B., Bucheli, T. D., 2007. Occurrence, behaviour and effects of nanoparticles in the
40 environment. Environ. Pollut. 150, 5–22.
41
- 42 Oren, A., 2005. A hundred years of *Dunaliella* research: 1905–2005. Saline systems 1 (2), 1–14.
43
- 44 Oukarroum, A., Bras, S., Perreault, F., Popovic, R., 2012. Inhibitory effects of silver nanoparticles
45 in two green algae, *Chlorella vulgaris* and *Dunaliella tertiolecta*. Ecotoxicol. Environ. Saf. 78, 80–
46 85.
47
- 48 Pinsino, A., Bergami, E., Della Torre, C., Vannuccini, M. L., Addis, P., Secci, M., Dawson, K. A.,
49 Matranga, V., Corsi, I., 2017. Amino modified polystyrene nanoparticles affect signalling pathways

- 1 of the sea urchin (*Paracentrotus lividus*) embryos. *Nanotoxicology* 11 (2), 201 – 209. DOI
2 <http://dx.doi.org/10.1080/17435390.2017.1279360>.
- 3
- 4 Popielarski, S. R., Pun, S. H., Davis, M. E., 2005. A nanoparticle-based model delivery system to
5 guide the rational design of gene delivery to the liver. *Synthesis and characterization. Bioconjug.*
6 *Chem.* 16, 1063–70.
- 7
- 8 Quigg, A., Chin, W. - C., Chen, C. - S., Zhang, S., Jiang, Y., Miao, A. - J., Schwehr, K. - A., Xu,
9 C., Santschi, P. H., 2013. Direct and Indirect Toxic Effects of Engineered Nanoparticles on Algae:
10 Role of Natural Organic Matter. *ACS Sustainable Chem. Eng.* 1, 686–702. DOI
11 dx.doi.org/10.1021/sc400103x.
- 12
- 13 Quik, J. T. K., Velzeboer, I., Wouterse, M., Koelmans, A. A., van de Meent, D., 2014.
14 Heteroaggregation and sedimentation rates for nanomaterials in natural waters. *Water Res.* 48, 269–
15 279.
- 16
- 17 Rand, G. M., 1995. *Fundamentals of aquatic toxicology: effects, environmental fate and risk*
18 *assessment.* CRC Press.
- 19
- 20 Rios, L. M., Moore, C., Jones, P. R., 2007. Persistent organic pollutants carried by synthetic
21 polymers in the ocean environment. *Mar. Pollut. Bull.* 54, 1230–1237.
- 22
- 23 Rochman, C. M., Manzano, C., Hentsche, B. T., Simonich, S. L., Hoh, E., 2013. Polystyrene
24 plastic: a source and sink for polycyclic aromatic hydrocarbons in the marine environment. *Environ.*
25 *Sci. Technol.* 47, 13976–84.
- 26
- 27 ~~Rossi, G., Barnoud, J., Monticelli, L., 2014. Polystyrene nanoparticles perturb lipid membranes. *J.*
28 *Phys. Chem. Lett.* 5, 241–246.~~
- 29
- 30 Rogach, A., Susha, A., Caruso, F., Sukhorukov, G., Kornowski, A., Kershaw, S., Möhwald, H.,
31 Eychmüller, A., Weller, H., 2000. Nano- and Microengineering: 3-D Colloidal Photonic Crystals
32 Prepared from Sub- μm -sized Polystyrene Latex Spheres Pre-Coated with Luminescent
33 Polyelectrolyte/Nanocrystal Shells. *Adv. Mater.* 12, 333–337.
- 34
- 35 ~~Rossi, G., Barnoud, J., Monticelli, L., 2014. Polystyrene nanoparticles perturb lipid membranes. *J.*
36 *Phys. Chem. Lett.* 5, 241–246.~~
- 37
- 38 Rotini, A., Manfra, L., Canepa, S., Tornambe, A., Migliore L., 2015. Can Artemia Hatching Assay
39 Be a (Sensitive) Alternative Tool to Acute Toxicity Test? *Bull. Environ. Contam. Toxicol.* DOI
40 [10.1007/s00128-015-1626-1](http://dx.doi.org/10.1007/s00128-015-1626-1).
- 41
- 42 ~~Rouillon, C., Bussiere, P. O., Desnoux, E., Collin, S., Vial, C., 2016. Is carbonyl index a
43 quantitative probe to monitor polypropylene photodegradation? *Polym. Degrad. Stab.* 128: 200–
44 208.~~
- 45
- 46 Ruiz-Orejón, L. F., 2016. Floating plastic debris in the Central and Western Mediterranean Sea.
47 *Mar. Environ. Res.* 120, 136 – 44.
- 48
- 49 Salvati, A., Åberg, C., dos Santos, T., Varela, J., Pinto, P., Lynch, I., Dawson, K. A., 2011.
50 Experimental and theoretical comparison of intracellular import of polymeric nanoparticles and

1 small molecules: Toward models of uptake kinetics. *Nanomedicine: Nanotechnology, Biology and*
2 *Medicine* 7, 818–826.

3

4 Savorelli, F., Palazzi, D., Gorbi, G., Invidia, M., Sei, S., Magaletti, E., Gelli, F., 2007. Messa a
5 punto di una metodologia di saggio a 14 giorni su *Artemia franciscana* e *A. partenogenetica*. *Biol.*
6 *Amb.* 21, 27–36.

7

8 SCENIHR (Scientific Committee on Emerging and Newly Identified Health Risks), 2007. Opinion
9 on the scientific aspects of the existing and proposed definitions relating to products of nanoscience
10 and nanotechnologies, Brussels, Belgium: European Commission (29 November 2007).

11

12 SIRA RSS www.sira.arpat.toscana.it/

13

14 Sjollema, S. B., Redondo-Hasselerharm, P., Leslie, H. A., Kraak, M. H. S., Vethaak, A. D., 2016.
15 Do plastic particles affect microalgal photosynthesis and growth? *Aquat. Toxicol.* 170, 259–261.

16

17 Stone, V., Nowack, B., Baun, A., van den Brink, N., von der Kammer, F., Dusinska, M., Handy, R.,
18 Hankin, S., Hassellöv, M., Joner, E., Fernandes, T. F., 2010. Nanomaterials for environmental
19 studies: Classification, reference material issues, and strategies for physico-chemical
20 characterisation. *Sci. Total Environ.* 408, 1745–1754.

21

22 US EPA, 1974. Marine assay procedure bottle test. Corvallis, Oregon. Pp. 43.

23

24 Van Lehn, R. C., Alexander-Katz, A., 2011. Penetration of lipid bilayers by nanoparticles with
25 environmentally-responsive surfaces: simulations and theory. *Soft Matter* 7, 11392–11404.

26

27 [Vannuccini, M. L., Grassi, G., Leaver, M. J., Corsi, I., 2015. Combination Effects of Nano-TiO₂ and](#)
28 [2,3,7,8-Tetrachlorodibenzo-P-Dioxin \(TCDD\) on Biotransformation Gene Expression in the Liver](#)
29 [of European Sea Bass *Dicentrarchus Labrax*. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 176-](#)
30 [177: 71-78.](#)

31

32 Velev, O. D., Kaler, E. W., 1999. In Situ Assembly of Colloidal Particles into Miniaturized
33 Biosensors. *Langmuir* 15, 3693–3698. DOI 10.1021/la981729c.

34

35 Velzeboer, I., Kwadijk, C. J. A. F., Koelmans, A. A., 2014. Strong sorption of PCBs to
36 nanoplastics, microplastics, carbon nanotubes and fullerenes. *Environ. Sci. Technol.* 48, 4869–
37 4876.

38

39 von Moos, N., Burkhardt-Holm, P., Koehler, A., 2012. Uptake and effects of microplastics on cells
40 and tissues of the blue mussel *Mytilus edulis* L. after experimental exposure. *Environ. Sci. Technol.*
41 46, 11327–11335.

42

43 Wang, T., Bai, J., Jiang, X., Nienhaus, G. U., 2012. Cellular Uptake of Nanoparticles by Membrane
44 Penetration: A Study Combining Confocal Microscopy with FTIR Spectroelectrochemistry. *ACS*
45 *Nano* 6 (2), 1251–1259.

46

47 Wang, F., Bexiga, M. G., Anguissola, S., Boya, P., Simpson, J. C., Salvati, A., Dawson, K. A.,
48 2013. Time resolved study of cell death mechanisms induced by amine-modified polystyrene
49 nanoparticles. *Nanoscale* 5, 10868-76.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Wegner, A., Besseling, E., Foekema, E. M., Kamermans, P., Koelmans, A. A., 2012. Effects of nanopolystyrene on the feeding behaviour of the blue mussel (*Mytilus edulis* L.). *Environ. Toxicol. Chem.* 31 (11), 2490–97.

Yap, F. L., Zhang, Y., 2007. Assembly of polystyrene microspheres and its application in cell micropatterning. *Biomaterials* 28, 2328–38.

Zhang, H., Kuo, Y-Y., Gerecke, A. C., Wang, J., 2012. Co-Release of Hexabromocyclododecane (HBCD) and Nano- and Microparticles from Thermal Cutting of Polystyrene Foams. *Environ. Sci. Technol.* 46, 10990–10996. DOI [dx.doi.org/10.1021/es302559v](https://doi.org/10.1021/es302559v).

1 **Tables with caption**
2

3 **Table 1.** Physico-chemical parameters of PS NPs in milli-Q water (mQW, T = 20°C), microalgae
4 medium (prepared in NSW 0.45 µm filtered, T 20°C, salinity 30‰, pH 8.3) and natural sea water
5 (NSW, 0.45 µm filtered, T 25°C, salinity 38‰, pH 8.3) using DLS analysis. Z-average (nm),
6 polydispersity index (PDI) and ζ-potential (mV), referred to PS NP concentration of 50 µg/ml are
7 reported. Values are shown as average ± standard deviation of 3 measurements.
8

	40 nm PS-COOH			50 nm PS-NH ₂		
	Z-Average (nm)	PDI	ζ-potential (mV)	Z-Average (nm)	PDI	ζ-potential (mV)
mQW	54.47 ± 0.82	0.116 ± 0.03	- 66 ± 1.1	56.48 ± 0.60	0.161 ± 0.014	+ 53.1 ± 2.1
microalgae medium	1237.5 ± 107	0.226 ± 0.03	- 12.4 ± 1.7	127 ± 5.07	0.316 ± 0.033	+ 16.5 ± 3.1
NSW	1064.2 ± 100.7	0.241 ± 0.07	- 9.18 ± 2.3	196 ± 7.46	0.287 ± 0.01	+ 17.5 ± 1.4

9

10

11 **Table 2.** EC₅₀ or LC₅₀ values (µg/ml) for different marine organisms belonging to phyto- (green
12 microalga *D. tertiolecta*) and zooplankton (sea urchin *P. lividus* embryos and brine shrimp *A.*
13 *franciscana* larvae), exposed to 40 nm negatively ly PS-COOH and 50 nm positively ly PS-NH₂ charged
14 nanoplastics in NSW media.


Species	Test	PS-COOH	PS-NH ₂	Reference
<i>P. lividus</i>	Embryotoxicity (48 h)	EC ₅₀ > 50 µg/ml	EC ₅₀ = 2.61 µg/ml	Della Torre et al. (2014)
<i>D. tertiolecta</i>	Growth Inhibition (72 h)	EC ₅₀ > 50 µg/ml	EC ₅₀ = 12.97 µg/ml	present study
<i>A. franciscana</i>	Acute Toxicity (48 h)	LC ₅₀ > 100 µg/ml	LC ₅₀ > 100 µg/ml	Bergami et al. (2016)
	Long-term Toxicity (14 d)	LC ₅₀ > 10 µg/ml	LC ₅₀ = 0.83 µg/ml	present study

15


16

17

1 **Figures with caption**



2




3


4 **Figure 1.** Intensity-based size distributions by DLS analysis of 40 nm PS-COOH (A) and 50 nm
 5 PS-NH₂ (B) at 50 µg/ml in milli-Q water (mQW), natural sea water (NSW) and microalgae
 6 medium. For each medium, one independent representative measurement is reported.
 7 The X-axis showing Z-average (nm) is set at 10 nm, logarithmic scale. Graphs were edited using
 8 GraphPad Prism5.

9

10




11




1
 2 **Figure 2.** Ecotoxicity of PS NPs to planktonic organisms. Average growth rate (μ_i , hr^{-1}) of green
 3 microalga *D. tertiolecta* after 72 h of exposure to increasing concentrations of PS NPs (~~growth~~
 4 ~~inhibition test~~) (A, growth inhibition test). Values corresponding to 0 – 5 – 10 – 25 – 50 $\mu\text{g/ml}$ are
 5 reported. Mortality (%) of brine shrimp *A. franciscana* after 14 d of exposure to increasing
 6 concentrations of PS NPs. Control groups showed an average mortality of $10 \pm 3.25\%$ and $14 \pm 6.81\%$
 7 for PS-COOH and PS-NH₂ experiments respectively (~~long term lethal toxicity test~~) (B, long-term
 8 lethal toxicity test). Error bars indicate standard deviation. Graphs were edited using GraphPad
 9 Prism5.

10




11

12 **Figure 3.** Disposition of yellow-green fluorescent PS-COOH at 5 $\mu\text{g/ml}$ in green microalga *D.*
 13 *tertiolecta* (growth inhibition test) by optical (A) and fluorescent (DAPI+FITC filter) (B)
 14 microscopy. Microalgae nuclei were stained with DAPI (blue). Images were taken at 63X. Scale
 15 bar: 10 μm .




1
2
3
4
5
6

Figure 4. Disposition of unlabelled PS-NH₂ in green microalga *D. tertiolecta* (growth inhibition test). SEM images showing control (A, B) and exposed to 5 µg/ml- PS-NH₂ (C, D). Images were taken at 1600X (A, C), 6400X (B) and 12800X (D). Scale bar: 10 µm (A, C) and 1 µm (B, D).



7



1

2 **Figure 5.** Expression of *clap* (A) and *cstb* (B) genes in brine shrimp *A. franciscana* larvae after
 3 short-term exposure (48 h) to positive surface charged PS-NH₂ at 0, 0.1 and 1 µg/ml. Results are
 4 shown as mean ± standard deviation. *** and * indicates significant differences respect to the
 5 control group, corresponding to p < 0.001 and p < 0.05 respectively.

6




7

8 **Figure 6.** Disposition of yellow-green fluorescent PS-COOH at 10 µg/ml in brine shrimp *A.*
 9 *franciscana* larvae after 14 days of exposure (long-term toxicity test). Fluorescent aggregates were
 10 observed inside the digestive tract (A, B), but also excreted as fecal pellets (C, D). Scale bar: 500
 11 µm.

12

13



1

2 **Figure 7.** Effect of PS-NH₂ at 10 µg/ml on brine shrimp *A. franciscana* growth (B), compared to
3 the control group (A) after 9 days of exposure (long-term toxicity test). Scale bar: 200 µm.

4

Long-term toxicity of surface charged polystyrene nanoplastics to marine planktonic species *Dunaliella tertiolecta* and *Artemia franciscana*

Bergami E.^{1*}, Pugnolini S.¹, Vannuccini M. L.¹, Manfra L.^{2,3}, Faleri C.⁴, Savorelli F.⁵, Dawson K. A.⁶, Corsi I.¹

¹Department of Physical, Earth and Environmental Sciences, University of Siena, Italy;

²Institute for Environmental Protection and Research (ISPRA), Rome, Italy;

³Department of Biology and Evolution of Marine Organisms, Anton Dohrn Zoological Station, 80121 Naples, Italy;

⁴Department of Life Sciences, University of Siena, 53100 Siena, Italy;

⁵Regional Agency for Prevention, Environment and Energy of Emilia-Romagna (Arpae ER), Ferrara, Italy

⁶Centre for BioNano Interactions, School of Chemistry and Chemical Biology, University College Dublin, Ireland;

***Corresponding author:** Elisa Bergami; bergami@student.unisi.it; Department of Physical, Earth and Environmental Sciences, University of Siena, Italy.

Supporting Information

The following [two pages](#) of supporting information contains two tables associated to this manuscript.

Materials and Methods

Polystyrene nanoparticles

Table S1. Conversion chart showing concentrations ($\mu\text{g/ml}$) and numbers (NPs/ml) of the nanoplastics adopted in this study. The particle numbers were calculated using the equations obtained from the suppliers (see FluoSpheres® Fluorescent Microspheres and TechNote 206) and considering the nominal size of the NPs (40 nm PS-COOH from Invitrogen and 50 nm PS-NH₂ from Bangs Laboratories Inc.).

PS NPs $\mu\text{g/ml}$	PS-NH ₂ NPs/ml	PS-COOH NPs/ml
50 $\mu\text{g/ml}$	$7.31 \cdot 10^{11}$	$1.42 \cdot 10^{12}$
25 $\mu\text{g/ml}$	$3.64 \cdot 10^{11}$	$7.11 \cdot 10^{11}$
10 $\mu\text{g/ml}$	$1.46 \cdot 10^{11}$	$2.84 \cdot 10^{11}$
5 $\mu\text{g/ml}$	$7.28 \cdot 10^{10}$	$1.42 \cdot 10^{11}$
2.5 $\mu\text{g/ml}$	$3.64 \cdot 10^{10}$	$7.11 \cdot 10^{10}$
1.5 $\mu\text{g/ml}$	$2.18 \cdot 10^{10}$	$4.27 \cdot 10^{10}$
1 $\mu\text{g/ml}$	$1.46 \cdot 10^{10}$	$2.84 \cdot 10^{10}$
0.5 $\mu\text{g/ml}$	$7.281 \cdot 10^9$	$1.42 \cdot 10^{10}$

RNA extraction, cDNA synthesis and Real Time q-PCR

Table S21. Prime sequences, length, accession number and annealing temperature for brine shrimp *A. franciscana* genes investigated through RT-qPCR analysis in this study. While *cstb* and *gapdh* sequences were already published (Chen et al., 2009), primers for *clap* were designed using NCBI Primer-BLAST.

Gene	Forward primer 5'-3'	Reverse primer 5'-3'	Acc number	Annealing T (°C)
<i>clap</i>	AGCACGACATGGAACAGTGA	GCATCGTGGTTCCTCCATT	AY307377.2	57
<i>cstb</i>	GCGAGAAGTCTTATCAAGT	TCTTTTACAGGAGTGATGG	Chen et al. (2009)	52
<i>gapdh</i> *	GTTGATGGCAAACCTCGTCATA	CCACCTTCCAAGTGAGCATTA	Chen et al. (2009)	55

*housekeeping gene

Results




Figure S1. Brine shrimp length after 14 days of exposure to PS-NH₂ (at 0.5 and 1 µg/ml) compared to the control group. Results are expressed as means ± s.d. and representative of three independent experiments. No significant difference among the groups (One-way ANOVA, $p > 0.05$) was observed. The length of the organisms was measured from the cephalic region (i.e. nauplius eye) to the telson.

References

Chen, W-H., Ge, X., Wang, W., Yu, J., Hu, S., 2009. A gene catalogue for post-diapause development of an anhydrobiotic arthropod *Artemia franciscana*. BMC Genomics 10: 52. DOI: 10.1186/1471-2164-10-52.

FluoSpheres® Fluorescent Microspheres (Molecular Probes):
https://www.google.it/url?sa=t&rct=j&q=&esrc=s&source=web&cd=2&cad=rja&uact=8&ved=0ahUKEwiwwf3T8oXUAhXJXhoKHVykCrIQFgg7MAE&url=https%3A%2F%2Ftools.thermofisher.com%2Fcontent%2Ffsfs%2Fmanuals%2Fmp05000.pdf&usg=AFQjCNE_N1Hep7jhmXJchJJbbQ_hzk8-tg

NCBI Primer-BLAST <http://www.ncbi.nlm.nih.gov/tools/primer-blast/>

TechNote 206 (Bangs Laboratories Inc.):
www.bangslabs.com/sites/default/files/imce/docs/TechNote%20206%20Web.pdf