

Non-granulomatous cerebellar infection by *Acanthamoeba* spp. in an immunocompetent host

This is the peer reviewed version of the following article:

Original:

Modica, S., Miracco, C., Cusi, M.G., Tordini, G., Muzii, V.F., Iacoangeli, F., et al. (2018). Non-granulomatous cerebellar infection by *Acanthamoeba* spp. in an immunocompetent host. *INFECTION*, 46(6), 885-889 [10.1007/s15010-018-1231-4].

Availability:

This version is available <http://hdl.handle.net/11365/1066810> since 2019-01-10T11:18:50Z

Published:

DOI: <http://doi.org/10.1007/s15010-018-1231-4>

Terms of use:

Open Access

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.

(Article begins on next page)

Infection

Non-granulomatous cerebellar infection by Acanthamoeba spp. in an immunocompetent host

--Manuscript Draft--

Manuscript Number:	INFE-D-18-00433R1
Full Title:	Non-granulomatous cerebellar infection by Acanthamoeba spp. in an immunocompetent host
Article Type:	Case Report
Keywords:	Acanthamoeba; Free living amoebas; Encephalitis; Immunocompetent host.
Corresponding Author:	Francesca Montagnani, MD, PhD University of Siena Siena, ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	University of Siena
Corresponding Author's Secondary Institution:	
Corresponding Author E-Mail:	francesca.montagnani@unisi.it
First Author:	Sara Modica
First Author Secondary Information:	
Order of Authors:	Sara Modica Clelia Miracco Maria Grazia Cusi Giacinta Tordini Vitaliano Francesco Muzii Francesco Iacoangeli Claudia Nocentini Ibne Karim M. Ali, Ph.D Shantanu Roy Alfonso Cerase Giacomo Zanelli Andrea De Luca, M.D. Francesca Montagnani, MD, PhD
Order of Authors Secondary Information:	
Funding Information:	
Abstract:	Acanthamoeba spp is a free-living amoeba, frequently involved in keratitis by contact lens in immunocompetent hosts. Anecdotal reports associate Acanthamoeba spp as a cause of severe granulomatous encephalitis in immunocompromised and, less frequently, in immunocompetent subjects. Data regarding clinical and therapeutic management are scanty and no defined therapeutic guidelines are available. We describe an unusual case of non-granulomatous Acanthamoeba cerebellitis in an immunocompetent adult male, with abrupt onset of neurological impairment, subtle hemorrhagic infarction at magnetic resonance imaging, and initial suspicion of cerebellar neoplasm. Histopathological findings of excised cerebellar mass revealed the presence of necrosis and inflammation with structure resembling amoebic

	trophozoites, but without granulomas. Polymerase chain reaction from cerebellar tissue was positive for Acanthamoeba T4 genotype. Due to gastrointestinal intolerance to miltefosine, the patient was treated with long-term course of fluconazole and trimethoprim/sulphamethoxazole, obtaining complete clinical and neuroradiological resolution.
Author Comments:	



UNIVERSITÀ
DI SIENA
1240

Dipartimento di Biotecnologie Mediche

To the Editor in Chief of Infection

Siena, September-19-2018

Dear Sir,


We submit to your attention our revised paper titled “**Non-granulomatous cerebellar infection by *Acanthamoeba* spp. in an immunocompetent host**”.

We thank you and the Reviewers for the interest and the constructive criticism regarding our paper:
we modified text accordingly to suggestions and we performed a point by point reply.

Hoping this new version will be suitable for publication in Infection,

best regards,

Francesca Montagnani



Department of Medical Biotechnologies, University of Siena, University Division of
Infectious Diseases, Policlinico Le Scotte, 4° lotto piano 0
viale Bracci, 16
53100 Siena, Italy
Tel. +39(0)577 586562
Fax: +39(0)577 233462
Email: francesca.montagnani@unisi.it

Malattie Infettive

Policlinico Santa Maria Le Scotte, IV lotto, piano 0
V.le Bracci, 16 – 53100 Siena (Italia)

Tel. +39 0577 233461 Fax +39 0577 233462 – e-mail segmalinf@unisi.it – <http://www.unisi.it>

CF 80002070524 P.IVA 00273530527

POINT by POINT REPLY

Reviewer #1: The article deals with the important topic of diagnosis and treatment of *Acanthamoeba* spp. Infections in humans. The article can be helpful in similar cases of infection.

Thank you for your comment and appreciation.

Reviewer #2: Very interesting case that reminds that when an infection is caused by a rare pathogen the diagnosis may be very difficult

Thank you for your useful suggestions, we have changed text accordingly.

1) Page 3 line 2

I would suggest to add: are ubiquitous protozoa and are commonly found in lakes, swimming pools, tap water, and heating and air conditioning units.

Added as request

2) Page 3 line 59 and page 4 line 0

Due to the fact that the first suspect is linked to the haematoxylin and eosin tissue staining and microscopy, I would suggest to describe better what you have seen in Figure 2

This is what other colleagues must look for and keep in mind (as described at the CDC link <https://www.cdc.gov/dpdx/freelivingamebic/index.html>) :

The cysts of *Acanthamoeba* spp. are typically 10-25 µm in diameter. The cysts have two walls: a wrinkled fibrous outer wall (exocyst) and an inner wall (endocyst) that may be hexagonal, spherical, star-shaped or polygonal. Cysts contain only one nucleus with a large karyosome. Trophozoites of *Acanthamoeba* spp. are pleomorphic and measure approximately 15-45 µm. They often produce many spine-like processes called acanthopodia. Trophozoites contain a large nucleus with a large, centrally-located karyosome but no peripheral chromatin.

Thank you for this suggestion. To better clarify staining description, we specify “and they do not show peripheral chromatin.” into the text (page 4 line 0), we revised graphic of Figure 2 (see Figure 2_REV file), we added an inset, to emphasize the difference in size of amoeba-like structures, and we modify related legend, as follow:

“Haemorrhages and necrosis in the white matter (a). Variably-sized amoeba-like structures (b, arrows). Small (b, upper inset, thin arrows) and large (b, lower inset, thin arrow) (bar = 5 micron) amoeba-like structures admixed to small cerebellar granules (thick arrows); the asterisk on a macrophage; Unlike macrophages, they do not show peripheral chromatin. Numerous macrophages (b, thick arrow), positive for CD68 (c, thick arrow), which instead does not stain the ameboid forms (c, thin arrows).

a, Haematoxylin and eosin, original magnification (OM): x 100;

b, Haematoxylin and Eosin, OM: x 200; b, upper and lower insets, Haematoxylin and eosin, OM: x 1000;

c, CD68 immunohistochemistry, Leica bond III automated stainer (Band Polymer Refine Detection system); chromogen: diaminobenzidine; OM: x 400”

3) Page 5 line 24

Due to the lack of experience and the difficulties of microscopic diagnosis, in case of hemorrhagic cerebellitis or cerebellar abscesses without any diagnosis it could be useful to remind that a sample of cerebellar lesion or purulent fluid should be preserved frozen together with two paired serum specimen for additional molecular or serological tests in selected reference diagnostic laboratories.

Useful remind: we added the following text “Moreover, due to the lack of experience and the difficulties of microscopic diagnosis, in case of hemorrhagic cerebellitis or cerebellar abscesses without any diagnosis it could be useful to remind that a sample of cerebellar lesion or purulent fluid should be preserved frozen together with two paired serum specimens for

additional molecular or serological tests in selected reference diagnostic laboratories. An appropriately preserved tissue sample is also essential to perform a culture, which is often considered gold standard for *Acanthamoeba* detection [12] in the absence of molecular testing.” Moreover, in order to better underline the difficult clinical and radiological diagnosis at onset, a minor adjustment to the abstract was added: “We describe an unusual case of non-granulomatous *Acanthamoeba* cerebellitis in an immunocompetent adult male, with abrupt onset of neurological impairment, subtle hemorrhagic infarction at magnetic resonance imaging, and initial suspicion of cerebellar neoplasm.”

4) Page 5 line 24 after note 3

Some authors (An update on *Acanthamoeba* keratitis: diagnosis, pathogenesis and treatment Jacob Lorenzo-Morales^{1,a,*}, Naveed A. Khan^{2,a}, and Julia Walochnik^{3,a}; Parasite 2015, 22, 10 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4330640/pdf/parasite-22-10.pdf>) still wrote:

The gold standard for *Acanthamoeba* detection is still the plate culture technique . The material (corneal scrapings/biopsies or transport medium/contact lenses/swabs, etc.) is applied centrally onto a 90 mm 1.5% non-nutrient (NN) agar plate covered with a lawn (100 IL) of a 24 h old culture of non-mucous bacteria (e.g. *Escherichia coli*). Plates are sealed with Parafilm , incubated at 30 C and screened daily for amoebae, optimally by inverted phase contrast microscopy. In cases of severe infection, amoebae are usually already visible after 24-48 h. However, samples should be observed for up to 1 week to reliably prove a negative result. Alternatively, amoebae can be cultured in tissue culture flasks in a suspension of bacteria in PBS.

We provided to insert suggested reference and the following sentence was added into discussion “An appropriately preserved tissue sample is also essential to perform a culture, which is often considered gold standard for *Acanthamoeba* detection [12] in the absence of molecular testing.”

TITLE PAGE

AUTHORS: Sara Modica, Clelia Miracco, Maria Grazia Cusi, Giacinta Tordini, Vitaliano Francesco Muzii, Francesco Iacoangeli, Claudia Nocentini, Ibne Karim M. Ali, Shantanu Roy, Alfonso Cerase, Giacomo Zanelli, Andrea De Luca, Francesca Montagnani*

TITLE: Non-granulomatous cerebellar infection by *Acanthamoeba* spp. in an immunocompetent host

AUTHORS AFFILIATIONS:

Sara Modica, Giacomo Zanelli, Andrea De Luca, Francesca Montagnani: Hospital Department of Specialized and Internal Medicine, Infectious Diseases Unit, University Hospital of Siena, Siena, Italy, and Department of Medical Biotechnologies, Infectious Diseases Division, University of Siena, Siena, Siena, Italy

Clelia Miracco: Department of Medicine, Surgery and Neuroscience, Pathological Anatomy Division, University of Siena, Siena, Italy

Maria Grazia Cusi, Giacinta Tordini: Hospital Department of Emergency Urgency and Diagnostic Services, Microbiology and Virology Unit, University Hospital of Siena, Siena, Italy, and Department of Medical Biotechnologies, Infectious Diseases Division, University of Siena, Siena, Siena, Italy

Vitaliano F. Muzii, Francesco Iacoangeli: Department of Medicine, Surgery and Neurosciences, Section of Neurosurgery, University of Siena, Siena, Italy

Claudia Nocentini: Radiology Unit, “Santa Maria la Gruccia” Hospital, Montevarchi (AR), Italy

Ibne Karim M. Ali, Shantanu Roy: Free-Living and Intestinal Amebas (FLIA) Lab, Waterborne Disease Prevention Branch Centers for Disease Control and Prevention, Atlanta, Georgia, USA

Alfonso Cerase: Department of Neurological and Sensorineural Sciences, Neuroimaging and Neurointervention Unit, University Hospital of Siena, Siena, Italy

* **Corresponding author:** Francesca Montagnani

Department of Medical Biotechnologies, University of Siena, University Division of Infectious Diseases,
Policlinico Le Scotte, 4° lotto piano 0, viale Bracci, 16 - 53100 Siena, Italy Tel. +39(0)577 586562 - Fax:
+39(0)577 233462 – Email: francesca.montagnani@unisi.it

KEYWORDS

Acanthamoeba; Free living amoebas; Encephalitis; Immunocompetent host.

ABSTRACT

Acanthamoeba spp is a free-living amoeba, frequently involved in keratitis by contact lens in immunocompetent hosts. Anecdotal reports associate *Acanthamoeba* spp as a cause of severe granulomatous encephalitis in immunocompromised and, less frequently, in immunocompetent subjects. Data regarding clinical and therapeutic management are scanty and no defined therapeutic guidelines are available.

We describe ~~an~~ unusual case of non-granulomatous *Acanthamoeba* ~~encephalitis-cerebellitis~~ in an immunocompetent adult male, with abrupt onset of neurological impairment, subtle hemorrhagic infarction at magnetic resonance imaging, and initial suspicion of cerebellar neoplasm. ~~H~~The histopathological findings of excised cerebellar mass revealed the presence of necrosis and inflammation with structure resembling amoebic trophozoites, but without granulomas. Polymerase chain reaction from cerebellar tissue was positive for *Acanthamoeba* T4 genotype. Due to gastrointestinal intolerance to miltefosine, the patient was treated with long-term course of fluconazole and trimethoprim/sulphamethoxazole, obtaining complete clinical and neuroradiological resolution.

INTRODUCTION

Free living amoebas (FLAs) are ubiquitous protozoa: and are commonly found in lakes, swimming pools, tap water, and heating and air conditioning units. Four genera of FLAs are actually identified as cause of human infections: *Balamuthia*, *Naegleria*, *Sappinia* and *Acanthamoeba* [1]. Although the infections of these FLAs are thought to be very rare, they may cause severe diseases in humans. *Acanthamoeba* spp are usually involved in eye-blinding keratitis in immunocompetent contact lens wearers and, less frequently, they are the cause of granulomatous amoebic encephalitis (GAE), mainly in immunocompromised subjects [1, 2]. GAE is a severe life-threatening disease, with a case fatality rate of >90%. The poor outcome of GAE is associated with the lack of a defined therapeutic management strategy, which results from scarcity of reports and insufficient clinical evidence. Since 1968, only 23 cases of GAE by *Acanthamoeba* spp. have been reported in immunocompetent patients [2–4]. Diagnosis of GAE is challenging, and misdiagnosis is frequent [5, 6]. Neuroimaging studies typically reveal a nonspecific focal brain lesion, which is not helpful in suggesting *Acanthamoeba* etiology. Search of amoebas in brain biopsies by expert microscopists is useful, but it may not resolve amoeba identification unless the characteristic amoebic cyst structures are observed [7].

CASE

On March 2016, a previously healthy 35 years-old man complained of a recent history of speech disturbance, gait instability, and clumsy hands, followed by abrupt onset of left hemiparesis, left hemihypoesthesia, dysarthria, and drowsiness. His medical history was formerly silent, except for recurrent headache in the last 20 years. The patient was born in Santo Domingo, Dominican Republic, and had moved to Italy two years prior to symptoms onset. At the Emergency Room, brain computed tomography and magnetic resonance imaging (MRI) (Figure 1, a-c) revealed a inhomogeneous contrast-enhancing right cerebellar lesion with subtle areas of hemorrhagic infarction. The patient was admitted to the Neurosurgery Unit of the Siena University Hospital (Central Italy) and a lumbar puncture was performed. The cerebrospinal fluid (CSF) showed normal chemical-physical features and was sterile. The patient was afebrile, and blood cell counts and standard blood chemistry were normal. A neoplasm was suspected and an initial neurological improvement was obtained with steroids. After 2 weeks, neurological deterioration occurred with ataxia, right dysmetria, and left VI cranial nerve palsy and surgical excision was planned. Preoperative MRI (Figure 1, d) showed restricted diffusion within the cerebellar lesion, consistent with suppuration. At surgery on April 12, the cerebellar mass grossly resembled a tumor but a macroscopically purulent fluid was also found. Intraoperative inspection of squash smears demonstrated inflammatory cells, hemorrhage and necrosis. The lesion was drained and a biopsy was performed. A haematoxylin and eosin (H&E) tissue staining revealed numerous macrophages that were further confirmed by CD68 staining (Figure 2). In addition, throughout the inflammatory areas, a few round and variably-sized structures with single nucleus and vacuolated cytoplasm, resembling amoebic trophozoites, were seen:

these did not stain with CD68 and they do not show peripheral chromatin. There were no neutrophils, epithelioid cells or giant cells, and macrophages did not aggregate to form organized granulomas.

Postoperative course was uneventful, except for wound revision, with gradual neurological improvement. The patient was transferred to the Infectious Diseases Unit and empiric intravenous (i.v.) ceftriaxone 2 gr q12h and metronidazole 750 mg q8h were started. Complete physical examination was confirmed to be negative except for previously described neurological alteration. Negative serology for *Entamoeba histolytica*, *Echinococcus* spp., cysticercosis, *Trypanosoma cruzi*, HTLV, *Treponema pallidum*, and HIV was obtained; *Toxoplasma gondii* IgG were positive. No alteration was noted by complete hematologic and immunologic evaluation. The chest X-ray and a complete ultrasound study of the abdomen were normal.

In our setting, immunohistochemistry (IHC) assays specific for amoebas such as *Acanthamoeba* spp. were not available. Therefore, DNA extracted from the paraffin-embedded biopsy tissue was sent to the Microbiology Unit, for molecular testing, which detected *Acanthamoeba* spp. DNA by polymerase chain reaction (PCR) [8]. Sequencing of the 18S ribosomal DNA gene fragment revealed 98% identity with an *Acanthamoeba* T4 genotype (GenBank Accession number KJ652989.1). Additionally, two paired serum specimens (collected on May 10 and May 30, respectively) and paraffin embedded blocks of biopsy samples were sent to the Free-Living and Intestinal Amoebas Laboratory, CDC, Atlanta, GA, USA for further confirmation. Both sera were positive for *Acanthamoeba* antibodies by an indirect immunofluorescence assay, with a titer of 1:64 and 1:128, respectively. Moreover, at the FLIA Laboratory, further molecular analysis by a multiplex real-time PCR [8] of the brain tissues was positive for *Acanthamoeba* spp. and negative for *Balamuthia mandrillaris* and *Naegleria fowleri* DNA.

On May 27, specific therapy was initiated with fluconazole 400 mg/day i.v, trimethoprim/sulphamethoxazole 15/75 mg/kg/day i.v. and oral miltefosine 50 mg q8h. The latter was discontinued after 14 days and the dosage of trimethoprim/sulphamethoxazole was reduced due to gastrointestinal intolerance. On June 30, the patient was discharged with oral fluconazole 400 mg/day and trimethoprim/sulphamethoxazole 8/40 mg/kg/day. No major alterations of laboratory parameters were observed during the treatment except for occasional hyperkalemia.

Forty-five and seventy-five day-postoperative follow-up MRI documented a progressive reduction of the size and of the gadolinium enhancement of the purulent foci, (Figure 1, e-f). On October 26, a new serum sample was negative for *Acanthamoeba* antibodies. The treatment was discontinued in January 2017.

In May 2017, brain MRI was negative for purulent foci, showing scarring evolution. The neurological examination on July 2017 was normal.

DISCUSSION

1 GAE due to *Acanthamoeba* species occurs predominantly in immunocompromised hosts. In this study, the patient was
2 immunocompetent. MRI showed evolution from hemorrhagic cerebellitis to cerebellar abscesses. Histopathological
3 findings, although suggestive of amoebic cerebellitis, did not allow a certain histopathological diagnosis: there were no
4 specific and classical diagnostic clues, *Acanthamoeba* immunohistochemistry was not available, and brain granulomas
5 were absent, which is unusual for *Acanthamoeba* brain infections and described in few reports in immunocompromised
6 hosts [9-11].

7
8 The accurate pathological diagnosis of amoebic brain infections is challenging. This is partly because the amoebas are
9 not considered in the initial differential diagnosis due to their rare role in causing infections in humans, and because of
10 the inherent complexity in distinguishing amoebas from the amoeba-shaped host macrophages. In our case, the
11 exclusion of other causes, along with a combination of several diagnostic approaches, including histopathology, nucleic
12 acid amplification and serology and the therapeutic response, lead to the final diagnosis. This case emphasizes the
13 complexity of the diagnosis of certain brain ~~*Acanthamoeba*~~*Acanthamoeba* infections and suggests how molecular
14 techniques and serology may be helpful. Moreover, due to the lack of experience and the difficulties of microscopic
15 diagnosis, in case of hemorrhagic cerebellitis or cerebellar abscesses without any diagnosis it could be useful to remind
16 that a sample of cerebellar lesion or purulent fluid should be preserved frozen together with two paired serum
17 specimens for additional molecular or serological tests in selected reference diagnostic laboratories. An appropriately
18 preserved tissue sample is also essential to perform a culture, which is often considered gold standard for
19 *Acanthamoeba* detection [12] in the absence of molecular testing.

20
21 Therapeutic management in our case was problematic due to tolerability issues, however a complete recovery was
22 obtained using two antimicrobials for 7 months. Overall, there are no established antimicrobial regimens against
23 *Acanthamoeba* brain infections. Recently, miltefosine has been added to the treatment armamentarium, in combination
24 with other antimicrobials, and almost all recent *Acanthamoeba* brain infection survivors received miltefosine in their
25 treatment regimens. However its effectiveness remains unclear since the same drug regimens containing miltefosine
26 were effective only in a part of the treated patients [1, 2]. In our case, miltefosine treatment probably played a marginal
27 role, given its short duration, which was due to drug intolerance. Therefore, we conclude that a “patient-customized”
28 treatment supported by the available literature evidence appears to be the most effective way to manage such rare
29 parasitic infections.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

NOTES

Compliance with ethical standards

Funding sources None

Acknowledgments We thank Marguerite Kelly Keating, formerly Infectious Disease Pathology Branch, Centers for Disease Control and Prevention (CDC), for critical revision of the manuscript.

Conflict of interest SM, CM, MGC, GC, VFM, FI, CN, IKMA, SR, AC, GZ: No conflict.

ADL reports research grants to his Institutions by ViiV Healthcare, Gilead Sciences (fellowship program) and MSD Italy; he has received consultant fees from ViiV Healthcare, Gilead Sciences, MSD Italy, Janssen, Bristol-Myers Squibb and Abbvie; all outside the submitted work.

FM has received non financial support from Angelini and Astellas, outside the submitted work. She has done contract research for Novartis Vaccine and Diagnostic S.rl. (now GSK Vaccine S.r.l.) on behalf of the University Hospital of Siena; she is Infectious Diseases Consultant for GSK (consultancy fee on behalf of University of Siena).

REFERENCES

1. Koshy AA, Blackburn BG SU. Free living amebae. In: Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases. Philadelphia: Saunders; 2015. pp. 3059–69.
2. Zamora A, Henderson H, Swiatlo E. *Acanthamoeba* encephalitis: a case report and review of therapy. Surg Neurol Int. 2014; 5:68.
3. Ong TYY, Khan NA, Siddiqui R. Brain-eating amoebae: predilection sites in the brain and disease outcome. J Clin Microbiol. 2017; 55:1989–97.
4. Rodríguez-Pérez EG, Escandón-Vargas K, Ancer A. Granulomatous amebic encephalitis caused by *Acanthamoeba* sp. in an immunocompetent Mexican adult. Rev Soc Bras Med Trop. 2017; 50:432.
5. McKellar MS, Mehta LR, Greenlee JE, Hale DC, Booton GC, Kelly DJ, et al. Fatal granulomatous *Acanthamoeba* encephalitis mimicking a stroke, diagnosed by correlation of results of sequential magnetic resonance imaging, biopsy, in vitro culture, immunofluorescence analysis, and molecular analysis. J Clin Microbiol. 2006; 44:4265-9.
6. Singh P, Kochhar R, Vashishta RK, Khandelwal N, Prabhakar S, Mohindra S, et al. Amebic meningoencephalitis: spectrum of imaging findings. Am J Neuroradiol. 2006; 27:1217–21.
7. Guarner J, Bartlett J, Shieh WJ, Paddock CD, Visvesvara GS, Zaki SR. Histopathologic spectrum and immunohistochemical diagnosis of amebic meningoencephalitis. Modern Pathology. 2007; 20:1230–7.
8. Qvarnstrom Y, Visvesvara GS, Sriram R, Da Silva AJ. Multiplex real-time PCR assay for simultaneous detection of *Acanthamoeba* spp., *Balamuthia mandrillaris*, and *Naegleria fowleri*. J Clin Microbiol. 2006; 44:3589–95.
9. Satlin MJ, Graham JK, Visvesvara GS, Mena H, Marks KM, Saal SD, et al. Fulminant and fatal encephalitis caused by *Acanthamoeba* in a kidney transplant recipient: case report and literature review. Transpl Infect Dis. 2013; 15:619–26.
10. Meersseman W, Lagrou K, Sciot R, Jonckheere J, Haberler C, Walochnik J, et al. Rapidly fatal *Acanthamoeba* encephalitis and treatment of cryoglobulinemia. Emerg Infect Dis. 2007; 13:469–71.
11. Shirwadkar CG, Samant R, Sankhe M, Deshpande R, Yagi S, Schuster FL, et al. *Acanthamoeba* encephalitis in patient with systemic lupus, India. Emerg Infect Dis. 2006; 12:984-6.
12. Lorenzo-Morales J, Khan NA, Walochnik J. An update on *Acanthamoeba* keratitis: diagnosis, pathogenesis

and treatment. Parasite. 2015; 22:10

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

FIGURE CAPTIONS

Fig. 1 Neuroradiological work-up At diagnosis, unenhanced nonconsecutive axial computed tomography (a), gradient-echo axial (b, upper row), T2-weighted coronal (b, lower row), and gadolinium-enhanced T1-weighted axial (c, upper row) and coronal (c, lower row) magnetic resonance (MR) images show an irregularly gadolinium-enhancing cortico-subcortical right cerebellar hemisphere lesion (white arrows), with subtle areas of hemorrhagic infarction (black arrows) resulting in mass effect. Preoperative axial diffusion-weighted (b: 1000 sec/mm²) MR images obtained four days later (d, upper row) and at six-day follow-up (d, lower row) clearly show appearance of cortico-subcortical restricted diffusion (black open arrowheads), consistent with purulent cerebellitis. Nonconsecutive axial diffusion-weighted (b: 1000 sec/mm²) MR images obtained 45 days after surgery (e) show clear-cut multiple foci (black open arrowheads) of restricted diffusion, consistent with abscesses, which are less evident one month later (f)

Fig. 2 Histopathologic findings

Haemorrhages and necrosis in the white matter (a). Variably-sized amoeba-like structures (b, arrows). Small (b, upper inset, thin arrows) and large (b, lower inset, thin arrow) (bar = 5 micron) amoeba-like structures admixed to small cerebellar granules (thick arrows); the asterisk on a macrophage. Unlike macrophages, they do not show peripheral chromatin. Numerous macrophages (b, thick arrow), positive for CD68 (c, thick arrow), which instead does not stain the ameboid forms (c, thin arrows).

a, Haematoxylin and eosin, original magnification (OM): x 100;

b, Haematoxylin and Eosin, OM: x 200; b, upper and lower insets, Haematoxylin and eosin, OM: x 1000;

c, CD68 immunohistochemistry, Leica bond III automated stainer (Bond Polymer Refine Detection system); chromogen: diaminobenzidine; OM: x 400
~~Haemorrhages and necrosis in the white matter (a). Amoeba-like structures (b, inset, thin arrows) admixed to small cerebellar granules (b, inset, thick arrow). Numerous macrophages (b, thick arrow), positive for CD68 (c, thick arrow), which instead does not stain the ameboid forms (c, thin arrows).~~

~~a, Haematoxylin and eosin, original magnification (OM): x 100;~~

~~b, Haematoxylin and eosin, OM: x 200; C D inset, Haematoxylin and eosin, OM: x 1000;~~

~~c, CD68 immunohistochemistry, Leica bond III automated stainer (Bond Polymer Refine Detection system); chromogen: diaminobenzidine; OM: x 400~~

Figure 1

Figure 1




Figure 2

