


Clinical Practice & Epidemiology in Mental Health

Content list available at: <https://clinical-practice-and-epidemiology-in-mental-health.com>



LETTER

Propensity to Sexual Response among Adults with Obsessive-Compulsive Disorder

Andrea Pozza^{1,*}, Donatella Marazziti², Federico Mucci² and Davide Dèttore¹

¹Department of Health Sciences, University of Florence, Florence, Italy

²Department of Clinical and Experimental Medicine, Section of Psychiatry, University of Pisa, Pisa, Italy

Abstract:

Background:

Propensity to sexual excitation and inhibition is one of the key dimensions of sexuality. Clinicians working with Obsessive-Compulsive Disorder (OCD) patients rarely assess this and other aspects of sexuality, since treatment targets generally symptom reduction. Literature on sexual functioning in OCD patients is scarce and no study has focused on symptom subtypes, nor investigated the psychological processes related to sexual response.

Objective:

In the present short report, we describe an exploratory study investigating the association between symptom subtypes and propensity towards sexual excitation/inhibition in OCD patients, controlling for gender, age and antidepressant treatment.

Methods:

Seventy-two OCD patients (mean age = 34.50 years, 37.50% women) completed the Obsessive-Compulsive Inventory-Revised and the Sexual Inhibition/Sexual Excitation Scales.

Results:

Patients with more severe compulsive washing habit had a lower propensity towards excitation and a higher one towards inhibition due to threat of performance consequences (*i.e.*, contamination with sexually transmitted diseases/having an unwanted pregnancy). Patients with more severe symptoms of checking showed a higher propensity towards inhibition due to the threat of performance consequences. Gender, age and antidepressant treatment were not related to sexual functioning.

Conclusion:

Specific OCD symptom subtypes may be associated with some psychological processes involved in sexual response. Sexual well-being should be carefully evaluated by practitioners and should be regarded as a treatment target. Future studies should investigate more comprehensively the processes involved in sexuality.

Keywords: Sexual functioning, Obsessive-compulsive disorder, Washing, Checking, Antidepressants, Selective serotonin reuptake inhibitors, Symptom subtypes, Sexual therapy.

Article History

Received: August 01, 2019

Revised: September 12, 2019

Accepted: September 13, 2019

1. INTRODUCTION

Sexuality is a multifaceted construct and one of the most important dimensions which contributes to the identity and relational well-being of the individual [1]. Obsessive-Compulsive Disorder (OCD) is a psychiatric condition which

can strongly impact intimate relationships [2, 3]. Clinicians working with OCD patients rarely assess sexual well-being, since the targets of pharmacological/psychotherapeutic treatment generally are the reduction of symptoms [4]. The literature on sexual functioning in OCD patients is insufficient. The existing studies showed that OCD is often associated with a number of sexual dysfunctions, such as increased sexual dissatisfaction and avoidance, less desire and excitation, or

* Address correspondence to this author at the Department of Health Sciences, University of Florence, Via Di San Salvi 12-50135, Florence, Italy; Tel/Fax: +39 055 27 55067; E-mail: andrea.pozza@unifi.it

orgasm difficulties. Compared with the 10-40% prevalence observed in the general population and in patients with anxiety disorders, sexual dysfunctions affect 50-80% of OCD patients [5 - 10].

The Dual Control Model of human sexual functioning assumes that a satisfactory sexual response is based upon the balance of the sexual excitation and sexual inhibition neurophysiological systems, which in turn are influenced by individual psychological characteristics [11]. The model identifies three psychological processes which drive sexual response: (a) sexual excitation, (b) inhibition due to threat of performance failure, (c) inhibition due to threat of performance consequences (e.g. contagion with sexually transmitted diseases, being seen by others during sexual encounters, having an undesired pregnancy). According to this model, low and high propensity of excitation and inhibition are indicative of a higher probability of having a sexual dysfunction confirming that such propensity is one of the most important dimensions related to sexuality [12, 13]. The empirical validity of this model has been further supported by cross-sectional and longitudinal studies demonstrating that propensity to sexual excitation/inhibition predicted sexual satisfaction and the development of sexual dysfunctions in both women and men beyond the effects of demographic, relationship and medical factors [11, 14 - 18]. In addition, research based on neuro-imaging methods showed that sexual excitation/inhibition is correlated with the activation of specific brain regions involved in sexual response [19, 20]. Some evidence showed gender differences, with women reporting a lower and higher propensity towards excitation and inhibition than men, respectively [21, 22].

A strength of this model is that it considers the psychological processes associated with the sexual response instead of focusing on sexual dysfunctions [12]. This strength can help researchers and clinicians in distinguishing these processes from the effects of pharmacological treatments in psychiatric patients. The knowledge of the psychological processes involved in sexuality may support the development of strategies aimed at improving patients' sexual well-being.

Given the dimensional nature of OCD [e.g 23.], it may be expected that propensity to sexual response is impaired to a different extent across the various symptom subtypes. OCD is generally associated with a poorer perception of physical health and body than control individuals [24, 25]. There are a number of potential explanations why some subtypes may be related to a lower propensity towards excitation and inhibition. Symptoms of contamination/washing which include concerns regarding interpersonal and body/sexual contact, may be hypothesized to be more strongly related to impaired sexuality, particularly inhibition due to threat of performance consequences. Compulsive checking behaviour may also be related to impaired sexual response through the constant tendency to check bodily sexual signals and inner experiences (*i.e.* sexual fantasies/arousal). Moreover, the obsessing subtype (*i.e.* unacceptable/forbidden thoughts) may also be related to a poorer sexual response because this subtype involves morally and sexually unacceptable fantasies/thoughts which provoke anxiety/disgust. No study investigated the relationship between

symptom subtypes and psychological processes related to sexual response in OCD, since the majority of the previous studies evaluated only sexual dysfunctions.

In the present short report, we describe an exploratory study that aimed to investigate the association between symptom subtypes and propensity to sexual response based on the Dual Control Model in a group of OCD patients, controlling for the effects of gender, age and antidepressant treatment. We hypothesized that washing, checking and obsessing (forbidden thoughts) subtypes were the most closely associated with a lower propensity to excitation and a higher propensity to inhibition. Given gender differences found in the literature [21], we expected that female gender was associated with a lower excitation and a higher inhibition. We also focused on the impact of a concurrent antidepressant treatment given its expected negative effects on propensity to sexual response [26, 27].

2. MATERIALS AND METHODS

2.1. Participants' Eligibility Criteria and Procedure

To be included, subjects had (a) to meet the criteria for a primary diagnosis of OCD made by a mental health professional through the Structured Clinical Interview for DSM-IV-TR Axis I Disorders [SCID-I; 28], (b) to be 18-65 years old, and (c) to sign informed consent. Subjects were excluded if they had (a) any psychotic/bipolar/unipolar depressive disorder according to the SCID-I, (b) intellectual disability, (c) neurological diseases, (d) obsessive-compulsive symptoms induced by drugs/medical diseases, and (e) general medical diseases. The rationale for selecting the above-mentioned age group was related to the fact that the study focused on propensity to sexual response in adults with OCD. Propensity to sexual excitation/inhibition may be expected to vary across different age cohorts as shown by the previous research [29], particularly in adolescent (age lower than 18 years old) and older adult groups (age higher than 65 years old). The antidepressant treatment was not chosen as exclusion criterion because this type of medication is the first-line therapy for mild to moderate OCD and a concurrent antidepressant treatment is a rule rather the exception in majority of the OCD patients accessing mental health services [30]. Subjects were recruited through referral from mental health specialists at public/private institutes.

Participation was voluntary and uncompensated. All the subjects were asked to provide written informed consent to their participation after a description of the aims and were informed about the possibility of withdrawing their consent at any time. Materials containing personal information about participants were stored on electronic supports protected by passwords. Ethical approval was obtained from the University of Florence ethics committee. The procedures followed were in accordance with the ethical standards of the University of Florence institutional committee on human experimentation and with the Helsinki Declaration of 1975, as revised in 1983.

2.2. Measures

2.2.1. Obsessive Compulsive Inventory - Revised [OCI-R; 31]

The OCI-R measures the severity of obsessive-compulsive

[illegible]

	Not on antidepressants	.001	.978	.008	.028	.978	-.602	.619	.000
	On antidepressants	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Age (years)	.081	.778	-.002	-.284	.778	-.015	.011	.001
	OCI-R Hoarding	.026	.871	.003	.081	.936	-.077	.084	.000
	OCI-R Checking	.205	.653	-.020	-.458	.649	-.108	.068	.004
	OCI-R Ordering	.794	.377	.039	1.068	.290	-.034	.112	.020
	OCI-R Obsessing	1.644	.205	-.071	-2.015	.050	-.141	.000	.068
	OCI-R Mental Neutralizing	.199	.657	-.001	-.034	.973	-.086	.083	.000
	OCI-R Washing	6.293	.015	.122	2.921	.005	.038	.206	.132
	Not on antidepressants * OCI-R Hoarding	.083	.775	-.015	-.288	.775	-.118	.088	.001
	On antidepressants * OCI-R Hoarding	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Not on antidepressants * OCI-R Checking	1.470	.230	.065	1.212	.230	-.042	.171	.026
	On antidepressants * OCI-R Checking	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Not on antidepressants * OCI-R Ordering	.634	.429	-.037	-.796	.429	-.130	.056	.011
	On antidepressants * OCI-R Ordering	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Not on antidepressants * OCI-R Obsessing	4.064	.049	.086	2.016	.050	.001	.172	.068
	On antidepressants * OCI-R Obsessing	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Not on antidepressants * OCI-R Mental Neutralizing	.260	.612	.031	.509	.612	-.090	.151	.005
	On antidepressants * OCI-R Mental Neutralizing	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Not on antidepressants * OCI-R Washing	5.596	.021	-.115	-2.366	.021	-.213	-.018	.091
	On antidepressants * OCI-R Washing	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
							95% CI		
	Outcome: SIS1	F _(d,7d)	p-value	β	t	p-value	Lower	Upper	Partial η ²
	Intercept	53.838	<.001	2.308	6.044	<.001	1.543	3.073	.395
	Male gender	.020	.887	.024	.142	.887	-.310	.358	.000
	Female gender	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Not on antidepressants	.064	.802	-.089	-.252	.802	-.794	.616	.001
	On antidepressants	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Age (years)	1.205	.277	.008	1.098	.277	-.007	.023	.021
	OCI-R Hoarding	.393	.534	.038	.820	.416	-.055	.131	.012
	OCI-R Checking	.657	.421	-.081	-1.602	.115	-.183	.020	.044
	OCI-R Ordering	.034	.854	.005	.118	.907	-.080	.090	.000
	OCI-R Obsessing	1.405	.241	-.013	-.319	.751	-.094	.068	.002
	OCI-R Mental Neutralizing	.119	.731	.022	.446	.657	-.076	.120	.004
	OCI-R Washing	.030	.863	-.002	-.048	.962	-.099	.094	.000
	Not on antidepressants * OCI-R Hoarding	.426	.517	-.039	-.653	.517	-.158	.080	.008
	On antidepressants * OCI-R Hoarding	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Not on antidepressants * OCI-R Checking	3.326	.074	.112	1.824	.074	-.011	.235	.056
	On antidepressants * OCI-R Checking	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Not on antidepressants * OCI-R Ordering	.000	.999	-7.759E-	-.001	.999	-.107	.107	.000
	On antidepressants * OCI-R Ordering	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Not on antidepressants * OCI-R Obsessing	.451	.505	-.033	-.672	.505	-.132	.066	.008
	On antidepressants * OCI-R Obsessing	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
	Not on antidepressants * OCI-R Mental Neutralizing	.074	.786	-.019	-.272	.786	-.157	.120	.001
	On antidepressants * OCI-R Mental Neutralizing	0 ^a	0 ^a	0 ^a	0 ^a	0<			

(Table 2) *contd.....*

Age (years)	.010	.922	.001	.098	.922	-.013	.014	.000
OCI-R Hoarding	2.114	.152	.014	.328	.744	-.069	.096	.002
OCI-R Checking	5.554	.022	-.100	-2.217	.031	-.191	-.010	.081
OCI-R Ordering	.847	.361	.037	.989	.327	-.038	.113	.017
OCI-R Obsessing	1.235	.271	.053	1.476	.146	-.019	.126	.037
OCI-R Mental Neutralizing	.030	.864	.039	.891	.377	-.049	.126	.014
OCI-R Washing	6.604	.013	-.082	-1.899	.063	-.168	.004	.061
Not on antidepressants * OCI-R Hoarding	.873	.354	.050	.934	.354	-.057	.156	.015
On antidepressants * OCI-R Hoarding	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
Not on antidepressants * OCI-R Checking	1.634	.206	.070	1.278	.206	-.040	.180	.028
On antidepressants * OCI-R Checking	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
Not on antidepressants * OCI-R Ordering	.413	.523	-.031	-.643	.523	-.126	.065	.007
On antidepressants * OCI-R Ordering	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
Not on antidepressants * OCI-R Obsessing	1.697	.198	-.057	-1.303	.198	-.146	.031	.029
On antidepressants * OCI-R Obsessing	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
Not on antidepressants * OCI-R Mental Neutralizing	1.167	.285	-.067	-1.080	.285	-.190	.057	.020
On antidepressants * OCI-R Mental Neutralizing	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
Not on antidepressants * OCI-R Washing	.299	.587	.027	.547	.587	-.073	.128	.005
On antidepressants * OCI-R Washing	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a

Note. CI = confidence interval, OCI-R = Obsessive Compulsive Inventory-Revised, SES = Sexual Excitation scale, SIS1 = Inhibition Due to Threat of Performance Failure scale, SIS2 = Inhibition Due to Threat of Performance Consequences scale, ^a = This parameter is set to zero because it is redundant in the model.

4. DISCUSSION

While the focus of research and clinical practice is on symptom reduction, the propensity to sexual excitation and inhibition in OCD patients is under-assessed. An important area is the investigation of specific well-being domains beyond the symptomatology, because well-being in OCD may be impaired even when the patient shows mild symptoms [37]. The few existing studies in the literature focused on prevalence rates of sexual dysfunctions in OCD. In addition, the role of symptom subtypes has not been investigated. It may be useful to understand the psychological processes driving propensity to sexual activity across specific symptom subtypes. The present short report described the findings of the first exploratory study investigating the association between symptom subtypes and propensity to sexual excitation/inhibition in OCD patients. The strength of the work is the fact that we excluded comorbid major psychiatric conditions (*e.g.*, psychotic/bipolar/unipolar depressive disorders) and medical disorders, quite often observed in OCD patients, potentially impacting sexual functioning and intimate relations [38, 39]. We think that this a-priori methodological choice can control for potentially biased effects of factors such as other major psychiatric disorders and other diseases which can negatively impact propensity to sexual response.

We detected no gender differences in the propensity towards excitation and inhibition, in disagreement with previous data and even our hypotheses [*e.g.* 21.]. These findings might be explained by the relatively small sample size with a preponderance of men. In addition, since symptoms of washing are often more severe among females than among males [40], it may be hypothesized that there is an interaction effect between gender and this subtype: women with more severe washing habits may be expected to have lower and higher tendencies towards excitation and inhibition, respectively. Due to the small sample size, we were not able to add other predictors in the statistical model to test such an interaction effect.

Approximately 40% of patients were on antidepressant treatment. We did not find a direct association between concurrent antidepressant treatment and both the propensity towards excitation and inhibition. These findings appear in contrast with the evidence showing that most of the antidepressants can influence the sexual response, particularly when prescribed at high dosages [26]. It should be noted however, that in our OCD group, only one-third of the patients on antidepressants were taking sertraline, which has been found to be one of the antidepressants associated with the highest incidence of sexual dysfunctions [26]. This lack of difference may also be due to the fact that the SIS/SES specifically measures the psychological processes related to impaired sexual functioning such as the fear of failure during sexual performance instead of focusing on dysfunctions. We may speculate that these mechanisms are not completely affected by antidepressant medication as they involve catastrophic misinterpretations of the outcome of sexual encounters. This evidence may support the hypothesis that impaired sexual functioning may not be related exclusively to the effects of an antidepressant treatment but also to psychological cognitive and emotional interpretations of sexuality to some extent. However, it should be noted that our study did not control for some variables potentially moderating the effects of antidepressant medication such as the type of antidepressants or the dosages, as a variety of antidepressants were prescribed to our patients. In addition, the cross-sectional design does not allow us to draw conclusions about the causal effect of medication and exclude the selection bias.

In accordance with our hypotheses, we observed that OCD patients with more severe compulsive washing habit had a lower propensity towards excitation and a higher one towards inhibition due to the threat of performance failure. It may be that patients with symptoms of washing experience lower excitation because the symptomatology focuses on interpersonal contact and the risk of contagion with sexually

transmitted diseases. Several processes may explain this relationship: for example, symptoms of washing are typically related to a more intense experience of disgust and also to a stronger focus on bodily signals which in turn may decrease excitation and increase inhibition [41, 42]. The lower levels of sexual excitation related to the washing subtype may be the effect of an avoidance coping mechanism protecting the patient from external and internal triggers of obsessions (*i.e.* sexual fantasies).

Another interesting result showed that patients with more intense washing behaviour and not on antidepressants experienced higher sexual excitation indicating an interaction effect between antidepressant treatment and this symptom subtype. This finding perhaps suggests that for washers with more severe symptoms, absence of an antidepressant treatment might have a protective effect on sexual functioning. The cross-sectional design of the study does not allow to ascertain this explanation and future longitudinal research is needed.

Consistent with our predictions, we also found a significant relationship between inhibition due to the threat of performance failure and the checking subtype. An explanation might be that symptoms of checking involve a constant self-monitoring of one's internal states or body signals (*i.e.* thoughts, mental images, impulses, emotions) which might increase the fear of performance consequences (*i.e.* undesired pregnancy and/or sexually transmitted diseases) thus inhibiting sexual functioning. Repetitive checking behaviours may also be directed to body contact during the sexual encounter and this might inhibit sexual response.

The association between symptom subtypes and propensity to sexual response may be interpreted as if sexuality has an effect on symptom subtypes. A clinical implication may be that adding elements of sexual therapy during treatment may lead to an improvement in symptom severity for patients with severe symptoms of washing/checking. In addition, using exposure therapy with exercises involving sexual/body contact might have beneficial effects on symptoms of washing.

In contrast with our hypotheses, we found no significant association between the subtype of obsession and propensity to sexual response. This result may be attributed perhaps to the method of assessment. It may be that the content of unacceptable/forbidden thoughts has a stronger impact on sexuality than the general subtype of obsession. The specific content of the obsessions/compulsions may be an alternative way to explore the effects of symptom subtypes on sexuality in OCD: *e.g.*, religious/aggressive/sexual obsessions may be hypothesized to be significant predictors of impaired sexual functioning.

We also did not observe any association between the other subtypes, including hoarding, mental neutralizing and ordering, and propensity to sexual response. It may be that the content of these subtypes is not focused on aspects related to sexuality. This may suggest that patients with higher symptoms of these subtypes may not need an assessment of sexual functioning. However, the assessment of the intensity of the symptom subtypes used in this study might have introduced a methodological bias which can explain the absence of a

significant association between sexuality and these subtypes. It may be that these subtypes, which are typically less frequent in the OCD population were under-represented in our sample. Future studies should use an alternative approach and should compare sexual functioning between OCD patients diagnosed with different subtypes instead of coding the symptom subtype as a dimensional variable.

In contrast with our expectations, we did not see an association between any of the subtypes and inhibition due to performance failure. This suggests that this sexual domain may not be relevant to OCD symptoms. An interpretation may be that most of the mental energy of OCD patients is spent on struggling with repetitive, distressing thoughts (*i.e.* obsessions) and this might make sexual performance failure a less important concern for the patient. Perhaps, other variables may have a role in inhibition due to performance failure such as fear of negative evaluation which is more specific to social anxiety disorder.

Future studies should establish more reliably the causal role of symptom subtypes on sexuality and also better control for other variables (*i.e.* type of antidepressant) through a longitudinal design. In addition, they should compare OCD patients with patients with other disorders (*i.e.* anxiety disorders) to ascertain whether the effects on propensity to sexual response are specific to OCD or represent a general consequence of emotional distress. Patients with other obsessive-compulsive spectrum conditions may also be considered as a comparator group, such as body dysmorphic disorder or skin picking disorder/trichotillomania because these patients have a negative relationship with their body [43, 44]. It may be interesting to explore whether the association between subtypes and excitation/inhibition processes mediates sexual dysfunctions, such as hypoactive sexual desire disorder which is quite frequent among OCD patients [45].

A main limitation of the present study was the lack of a systematic follow-up or stratification of psychopharmacological outcomes across different subtypes of OCD. Future research should investigate prospectively whether a concurrent psychopharmacological treatment is a predictor of long-term changes in propensity to sexual excitation/inhibition and also sexual dysfunctions. It may be interesting to assess whether the psychopharmacological treatment has a differential impact on propensity to sexual response for specific symptom subtypes.

Since sexuality is a multifaceted construct [46], another limitation concerns the lack of measures covering other aspects related to it. In addition to propensity to excitation/inhibition, other measures should be used to capture the wide complexity of sexuality such as psychophysiological response patterns, beliefs about sexual functioning, self-perceived sexual attractiveness, and sexual preferences and orientation [47-49].

Finally, further research should evaluate the role of other clinical characteristics related to symptoms such as comorbid personality disorders [50], which may impact interpersonal and intimate relations, including sexual life.

CONCLUSION

Sexuality is an overlooked dimension in OCD patients.

Our study assessed the psychological processes related to sexual response in this clinical population. It is the first investigation of the relationship between symptom subtypes and propensity to sexual response. Our findings support the heterogeneity of the clinical correlates of OCD, demonstrating that perhaps there are specific relationships between some subtypes and propensity towards excitation and inhibition. These results point out the importance of considering propensity to sexual response during the assessment of patients with OCD.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Ethical approval was obtained from the Ethics Committee of University of Florence.

HUMAN AND ANIMAL RIGHTS

No Animals were used in this research. All human research procedures followed were in accordance with the ethical standards of the committee responsible for human experimentation (institutional and national), and with the Helsinki Declaration of 1975, as revised in 2013.

CONSENT FOR PUBLICATION

Written and informed consent was obtained from all the participants prior to the study.

AVAILABILITY OF DATA AND MATERIALS

The data supporting the findings of the article are not publicly available but it can be made available by the corresponding author on reasonable request.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

AP designed the study, conducted the literature searches, collected the data, conducted the statistical analysis, wrote the paper. DM designed the study and critically reviewed the first and final drafts of the paper. FM reviewed the final draft of the paper. DD reviewed and edited the final version of the paper.

REFERENCES

- [1] Stephenson KR, Meston CM. The conditional importance of sex: exploring the association between sexual well-being and life satisfaction. *J Sex Marital Ther* 2015; 41(1): 25-38. [http://dx.doi.org/10.1080/0092623X.2013.811450] [PMID: 24308354]
- [2] Ali N, Butt AJ, Bokharey IZ. Familial and Interpersonal Relations in patients with Obsessive Compulsive Disorder (OCD). *J Psychol Clin Psychiatry* 2015; 4: 00194. [http://dx.doi.org/10.15406/jpcpy.2015.04.00194]
- [3] Coluccia A, Fagiolini A, Ferretti F, Pozza A, Goracci A. Obsessive-Compulsive Disorder and quality of life outcomes: Protocol for a systematic review and meta-analysis of cross-sectional case-control studies. *Epidemiol Biostat Public Health* 2015; 12: 2.
- [4] Norberg MM, Calamari JE, Cohen RJ, Riemann BC. Quality of life in obsessive-compulsive disorder: An evaluation of impairment and a preliminary analysis of the ameliorating effects of treatment. *Depress Anxiety* 2008; 25(3): 248-59. [http://dx.doi.org/10.1002/da.20298] [PMID: 17352377]
- [5] Fontenelle LF, de Souza WF, de Menezes GB, *et al.* Sexual function and dysfunction in Brazilian patients with obsessive-compulsive disorder and social anxiety disorder. *J Nerv Ment Dis* 2007; 195(3): 254-7. [http://dx.doi.org/10.1097/01.nmd.0000243823.94086.6f] [PMID: 17468686]
- [6] Ghassemzadeh H, Raisi F, Firoozkikhastefar R, *et al.* A study on sexual function in obsessive-compulsive disorder (OCD) patients with and without depressive symptoms. *Perspect Psychiatr Care* 2017; 53(3): 208-13. [http://dx.doi.org/10.1111/ppc.12160] [PMID: 27061854]
- [7] McCabe MP, Sharlip ID, Lewis R, *et al.* Incidence and prevalence of sexual dysfunction in women and men: A consensus statement from the Fourth International Consultation on Sexual Medicine 2015. *J Sex Med* 2016; 13(2): 144-52. [http://dx.doi.org/10.1016/j.jsxm.2015.12.034] [PMID: 26953829]
- [8] Real E, Montejo A, Alonso P, Mencho JM. Sexuality and obsessive-compulsive disorder: The hidden affair. *Neuropsychiatry (London)* 2013; 3: 23-31. [http://dx.doi.org/10.2217/np.12.72]
- [9] Thakurta RG, Dhar OPSP, Sarkar S, *et al.* Prevalence and nature of sexual dysfunctions in OCD in a tertiary medical college. *Eastern Journal of Psychiatry* 2016; 17: 2.
- [10] Vulink NC, Denys D, Bus L, Westenberg HG. Sexual pleasure in women with obsessive-compulsive disorder? *J Affect Disord* 2006; 91(1): 19-25. [http://dx.doi.org/10.1016/j.jad.2005.12.006] [PMID: 16434108]
- [11] Janssen E, Vorst H, Finn P, Bancroft J. The Sexual Inhibition (SIS) and Sexual Excitation (SES) Scales: I. Measuring sexual inhibition and excitation proneness in men. *J Sex Res* 2002; 39(2): 114-26. [http://dx.doi.org/10.1080/00224490209552130] [PMID: 12476243]
- [12] Bancroft J. Sexual behavior that is "out of control": A theoretical conceptual approach. *Psychiatr Clin North Am* 2008; 31(4): 593-601. [http://dx.doi.org/10.1016/j.psc.2008.06.009] [PMID: 18996300]
- [13] Janssen E, Bancroft J. The dual-control model: The role of sexual inhibition and excitation in sexual arousal and behavior. *Psychophysiology of Sex* 2007; 15: 197-222.
- [14] Bloemendaal LBA, Laan ETM. The psychometric properties of the sexual excitation/sexual inhibition inventory for women (SESII-W) within a Dutch population. *J Sex Res* 2015; 52(1): 69-82. [http://dx.doi.org/10.1080/00224499.2013.826166] [PMID: 24215152]
- [15] Sanders SA, Graham CA, Milhausen RR, Graham AEA, Milhausen RR. Predicting sexual problems in women: The relevance of sexual excitation and sexual inhibition. *Arch Sex Behav* 2008; 37(2): 241-51. [http://dx.doi.org/10.1007/s10508-007-9235-7] [PMID: 17902042]
- [16] Velten J, Scholten S, Graham CA, Margraf J. Sexual excitation and sexual inhibition as predictors of sexual function in women: A cross-sectional and longitudinal study. *J Sex Marital Ther* 2017; 43(2): 95-109. [http://dx.doi.org/10.1080/0092623X.2015.1115792] [PMID: 26735804]
- [17] Sarin S, Amsel R, Binik YM. How hot is he? A psychophysiological and psychosocial examination of the arousal patterns of sexually functional and dysfunctional men. *J Sex Med* 2014; 11(7): 1725-40. [http://dx.doi.org/10.1111/jsm.12562] [PMID: 24820313]
- [18] Tavares IM, Laan ETM, Nobre PJ. Sexual inhibition is a vulnerability factor for orgasm problems in women. *J Sex Med* 2018; 15(3): 361-72. [http://dx.doi.org/10.1016/j.jsxm.2017.12.015] [PMID: 29396021]
- [19] Unterhorst K, Gerwinn H, Pohl A, *et al.* An exploratory study on the central nervous correlates of sexual excitation and sexual inhibition. *J Sex Res* 2018; 1-12. [http://dx.doi.org/10.1080/00224499.2018.1539462] [PMID: 30489159]
- [20] Mitchell KR, Jones KG, Wellings K, *et al.* Estimating the prevalence of sexual function problems: The impact of morbidity criteria. *J Sex Res* 2016; 53(8): 955-67. [http://dx.doi.org/10.1080/00224499.2015.1089214] [PMID: 26605494]
- [21] Carpenter D, Janssen E, Graham C, Vorst H, Wicherts J. Women's scores on the sexual inhibition/sexual excitation scales (SIS/SES): gender similarities and differences. *J Sex Res* 2008; 45(1): 36-48. [http://dx.doi.org/10.1080/00224490701808076] [PMID: 18321029]

- [22] Quinta Gomes AL, Janssen E, Santos-Iglesias P, Pinto-Gouveia J, Fonseca LM, Nobre PJ. Validation of the Sexual Inhibition and Sexual Excitation Scales (SIS/SES) in Portugal: Assessing gender differences and predictors of sexual functioning. *Arch Sex Behav* 2018; 47(6): 1721-32. [http://dx.doi.org/10.1007/s10508-017-1137-8] [PMID: 29536260]
- [23] Albert U, Bogetto F, Maina G, Saracco P, Brunatto C, Mataix-Cols D. Family accommodation in obsessive-compulsive disorder: Relation to symptom dimensions, clinical and family characteristics. *Psychiatry Res* 2010; 179(2): 204-11. [http://dx.doi.org/10.1016/j.psychres.2009.06.008] [PMID: 20483467]
- [24] Aguglia A, Signorelli MS, Albert U, Maina G. The impact of general medical conditions in obsessive-compulsive disorder. *Psychiatry Investig* 2018; 15(3): 246-53. [http://dx.doi.org/10.30773/pi.2017.06.17.2] [PMID: 29475243]
- [25] Pozza A, Ferretti F, Coluccia A. Perceived physical health in obsessive-compulsive disorder: A protocol for a systematic review and meta-analysis. *BMJ Open* 2019; 9(6):e026261 [http://dx.doi.org/10.1136/bmjopen-2018-026261] [PMID: 31175195]
- [26] Albert U, Di Salvo G, Solia F, Rosso G, Maina G. Combining Drug and Psychological Treatments for Obsessive-Compulsive Disorder: What is the Evidence, When and for Whom. *Curr Med Chem* 2018; 25(41): 5632-46. [http://dx.doi.org/10.2174/0929867324666170712114445] [PMID: 28707590]
- [27] Serretti A, Chiesa A. Treatment-emergent sexual dysfunction related to antidepressants: A meta-analysis. *J Clin Psychopharmacol* 2009; 29(3): 259-66. [http://dx.doi.org/10.1097/JCP.0b013e3181a5233f] [PMID: 19440080]
- [28] First MB, Spitzer RL, Gibbon M, Williams JB. User's guide for the Structured clinical interview for DSM-IV axis I disorders SCID-I: clinician version. American Psychiatric Pub 1997.
- [29] Pinxten W, Lievens J. Gender differences in the development of sexual excitation and inhibition through the life course: Preliminary findings from a representative study in Flanders. *J Sex Res* 2016; 53(7): 825-35. [http://dx.doi.org/10.1080/00224499.2015.1063575] [PMID: 26488561]
- [30] Pallanti S, Grassi G, Cantisani A. Emerging drugs to treat obsessive-compulsive disorder. *Expert Opin Emerg Drugs* 2014; 19(1): 67-77. [http://dx.doi.org/10.1517/14728214.2014.875157] [PMID: 24377420]
- [31] Foa EB, Huppert JD, Leiberg S, *et al.* The Obsessive-Compulsive Inventory: Development and validation of a short version. *Psychol Assess* 2002; 14(4): 485-96. [http://dx.doi.org/10.1037/1040-3590.14.4.485] [PMID: 12501574]
- [32] Sica C, Ghisi M, Altoè G, *et al.* The Italian version of the obsessive compulsive inventory: Its psychometric properties on community and clinical samples. *J Anxiety Disord* 2009; 23(2): 204-11. [http://dx.doi.org/10.1016/j.janxdis.2008.07.001] [PMID: 18701254]
- [33] Panzeri M, Dettore D, Altoè G, *et al.* T01-O-14 Factor structure of the Italian Sexual Inhibition/Excitation (SIS/SES) scales. *Sexologies* 2008; 17: S54. [http://dx.doi.org/10.1016/S1158-1360(08)72664-1]
- [34] Olejnik S, Algina J. Generalized eta and omega squared statistics: Measures of effect size for some common research designs. *Psychol Methods* 2003; 8(4): 434-47. [http://dx.doi.org/10.1037/1082-989X.8.4.434] [PMID: 14664681]
- [35] Cohen J. Statistical power analysis for the behavioral sciences. New York, NY: Routledge 1998.
- [36] I. B. M. Statistics for Windows. Version 24 0. Armonk, NY: IBM Corp 2016. Computer Software
- [37] Pozza A, Lochner C, Ferretti F, Cuomo A, Coluccia A. Does higher severity really correlate with a worse quality of life in obsessive-compulsive disorder? A meta-regression. *Neuropsychiatr Dis Treat* 2018; 14: 1013-23. [http://dx.doi.org/10.2147/NDT.S157125] [PMID: 29713171]
- [38] Pozza A, Coradeschi D, Dettore D. Do dysfunctional beliefs moderate the negative influence of comorbid severe depression on outcome of residential treatment for refractory OCD? A pilot study. *Clin Neuropsychiatry* 2013; 10: 72-83.
- [39] Isomura K, Brander G, Chang Z, *et al.* Metabolic and cardiovascular complications in obsessive-compulsive disorder: A total population, sibling comparison study with long-term follow-up. *Biol Psychiatry* 2018; 84(5): 324-31. [http://dx.doi.org/10.1016/j.biopsych.2017.12.003] [PMID: 29395042]
- [40] Labad J, Menchon JM, Alonso P, *et al.* Gender differences in obsessive-compulsive symptom dimensions. *Depress Anxiety* 2008; 25(10): 832-8. [http://dx.doi.org/10.1002/da.20332] [PMID: 17436312]
- [41] Olatunji BO, Kim J, Cox RC, Ebesutani C. Prospective associations between disgust proneness and OCD symptoms: Specificity to excessive washing compulsions. *J Anxiety Disord* 2019; 65: 34-40. [http://dx.doi.org/10.1016/j.janxdis.2019.05.003] [PMID: 31158647]
- [42] Summerfeldt LJ, Endler NS. Examining the evidence for anxiety-related cognitive biases in obsessive-compulsive disorder. *J Anxiety Disord* 1998; 12(6): 579-98. [http://dx.doi.org/10.1016/S0887-6185(98)00035-8] [PMID: 9879037]
- [43] Schneider SC, Turner CM, Storch EA, Hudson JL. Body dysmorphic disorder symptoms and quality of life: The role of clinical and demographic variables. *J Obsessive Compuls Relat Disord* 2019; 21: 1-5. [http://dx.doi.org/10.1016/j.jocrd.2018.11.002]
- [44] Pozza A, Giaquinta N, Dettore D. Borderline, avoidant, sadistic personality traits and emotion dysregulation predict different pathological skin picking subtypes in a community sample. *Neuropsychiatr Dis Treat* 2016; 12: 1861-7. [http://dx.doi.org/10.2147/NDT.S109162] [PMID: 27536108]
- [45] Aksoy UM, Aksoy SG, Maner F, Gokalp P, Yanik M. Sexual dysfunction in obsessive compulsive disorder and panic disorder. *Psychiatr Danub* 2012; 24(4): 381-5. [PMID: 23132189]
- [46] Brotto L, Atallah S, Johnson-Agbakwu C, *et al.* Psychological and interpersonal dimensions of sexual function and dysfunction. *J Sex Med* 2016; 13(4): 538-71. [http://dx.doi.org/10.1016/j.jsxm.2016.01.019] [PMID: 27045257]
- [47] Pascoal PM, Alvarez MJ, Pereira CR, Nobre P. Development and initial validation of the beliefs about sexual functioning scale: A gender invariant measure. *J Sex Med* 2017; 14(4): 613-23. [http://dx.doi.org/10.1016/j.jsxm.2017.01.021] [PMID: 28259502]
- [48] Morgan EM. Associations between young adults' use of sexually explicit materials and their sexual preferences, behaviors, and satisfaction. *J Sex Res* 2011; 48(6): 520-30. [http://dx.doi.org/10.1080/00224499.2010.543960] [PMID: 21259151]
- [49] Rosen RC, Beck JG. Patterns of sexual arousal: Psychophysiological processes and clinical applications. New York, NY: Guilford Press 1988.
- [50] Dettore D, Pozza A, Coradeschi D. Does time-intensive ERP attenuate the negative impact of comorbid personality disorders on the outcome of treatment-resistant OCD? *J Behav Ther Exp Psychiatry* 2013; 44(4): 411-7. [http://dx.doi.org/10.1016/j.jbtep.2013.04.002] [PMID: 23770674]