

Hadamard products of hypersurfaces

This is a pre print version of the following article:

Original:

Bocci, C., Carlini, E. (2022). Hadamard products of hypersurfaces. JOURNAL OF PURE AND APPLIED ALGEBRA, 226(11) [10.1016/j.jpaa.2022.107078].

Availability:

This version is available <http://hdl.handle.net/11365/1218535> since 2022-11-02T11:20:42Z

Published:

DOI: <http://doi.org/10.1016/j.jpaa.2022.107078>

Terms of use:

Open Access

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.

(Article begins on next page)

HADAMARD PRODUCTS OF HYPERSURFACES

CRISTIANO BOCCI, ENRICO CARLINI

ABSTRACT. In this paper we characterize hypersurfaces for which their Hadamard product is still a hypersurface. Then we study hypersurfaces and, more generally, varieties which are idempotent under Hadamard powers.

1. INTRODUCTION

According to the definition in [7, 8], the Hadamard product between projective varieties $V, W \subset \mathbb{P}^n$, is the closure of the image of the rational map

$$V \times W \dashrightarrow \mathbb{P}^n, \quad ([a_0 : \cdots : a_n], [b_0 : \cdots : b_n]) \mapsto [a_0 b_0 : a_1 b_1 : \cdots : a_n b_n].$$

For any projective variety V , we may consider its Hadamard square $V^{\star 2} = V \star V$ and its higher Hadamard powers $V^{\star r} = V \star V^{\star(r-1)}$.

In [7], the authors use this definition to describe the algebraic variety associated to the restricted Boltzmann machine, which is the undirected graphical model for binary random variables specified by the bipartite graph $K_{r,n}$. This variety is the r -th Hadamard power of the first secant variety of $(\mathbb{P}^1)^n$. Note that [8] concerns the case $r = 2, n = 4$.

Hadamard products and powers are well-connected to other operations of varieties. They are the multiplicative analogs of joins and secant varieties, and in tropical geometry, tropicalized Hadamard products equal Minkowski sums. It is then natural to study properties of this new operation, and see its effects on various varieties.

For this reasons, in the last few years, the Hadamard product of the projective varieties has been widely studied from the point of view of Projective Geometry. The main problem in this setting is the behaviour of the Hadamard product between varieties with many zero. The paper [2], where the Hadamard products of linear space are studied, can be considered the first step in this direction. Successively, the first author, with Calussi, Fatabbi and Lorenzini ([3]) address the Hadamard product of linear varieties not necessarily in general position, obtaining, in \mathbb{P}^2 a complete description of the possible outcomes. In \mathbb{P}^3 , under suitable conditions (which can be prove to be generic), can be shown that, for two set V and V' of collinear points, $V \star V'$ consists of $|V||V'|$ points on the two different rulings of a non-degenerate quadric. Then, in [4], they address the Hadamard product of not necessarily generic linear varieties and show that the Hilbert function of the Hadamard product $V \star W$ of two varieties, with $\dim(V), \dim(W) \leq 1$, is the product of the Hilbert functions of the original varieties V and W . Moreover, they show that the Hadamard product of two generic linear varieties V and W is projectively equivalent to a Segre embedding. In [5], the second author, with Calussi, Fatabbi and Lorenzini consider generic degenerate subvarieties $V_i \subset \mathbb{P}^n$

and compute dimension and degree formulas for the Hadamard product of the varieties V_i .

The construction of star configurations of points, via Hadamard product, described in [2], found a generalization in [6] where the authors introduce a new construction using the Hadamard product to present star configurations of codimension c of \mathbb{P}^n and which they called Hadamard star configurations. Successively, Bahmani Jafarloo and Calussi introduce a more general type of Hadamard star configuration; any star configuration constructed by their approach is called a weak Hadamard star configuration. In [1] they classify weak Hadamard star configurations and, in the case $c = n$, they investigate the existence of a (weak) Hadamard star configuration which is apolar to the generic homogeneous polynomials of degree d .

In [9], the Hadamard product is studied for projective varieties and, in particular, the authors consider Hadamard products of varieties of matrices with fixed rank also due to their connection with problems related to algebraic statistics and quantum information. The authors introduce the notion of a Hadamard decomposition and the Hadamard rank of a matrix (multiplicative versions of the well-studied additive decomposition of tensors and tensor ranks). One of the result in [9] is the following:

Proposition 1.1. *Let $V \subset \mathbb{P}^n$ be a projective variety generated by an ideal with a minimal set of generators of the type $f_{\alpha,\beta} = X^\alpha - X^\beta$. with $\alpha, \beta \in \mathbb{N}^{n+1}$. Then*

$$V^{\star 2} = V.$$

In this paper, starting from the previous result, we are interested to study which varieties are idempotent under Hadamard power. We focus, in particular on the case of hypersurfaces. Since, in general, the Hadamard product of two hypersurfaces V and W is the whole ambient space, we first study which conditions on V and W are required so that $V \star W$ is a hypersurface. We obtain a necessary and sufficient condition which is stated in Propositions 4.5 and 4.7. Successively we pass to study Hadamard powers of a hypersurface V obtaining a sufficient and necessary condition for which $V^{\star r} = V$. This results leads to Theorem 6.2 which generalizes the result of Proposition 1.1 to $V^{\star r} = V$.

The paper is organized in the following way.

In Section 2 we recall the definitions of a Hadamard product of varieties and Hadamard powers. In Section 3 we define the concept of a Hadamard transformation, $f^{\star P}$ of a polynomial f for a point P and we prove some related results leading to Theorem 3.5 which proves the connection between the ideals of V and $P \star V$.

In Section 4 we study the Hadamard product of hypersurfaces, considering also the case of union of coordinate hyperplanes. Proposition 4.5 shows that if $C, D \subset \mathbb{P}^n$ are binomial hypersurfaces with “similar” defining equations, then $C \star D$ is a binomial hypersurface. Proposition 4.7 show that this condition is also necessary.

The case of Hadamard powers of hypersurfaces, treated in Section 5, shows similar results for the case of the Hadamard product of hypersurfaces. In this case the binomial equation of the hypersurface has an extra condition on the coefficients leading us to introduce the concept of “binomial hypersurfaces of type (t, ϵ) ”.

The generalization to any power of the result in [9] is described in Section 6.

Acknowledgments: The authors are members of GNSAGA of INDAM. The second author was supported by MIUR grant Dipartimenti di Eccellenza 2018-2022 (E11G18000350001).

2. BASIC DEFINITIONS

We now recall the definitions of the Hadamard product of varieties and Hadamard powers, as introduced in [2]. We work over the field of complex numbers \mathbb{C} .

Definition 2.1. Let $H_i \subset \mathbb{P}^n, i = 0, \dots, n$, be the hyperplane $x_i = 0$ and set

$$\Delta_i = \bigcup_{0 \leq j_1 < \dots < j_{n-i} \leq n} H_{j_1} \cap \dots \cap H_{j_{n-i}}.$$

In other words, Δ_i is the i -dimensional variety of points having *at most* $i + 1$ non-zero coordinates. Thus Δ_0 is the set of coordinate points and Δ_{n-1} is the union of the coordinate hyperplanes. Note that elements of Δ_i have *at least* $n - i$ zero coordinates. We have the following chain of inclusions:

$$(2.1) \quad \Delta_0 = \{[1 : 0 : \dots : 0], \dots, [0 : \dots : 0 : 1]\} \subset \Delta_1 \subset \dots \subset \Delta_{n-1} \subset \Delta_n = \mathbb{P}^n.$$

Definition 2.2. Let $p, q \in \mathbb{P}^n$ be two points with coordinates $[a_0 : a_1 : \dots : a_n]$ and $[b_0 : b_1 : \dots : b_n]$. If $a_i b_i \neq 0$ for some i , their Hadamard product $p \star q$ of p and q , is defined as

$$p \star q = [a_0 b_0 : a_1 b_1 : \dots : a_n b_n].$$

If $a_i b_i = 0$ for all $i = 0, \dots, n$ then we say $p \star q$ is not defined.

Given varieties V and W in \mathbb{P}^n , their Hadamard product $V \star W$ is defined as

$$V \star W = \overline{\{p \star q : p \in V, q \in W, p \star q \text{ is defined}\}},$$

where the closure is in the Zariski topology.

Definition 2.3. Given a positive integer r and a variety $V \subset \mathbb{P}^n$, the r -th Hadamard power of V is

$$V^{\star r} = V \star V^{\star(r-1)},$$

where $V^{\star 0} = [1 : \dots : 1]$.

Note that $V \star W$ is a variety such that $\dim(V \star W) \leq \dim(V) + \dim(W)$ and that $V \star W$ can be empty even if neither V nor W is empty. Moreover, $\dim(V^{\star r}) \leq r \dim(V)$ and the r -th Hadamard power cannot be empty if V is not empty ([2]).

Note that if V and W are irreducible, then $V \star W$ is irreducible.

For the rest of the paper we denote by $Z(f)$ (respectively $Z(f_1, \dots, f_t)$ and $Z(J)$) the zero set of a polynomial f (respectively, of a set of polynomials f_1, \dots, f_t and of an ideal J).

3. PRELIMINARY RESULTS

Let R be the ring $\mathbb{C}[x_0, x_1, \dots, x_n]$. Given a vector of nonnegative integers $I = (i_0, \dots, i_n)$, we denote by X^I the monomial $x_0^{i_0} x_1^{i_1} \dots x_n^{i_n}$ and by $|I| = i_0 + \dots + i_n$. Similarly, if P is a point of \mathbb{P}^n with coordinates $[p_0 : p_1 : \dots : p_n]$, we denote by P^I the monomial X^I evaluated in P , that is $p_0^{i_0} p_1^{i_1} \dots p_n^{i_n}$. Moreover, if P is a point of $\mathbb{P}^n \setminus \Delta_{n-1}$ with coordinates $[p_0 : p_1 : \dots : p_n]$, we denote by $\frac{1}{P}$ the point with coordinates $[\frac{1}{p_0} : \frac{1}{p_1} : \dots : \frac{1}{p_n}]$.

Definition 3.1. Let $f \in \mathbb{C}[x_0, x_1, \dots, x_n]$ be a homogenous polynomial, of degree d , of the form $f = \sum_{|I|=d} a_I X^I$ and consider a point $P \in \mathbb{P}^n \setminus \Delta_{n-1}$. The Hadamard transformation of f by P is the polynomial

$$(3.1) \quad f^{\star P} = \sum_{|I|=d} \frac{a_I}{P^I} X^I.$$

Lemma 3.2. *If $P \in \mathbb{P}^n \setminus \Delta_{n-1}$, one has*

$$(f^{\star P})^{\star \frac{1}{P}} = (f^{\star \frac{1}{P}})^{\star P} = f.$$

Proof. Since $f^{\star \frac{1}{P}} = \sum_{|I|=d} a_I P^I X^I$, one has

$$(f^{\star P})^{\star \frac{1}{P}} = \left(\sum_{|I|=d} \frac{a_I}{P^I} X^I \right)^{\star \frac{1}{P}} = \sum_{|I|=d} \frac{a_I}{P^I} P^I X^I = \sum_{|I|=d} a_I X^I = f$$

and

$$(f^{\star \frac{1}{P}})^{\star P} = \left(\sum_{|I|=d} a_I P^I X^I \right)^{\star P} = \sum_{|I|=d} \frac{a_I P^I}{P^I} X^I = \sum_{|I|=d} a_I X^I = f.$$

□

Lemma 3.3. *Let $Q \in \mathbb{P}^n$ and $P \in \mathbb{P}^n \setminus \Delta_{n-1}$. Then $f(Q) = 0$ if and only if $f^{\star P}(P \star Q) = 0$.*

Proof. The statement easily follows from

$$\begin{aligned} f(Q) &= \left(\sum_{|I|=d} a_I Q^I \right) = \sum_{|I|=d} \frac{a_I P^I}{P^I} Q^I \\ (3.2) \quad &= \sum_{|I|=d} \frac{a_I}{P^I} (P^I Q^I) = \sum_{|I|=d} \frac{a_I}{P^I} (P \star Q)^I \\ &= f^{\star P}(P \star Q). \end{aligned}$$

□

For $P \in \mathbb{P}^n \setminus \Delta_{n-1}$ and $Q \in \mathbb{P}^n$ denote by $\frac{Q}{P}$ the Hadamard product $Q \star \frac{1}{P}$.

Lemma 3.4. *If $Q \in P \star V$ then $\frac{Q}{P} \in V$.*

Proof. If $Q \in P \star V$ then $\frac{Q}{P} \in \frac{1}{P} \star P \star V$. By the associativity of the Hadamard product one has $\frac{1}{P} \star P \star V = V$, and the claim follows. □

Theorem 3.5. *Let $V \subset \mathbb{P}^n$ be a variety and consider a point $P \in \mathbb{P}^n \setminus \Delta_{n-1}$. If $f_1, \dots, f_s \subset \mathbb{C}[x_0, \dots, x_n]$ is a generating set for $I(V)$, that is $I(V) = \langle f_1, \dots, f_s \rangle$, then $f_1^{\star P}, \dots, f_s^{\star P}$ is a generating set for $I(P \star V)$.*

Moreover, if f_1, \dots, f_s is a Groebner bases for $I(V)$, then $f_1^{\star P}, \dots, f_s^{\star P}$ is a Groebner bases for $I(P \star V)$.

Proof. We first show that the polynomials $f_1^{\star P}, \dots, f_s^{\star P}$ vanish on $P \star V$. If $Q \in P \star V$ then $Q = P \star C$ for some $C \in V$. By proof of Lemma 3.3 one has

$$f_i^{\star P}(Q) = f_i^{\star P}(P \star C) = f_i(C) = 0, \text{ for all } i = 1, \dots, s.$$

Suppose now that $g \in I(P \star V)$ is different from f_i for $i = 1, \dots, s$. Then $g(Q) = 0$ for all $Q \in P \star V$ and, by Lemma 3.3 and 3.4, one has

$$g^{\star \frac{1}{P}}\left(\frac{Q}{P}\right) = 0 \text{ for all } \frac{Q}{P} \in V$$

that is $g^{\star \frac{1}{P}} \in I(V)$. Hence, by hypothesis $g^{\star \frac{1}{P}} = \sum_i \alpha_i f_i$ and by Lemma 3.2

$$g = (g^{\star \frac{1}{P}})^{\star P} = \left(\sum_i \alpha_i f_i \right)^{\star P} = \sum_i (\alpha_i)^{\star P} f_i^{\star P}$$

and thus $f_1^{\star P}, \dots, f_s^{\star P}$ is a generating set for $I(P \star V)$.

Suppose now f_1, \dots, f_s is a Groebner bases for $I(V)$, that is, there is a monomial order such that

$$\langle LT(I(V)) \rangle = \langle LT(f_1), \dots, LT(f_s) \rangle.$$

Since $LT(f_i)$ and $LT(f_i^{*P})$ differs only by a constant, for all $i = 1, \dots, s$, one has

$$\langle LT(f_1), \dots, LT(f_s) \rangle = \langle LT(f_1^{*P}), \dots, LT(f_s^{*P}) \rangle.$$

Hence it is enough to show that $\langle LT(I(V)) \rangle = \langle LT(I(P \star V)) \rangle$. The monomials $LM(g)$, for all g in $I(V) \setminus \{0\}$ span $\langle LM(g) : g \in I(V) \setminus \{0\} \rangle$ and since $LM(g)$ and $LT(g)$ differ only by a constant one has

$$A := \langle LM(g) : g \in I(V) \setminus \{0\} \rangle = \langle LT(g) : g \in I(V) \setminus \{0\} \rangle = \langle LT(I(V)) \rangle$$

and similarly

$$B := \langle LM(g) : g \in I(P \star V) \setminus \{0\} \rangle = \langle LT(g) : g \in I(P \star V) \setminus \{0\} \rangle = \langle LT(I(P \star V)) \rangle.$$

Hence the proof is complete if we show $A = B$.

Let $h \in A$. Then there exists $g \in I(V)$ such that $h = LM(g)$. Hence $g = \sum \alpha_i f_i$ and clearly $g^{*P} = \sum \alpha_i^{*P} f_i^{*P}$ and moreover $LM(g^{*P}) = LM(g)$, since g^{*P} and g differ only by a constant, from which one has $h \in B$.

Conversely if $h \in B$, then there exists $g \in I(P \star V)$ such that $h = LM(g)$. Thus $g = \sum \alpha_i f_i^{*P}$ and, by Lemma 3.2, $g^{*\frac{1}{P}} = \sum \alpha_i^{*\frac{1}{P}} f_i$ with, again, $LM(g^{*P}) = LM(g)$, from which one has $h \in A$. \square

Remark 3.6. It is easy to prove that if f_1, \dots, f_s is a minimal Groebner bases for $I(V)$, then $f_1^{*P}, \dots, f_s^{*P}$ is a minimal Groebner bases for $I(P \star V)$.

4. HADAMARD PRODUCTS OF HYPERSURFACES

In general the Hadamard product of two hypersurfaces is not a hypersurface but the whole ambient space. However, the following results show that there are cases in which the Hadamard product of two hypersurfaces is a hypersurfaces.

First of all we analyze some pathological cases.

Lemma 4.1. *Let H_i be any coordinate hyperplane, for $i = 0, \dots, n$. Then*

- i) $H_{i_1} \star \dots \star H_{i_t}$ is the linear subspace $Z(x_{i_j} ; j = 1, \dots, t)$.
- ii) $H_i \star C = H_i$ for any hypersurface C different from a coordinate hyperplane.

Proof. For i) notice that, since the hyperplane H_{i_j} is the set of points with $x_{i_j} = 0$, the product $H_{i_1} \star \dots \star H_{i_t}$ consists of all points with $x_{i_j} = 0$, for $j = 1, \dots, t$, i.e. the desired linear space. Similarly, for ii), since the hyperplane H_i is the set of points with $x_i = 0$, the product $H_i \star C$ is contained in $H_i = 0$. To see that $H_i \star C = H_i$ it is enough to apply [2, Lemma 2.12] which yields that the tangent space to $H_i \star C$ in $P \star Q$ is

$$Q \star T_P(H_i)$$

for generic points P and Q . Note that $T_P(H_i) = H_i$ and that, since C is not contained in a coordinate hyperplane, $Q \star H_i = H_i$. \square

Before showing the results for the Hadamard product of hypersurface, we need the following definition.

Definition 4.2. An irreducible hypersurface $C \subset \mathbb{P}^n$ is called binomial hypersurface if the equation defining C is of the form

$$\alpha_1 X^{I_1} - \alpha_2 X^{I_2} = 0.$$

Remark 4.3. The condition of irreducibility of C in the previous definition forces I_1 and I_2 to be coprime (Lemma 4.30 in [10]).

Remark 4.4. From the definition of the binomial hypersurface C , if $I_1 + I_2$ has no zero entries (that is, all variables are involved in the binomial defining C), then C does not contain points in $\Delta_{n-1} \setminus \Delta_{n-2}$ since such points contain only one zero coordinate and so the equation on C does not vanish on them. If $I_1 + I_2$ has zero entries, say, for example $(I_1 + I_2)_i = 0$, then C contains points P with $P_i = 0$. However, in this case, it is still possible to apply the Hadamard transformation by such P since P_i will be not involved in the formula (3.1).

On the other side, the hypersurface C , given by the equation $\alpha_1 X^{I_1} - \alpha_2 X^{I_2} = 0$, contains all subspaces of equations

$$\left\{ \begin{array}{l} x_{i_1} = 0 \\ x_{i_2} = 0 \\ \vdots \\ x_{i_t} = 0 \\ x_{j_1} = 0 \\ x_{j_2} = 0 \\ \vdots \\ x_{j_s} = 0 \end{array} \right.$$

where $\emptyset \neq \{i_1, i_2, \dots, i_t\} \subset I_1$ and $\emptyset \neq \{j_1, j_2, \dots, j_s\} \subset I_2$. For any such subspace $W \subset C \subset \Delta_{n-2}$ and a point $P \in W$, the Hadamard product $P \star C$ is exactly W if the only zero entries of P are in coordinates $i_1, i_2, \dots, i_t, j_1, j_2, \dots, j_s$, while it is strictly contained in W if P has extra zero coordinates. In both case, however, one has $P \star C \subset C$.

Proposition 4.5. If $C, D \subset \mathbb{P}^n$ are the following binomial hypersurfaces

$$C = Z(\alpha_1 X^{I_1} - \alpha_2 X^{I_2}) \text{ and } D = Z(\beta_1 X^{I_1} - \beta_2 X^{I_2}),$$

then $C \star D$ is the binomial hypersurface

$$C \star D = Z(\alpha_1 \beta_1 X^{I_1} - \alpha_2 \beta_2 X^{I_2}).$$

Proof. Let $P \in C \setminus \Delta_{n-1}$. Since $P \in C$ one has

$$(4.1) \quad P^{I_1} = \frac{\alpha_2}{\alpha_1} P^{I_2}.$$

Moreover, by Theorem 3.5, the equation defining $P \star D$ is

$$\frac{\beta_1}{P^{I_1}} X^{I_1} - \frac{\beta_2}{P^{I_2}} X^{I_2} = 0.$$

Using (4.1) in the previous equation, we get the desired equation for $C \star D$.

If $P \in C \cap \Delta_{n-1}$ then, by Remark 4.4, P has a zero entry in a coordinate not involved in $\alpha_1 X^{I_1} - \alpha_2 X^{I_2}$ or $P \in C \cap \Delta_{n-2}$. In the first case we can still apply the first part of the proof. In the second case, P , and hence also $P \star D$ are contained

in a linear space $W \subset C, D$. Looking at the equations of C, D and $C \star D$, again by Remark 4.4, one has $P \star D \in Z(\alpha_1 \beta_1 X^{I_1} - \alpha_2 \beta_2 X^{I_2})$. \square

Remark 4.6. Theorem 2.5 in [3] states that if H and K are two hyperplanes in \mathbb{P}^n of equation respectively $a_i x_i + a_j x_j = 0$ and $b_i x_i + b_j x_j = 0$, then their Hadamard product $H \star K$ is the hyperplane of equation $a_i b_i x_i - a_j b_j x_j = 0$.

Observe that this result is a special case of Proposition 4.5 just taking as I_1 and I_2 respectively the i -th and the j -th coordinate vectors and

$$\alpha_1 = a_i, \alpha_2 = -a_j \text{ and } \beta_1 = b_i, \beta_2 = -b_j.$$

The following result shows that also the converse of Proposition 4.5 is true.

Proposition 4.7. *Let C, D be irreducible hypersurfaces not contained in Δ_{n-1} . If $C \star D$ is a hypersurface, then C and D are binomials, such that*

$$C = Z(\alpha_1 X^{I_1} - \alpha_2 X^{I_2}) \text{ and } D = Z(\beta_1 X^{I_1} - \beta_2 X^{I_2}).$$

Moreover, $C \star D = Z(\alpha_1 \beta_1 X^{I_1} - \alpha_2 \beta_2 X^{I_2})$.

Proof. Let $E = C \star D$ and suppose that

$$\begin{aligned} C &= Z(\alpha_1 X^{I_1} + \alpha_2 X^{I_2} + \dots + \alpha_s X^{I_s}), \\ D &= Z(\beta_1 X^{J_1} + \beta_2 X^{J_2} + \dots + \beta_r X^{J_r}), \\ E &= Z(\gamma_1 X^{K_1} + \gamma_2 X^{K_2} + \dots + \gamma_t X^{K_t}). \end{aligned}$$

By hypothesis $C \star Q \subseteq E$ for all $Q \in D$. But since D is not contained in Δ_{n-1} , for all points $Q \in D \setminus \Delta_{n-1}$ one has that $C \star Q$ is a hypersurface, hence $C \star Q = E$. Thus, by Theorem 3.5,

$$Z(\alpha_1 X^{I_1} + \alpha_2 X^{I_2} + \dots + \alpha_s X^{I_s})^{\star Q} = E$$

and hence $s = t$ and, possibly after relabelling, $I_i = K_i$ for $i = 1, \dots, t$.

The same argument applied to $P \star D = E$, for $P \in C$ gives $r = t$ and, possibly after relabelling, $J_i = K_i$ for $i = 1, \dots, t$.

Consider now a point $P \in C \setminus \Delta_{n-1}$. Again by Theorem 3.5 one has that the equation defining $P \star D$ is

$$\frac{\beta_1}{P^{I_1}} X^{I_1} + \frac{\beta_2}{P^{I_2}} X^{I_2} + \dots + \frac{\beta_s}{P^{I_s}} X^{I_s} = 0$$

and since $P \star D = E$ one has

$$\frac{\beta_i}{P^{I_i}} = \lambda \gamma_i, \text{ for } i = 1, \dots, s$$

for a suitable choice of λ . Hence the points of C satisfy

$$\frac{\gamma_1}{\beta_1} X^{I_1} - \frac{\gamma_i}{\beta_i} X^{I_i} = 0, \text{ for } i = 2, \dots, s.$$

Thus the ideal

$$\langle \frac{\gamma_1}{\beta_1} X^{I_1} - \frac{\gamma_2}{\beta_2} X^{I_2}, \dots, \frac{\gamma_1}{\beta_1} X^{I_1} - \frac{\gamma_s}{\beta_s} X^{I_s} \rangle$$

is contained in the ideal $\langle \alpha_1 X^{I_1} + \alpha_2 X^{I_2} + \dots + \alpha_s X^{I_s} \rangle$, but this is possible only if $s = 2$ and

$$(4.2) \quad \frac{\gamma_1}{\beta_1} = \rho \alpha_1, \quad \frac{\gamma_2}{\beta_2} = -\rho \alpha_2.$$

It follows, possibly changing the signs, that

$$\begin{aligned} C &= Z(\alpha_1 X^{I_1} - \alpha_2 X^{I_2}) \\ D &= Z(\beta_1 X^{I_1} - \beta_2 X^{I_2}) \end{aligned}$$

and, using (4.2) that

$$E = Z(\alpha_1 \beta_1 X^{I_1} - \alpha_2 \beta_2 X^{I_2}).$$

□

5. HADAMARD POWERS OF HYPERSURFACES

We study now which hypersurfaces are idempotent under Hadamard powers. The first easy case to study concerns union, eventually with multiplicities, of coordinate hyperplanes H_i .

Lemma 5.1. *Let $C \subset \mathbb{P}^n$ be the reducible hypersurface $C = H_{i_0} \cup H_{i_1} \cup \dots \cup H_{i_s}$, union of distinct coordinate hyperplanes. Then $C^{\star t} = C$ for all $t \geq 1$.*

Proof. Since $H_i^{\star t} = H_i$, for every t and $H_i \star H_j = H_i \cap H_j$ (by part i of Lemma 4.1), expanding all terms in $(H_{i_0} \cup H_{i_1} \cup \dots \cup H_{i_s})^{\star t}$ we get the desired equality. □

Definition 5.2. Let $C \subset \mathbb{P}^n$ be a binomial hypersurface. We say that C is of type (t, ϵ) if

$$\alpha_1 = 1 \text{ and } \alpha_2 = \xi^\epsilon$$

where ξ is a primitive $(t-1)$ -th root of unity and $1 \leq \epsilon \leq t-1$.

Proposition 5.3. *If $C \subset \mathbb{P}^n$ be a binomial hypersurface of type (t, ϵ) , then $C^{\star t} = C$. Moreover, if $\gcd(t-1, \epsilon) = 1$ then t is the minimal exponent for which the previous equality holds.*

Proof. By hypothesis the equation of C is $F(x_0, \dots, x_n) = X^{I_1} - \xi^\epsilon X^{I_2}$. Consider $t-1$ points $P_i \in C \setminus \Delta_{n-1}$. Each point satisfies $P_i^{I_1} = \xi^\epsilon P_i^{I_2}$ and

$$\begin{aligned} (P_1 \star \dots \star P_{t-1})^{I_1} &= P_1^{I_1} \dots P_{t-1}^{I_1} \\ &= \xi^\epsilon P_1^{I_2} \dots \xi^\epsilon P_{t-1}^{I_2} \\ (5.1) \quad &= (\xi^\epsilon)^{t-1} P_1^{I_2} \dots P_{t-1}^{I_2} \\ &= P_1^{I_2} \dots P_{t-1}^{I_2} \\ &= (P_1 \star \dots \star P_{t-1})^{I_2}. \end{aligned}$$

By Theorem 3.5, the equation for $P_1 \star \dots \star P_{t-1} \star C$ is

$$F^{\star(P_1 \star \dots \star P_{t-1})}(x_0, \dots, x_n) = \frac{X^{I_1}}{(P_1 \star \dots \star P_{t-1})^{I_1}} - \frac{\xi^\epsilon X^{I_2}}{(P_1 \star \dots \star P_{t-1})^{I_2}}$$

and, by (5.1), one has that $F^{\star(P_1 \star \dots \star P_{t-1})}(x_0, \dots, x_n)$ and $F(x_0, \dots, x_n)$ define the same hypersurface, hence $P_1 \star \dots \star P_{t-1} \star C = C$. Since this holds for every choice of $P_1, \dots, P_{t-1} \in C \setminus \Delta_{n-2}$ and, by Remark 4.4, $P_1 \star \dots \star P_{t-1} \star C \subset C$ if $P_1 \star \dots \star P_{t-1} \in C \cap \Delta_{n-2}$, one has $C^{\star t} = C$.

Suppose now that $\gcd(t-1, \epsilon) = \alpha > 1$. Since $\xi^\epsilon = e^{\frac{2\pi\epsilon i}{t-1}}$ and assuming that $t-1 = \alpha(r-1)$ and $\epsilon = \alpha\theta$ one has $\xi^\epsilon = e^{\frac{2\pi\alpha\theta i}{\alpha(r-1)}} = e^{\frac{2\pi\theta i}{r-1}}$. Hence ξ^ϵ is the power of a primitive $(r-1)$ -th root of unity and, by the first part of the proof, we get $C^{\star r} = C$.

□

The following results prove that being binomial hypersurface of type (t, ϵ) is also a necessary condition.

Proposition 5.4. *Let $C \subset \mathbb{P}^n$ be an irreducible hypersurface not contained in Δ_{n-1} . If $C^{\star t} = C$ then C is a binomial hypersurface of type (t, ϵ) . In particular, if $\gcd(t-1, \epsilon) = 1$ then t is the minimal exponent for which $C^{\star t} = C$ holds.*

Proof. Let C be a hypersurface satisfying $C^{\star t} = C$. Hence, by dimensional reasons, each successive Hadamard product in

$$C \star (C \star (C \star (\cdots (C \star C))))$$

is a hypersurface. In particular, by Proposition 4.7, applied to $C \star C$, one has that C must be binomial of equation $\alpha_1 X^{I_1} - \alpha_2 X^{I_2} = 0$. Then by Corollary 4.7 we get that the equation of $C^{\star t}$ is $\alpha_1^t X^{I_1} - \alpha_2^t X^{I_2} = 0$.

Finally, the condition $C^{\star t} = C$ implies that $\alpha_i^t = \lambda \alpha_i$, for $i = 1, 2$ and a suitable nonzero λ . Thus we get

$$\frac{\alpha_2}{\alpha_1} \left[\left(\frac{\alpha_2}{\alpha_1} \right)^{t-1} - 1 \right] = 0.$$

Excluding the trivial solution $\alpha_2 = 0$, we get that $\frac{\alpha_2}{\alpha_1}$ is a $(t-1)$ -th root of unity, showing that C is binomial of type (t, ϵ) .

If $\gcd(t-1, \epsilon) = 1$, then, by Proposition 5.3, we get the minimality of t . \square

We end this section with an analogue results for reducible hypersurfaces.

Proposition 5.5. *Let $C_1 \cup \cdots \cup C_s$ be a reducible hypersurface such that $C_i \not\subset \Delta_{n-1}$ for each component C_i , $i = 1, \dots, s$. Then $(C_1 \cup \cdots \cup C_s)^{\star r} = C_1 \cup \cdots \cup C_s$ if and only if*

1. *each component C_i is a binomial hypersurfaces of type (t, ϵ_i) and equation $X^{I_1} - \xi^{\epsilon_i} X^{I_2}$, for $i = 1, \dots, s$;*
2. *all products $\xi^{d_1 \epsilon_1} \cdots \xi^{d_s \epsilon_s}$, with $d_1 + d_2 + \cdots + d_s = r$, is equal to ξ^{ϵ_j} for some $j = 1, \dots, s$.*

Proof. Assume 1. and 2. holds. Any terms in $(C_1 \cup \cdots \cup C_s)^{\star r}$ has the form $C_1^{\star d_1} \star C_2^{\star d_2} \star \cdots \star C_s^{\star d_s}$, with $d_1 + d_2 + \cdots + d_s = r$. Condition 2, together with Proposition 4.5, applied to the binomial hypersurfaces C_i 's, of type (t, ϵ_i) , implies that $C_1^{\star d_1} \star C_2^{\star d_2} \star \cdots \star C_s^{\star d_s} = C_j$ for some $j = 1, \dots, s$.

For the other direction, the condition $(C_1 \cup \cdots \cup C_s)^{\star r} = C_1 \cup \cdots \cup C_s$ implies that $C_1^{\star d_1} \star C_2^{\star d_2} \star \cdots \star C_s^{\star d_s}$, with $d_1 + d_2 + \cdots + d_s = r$, is equal to C_j for some $j = 0, \dots, s$. In particular, if we consider $d_i = r$ we get $C_i^{\star r} = C_j$ for some $j = 1, \dots, s$ and for all $i = 1, \dots, s$. Repeatedly applying Proposition 4.7 (starting from $C_i \star C_i^{\star r-1}$) it follows that C_i is a binomial hypersurface for $i = 1, \dots, s$ and, moreover, that all these components are defined by the same binomial equation, with different coefficients. Let assume that $C_i = Z(X^{I_1} - \alpha_i X^{I_2})$, for $i = 1, \dots, s$. Hence, by the previous consideration we have

$$(5.2) \quad \alpha_1^{d_1} \cdots \alpha_s^{d_s} = \alpha_j \quad \text{for some } j = 1, \dots, s, \\ \text{with } d_1 + \cdots + d_s = r.$$

We claim that $|\alpha_i| = 1$ for $i = 1, \dots, s$. To this aim, assume, eventually reordering the components, that

$$|\alpha_1| \leq |\alpha_2| \leq \cdots \leq |\alpha_s|.$$

If there exists a index j such that $|\alpha_j| > 1$, then $|\alpha_s| > 1$. Taking the Hadamard power C_s^{*r} we get, by Proposition 4.5, a hypersurface defined by the polynomial $X^{I_1} - \alpha_s^r X^{I_2}$. Since $|\alpha_s^r| > |\alpha_s|$, condition (5.2) is not satisfied and we get a contradiction. Similarly, if there exists a index j such that $|\alpha_j| < 1$, then $|\alpha_1| < 1$. Taking the the Hadamard power C_1^{*1} we get a hypersurface defined by the polynomial $X^{I_1} - \alpha_1^r X^{I_2}$. Since $|\alpha_1^r| < |\alpha_1|$, condition (5.2) is not satisfied and, again, we get a contradiction.

Since α_i lies in the unitary circle, we can write $\alpha_i = \eta_i^{\rho_i}$ where η_i is a t_i -th root of unity, and $1 \leq \rho_i \leq t_i$, for $i = 1, \dots, s$.

Let $t-1 = \text{lcm}(t_1, t_2, \dots, t_s)$, then we can see each α_i as a $(t-1)$ -th root of unity and write $\alpha_i = \xi^{\epsilon_i}$, where ξ is a primitive $(t-1)$ -th root of unity and $1 \leq \epsilon_i \leq t-1$, for $i = 1, \dots, s$. Hence each component C_i is a binomial hypersurfaces, of type (t, ϵ_i) , $i = 1, \dots, s$, and condition 1. is verified.

Condition 2. now follows directly substituting $\alpha_i = \xi^{\epsilon_i}$ in condition (5.2). \square

Example 5.6. Let ξ be the 6-th root of unity $e^{\frac{2\pi i}{6}}$ and define the hypersurfaces

$$C_j = Z(h_j)$$

where $h_j = X^{I_1} - \xi^j X^{I_2}$, $j = 1, \dots, 6$, and X^{I_1} and X^{I_2} are coprime monomials of degree d .

According to Proposition 5.3 we have the following Hadamard multiplication table

	C_1	C_2	C_3	C_4	C_5	C_6
C_1	C_2	C_3	C_4	C_5	C_6	C_1
C_2	C_3	C_4	C_5	C_6	C_1	C_2
C_3	C_4	C_5	C_6	C_1	C_2	C_3
C_4	C_5	C_6	C_1	C_2	C_3	C_4
C_5	C_6	C_1	C_2	C_3	C_4	C_5
C_6	C_1	C_2	C_3	C_4	C_5	C_6

Consider $C = C_1 \cup C_3 \cup C_5$. If we compute C^{*2} , using the previous table, we get

$$(C_1 \cup C_3 \cup C_5)^{*2} = C_2 \cup C_4 \cup C_6.$$

Then, computing $(C_1 \cup C_3 \cup C_5)^{*3}$ we get

$$\begin{aligned} (C_1 \cup C_3 \cup C_5)^{*3} &= (C_2 \cup C_4 \cup C_6) \star (C_1 \cup C_3 \cup C_5) \\ &= C_1 \cup C_3 \cup C_5. \end{aligned}$$

Notice that Conditions 1. and 2. are satisfied for C_1, C_3, C_5 and $r = 3$.

6. ON HADAMARD POWERS OF VARIETIES

The result presentend in this section, generalizes the sufficient condition to have $V^{*2} = V$, in [9] and stated in Proposition 1.1 in the introduction of this paper, to the case of higher powers.

Similarly to the case of hypersurfaces, we define a specific class of varieties.

Definition 6.1. Let $C \subset \mathbb{P}^n$ be a binomial variety. We say that C is of type $[(t_1, \epsilon_1), \dots, (t_s, \epsilon_s)]$ if the ideal of C is generated by

$$X^{I_{1,1}} - \xi_1^{\epsilon_1} X^{I_{1,2}}, \dots, X^{I_{s,1}} - \xi_s^{\epsilon_s} X^{I_{s,2}},$$

where ξ_i is a primitive $(t_i - 1)$ -th root of unity and $1 \leq \epsilon_i \leq t_i - 1$ for $i = 1, \dots, s$.

Theorem 6.2. *Let $C \subset \mathbb{P}^n$ be a binomial variety of type $[(t_1, \epsilon_1), \dots, (t_s, \epsilon_s)]$ with*

$$(6.1) \quad \text{lcm} \left(\frac{t_1}{\gcd(t_1, \epsilon_1)}, \frac{t_2}{\gcd(t_2, \epsilon_2)}, \dots, \frac{t_s}{\gcd(t_s, \epsilon_s)} \right) = t - 1.$$

Then $C^{\star t} = C$ and t is the minimal exponent for which the previous equality holds.

Proof. By hypothesis the ideal of C is generated by $F_j(x_0, \dots, x_n) = X^{I_{j,1}} - \xi_j^{\epsilon_j} X^{I_{j,2}}$ for $j = 1, \dots, s$. Consider $t - 1$ points $P_i \in C \setminus \Delta_{n-1}$. Each point satisfies $P_i^{I_{j,1}} = \xi_j^{\epsilon_j} P_i^{I_{j,2}}$ and

$$(6.2) \quad \begin{aligned} (P_1 \star \dots \star P_{t-1})^{I_{j,1}} &= P_1^{I_{j,1}} \dots P_{t-1}^{I_{j,1}} \\ &= (\xi_j^{\epsilon_j})^{t-1} P_1^{I_{j,2}} \dots P_{t-1}^{I_{j,2}} \\ &= P_1^{I_{j,2}} \dots P_{t-1}^{I_{j,2}} \\ &= (P_1 \star \dots \star P_{t-1})^{I_{j,2}} \end{aligned}$$

where the third equality follows from the fact that $t - 1$ is the least integer such that $1 = (\xi_j^{\epsilon_j})^{t-1}$, for all $j = 1, \dots, s$ since (6.1) holds.

By Theorem 3.5, the equations for $P_1 \star \dots \star P_{t-1} \star C$ are

$$F_j^{\star(P_1 \star \dots \star P_{t-1})}(x_0, \dots, x_n) = \frac{X^{I_{j,1}}}{(P_1 \star \dots \star P_{t-1})^{I_{j,1}}} - \frac{\xi_j^{\epsilon_j} X^{I_{j,2}}}{(P_1 \star \dots \star P_{t-1})^{I_{j,2}}}$$

and, by (6.2), one has that $F_j^{\star(P_1 \star \dots \star P_{t-1})}(x_0, \dots, x_n)$ and $F_j(x_0, \dots, x_n)$ are equal up to a multiplicative constant, hence $P_1 \star \dots \star P_{t-1} \star C = C$. Since this holds for every choice of $P_1, \dots, P_{t-1} \in C \setminus \Delta_{n-2}$ and, by Remark 4.4, $P_1 \star \dots \star P_{t-1} \star C \subset C$ if $P_1 \star \dots \star P_{t-1} \in C \cap \Delta_{n-2}$, one has $C^{\star t} = C$. \square

REFERENCES

- [1] I. Bahmani Jafarloo and G. Calussi, *Weak Hadamard star configurations and apolarity*, Rocky Mountain J. Math. **50**(3) (2020), 851–862.
- [2] C. Bocci, E. Carlini and J. Kileel, *Hadamard Products of Linear Spaces*, J. of Algebra **448** (2016), 595–617.
- [3] C. Bocci, G. Calussi, G. Fatabbi and A. Lorenzini, *On Hadamard product of linear varieties*, Journal of Algebra and its Applications, **16**(8) (2017), 155–175.
- [4] C. Bocci, G. Calussi, G. Fatabbi and A. Lorenzini, *The Hilbert function of some Hadamard products*, Coll. Mathematica, **69**(2) (2018), 205–220.
- [5] G. Calussi, E. Carlini, G. Fatabbi, and A. Lorenzini, *On the Hadamard product of degenerate subvarieties*, Port. Math. **76**(2) (2019), 123–141.
- [6] E. Carlini, M. V. Catalisano, E. Guardo, and A. Van Tuyl, *Hadamard star configurations*, Rocky Mountain J. Math. **49**(2) (2019), 419–432.
- [7] M.A. Cueto, J. Morton and B. Sturmfels, *Geometry of the restricted Boltzmann machine*, Algebraic Methods in Statistics and Probability (eds. M. Viana and H. Wynn), American Mathematical Society, Contemporary Mathematics **516** (2010) 135–153.
- [8] M.A. Cueto, E.A. Tobis and J. Yu, *An implicitization challenge for binary factor analysis*, J. Symbolic Comput. **45**(12) (2010), 1296–1315.
- [9] N. Friedenberg, A. Oneto, R.L. Williams, *Minkowski sums and Hadamard products of algebraic varieties*, in Smith, G. G., Sturmfels, B. (edited by), Combinatorial Algebraic Geometry - Selected Papers From the 2016 Apprenticeship Program, New York: Springer New York LLC, 2017, p. 133–157.
- [10] J. Herzog, T. Hibi, H. Ohsugi, *Binomial Ideals*, Graduate Texts in Math., **279**, Springer (2018).

CRISTIANO BOCCI, DEPARTMENT OF INFORMATION ENGINEERING AND MATHEMATICS, UNIVERSITY OF SIENA, VIA ROMA, 56 SIENA, ITALY

Email address: `cristiano.bocci@unisi.it`

ENRICO CARLINI, DEPARTMENT OF MATHEMATICAL SCIENCES, POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI 24, TURIN, ITALY

Email address: `enrico.carlini@polito.it`