



Verification results for age-structured models of economic-epidemics dynamics

This is the peer reviewed version of the following article:

Original:

Fabbri, G., Gozzi, F., Zanco, G.A. (2021). Verification results for age-structured models of economic-epidemics dynamics. JOURNAL OF MATHEMATICAL ECONOMICS, 93, 1-11 [10.1016/j.jmateco.2020.102455].

Availability:

This version is available <http://hdl.handle.net/11365/1222417> since 2022-12-16T12:48:28Z

Published:

DOI: <http://doi.org/10.1016/j.jmateco.2020.102455>

Terms of use:

Open Access

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.

(Article begins on next page)

Journal of Mathematical Economics

Verification results for age-structured models of economic-epidemics dynamics

--Manuscript Draft--

Manuscript Number:	JME-D-20-00380R1
Article Type:	VSI: Economics of epidemics
Keywords:	COVID-19; macro-dynamic models; epidemiological dynamics; Hilbert spaces; verification theorem.
Corresponding Author:	Giorgio Fabbri Grenoble CEDEX 9, FRANCE
First Author:	Giorgio Fabbri
Order of Authors:	Giorgio Fabbri Fausto Gozzi Giovanni Zanco
Abstract:	<p>In this paper we propose a macro-dynamic age-structured set-up for the analysis of epidemics/economic dynamics in continuous time. The resulting optimal control problem is reformulated in an infinite dimensional Hilbert space framework where we perform the basic steps of dynamic programming approach.</p> <p>Our main result is a verification theorem which allows to guess the feedback form of optimal strategies. This will be a departure point to discuss the behavior of the models of the family we introduce and their policy implications.</p>

VERIFICATION RESULTS FOR AGE-STRUCTURED MODELS OF ECONOMIC-EPIDEMICS DYNAMICS

GIORGIO FABBRI, FAUSTO GOZZI, AND GIOVANNI ZANCO

ABSTRACT. In this paper we propose a macro-dynamic age-structured set-up for the analysis of epidemics/economic dynamics in continuous time.

The resulting optimal control problem is reformulated in an infinite dimensional Hilbert space framework where we perform the basic steps of dynamic programming approach.

Our main result is a verification theorem which allows to guess the feedback form of optimal strategies. This will be a departure point to discuss the behavior of the models of the family we introduce and their policy implications.

Keywords: COVID-19, macro-dynamic models, epidemiological dynamics, Hilbert spaces, verification theorem.

JEL Classification: E60, I10, C61.

1. INTRODUCTION

The outbreak of the COVID-19 pandemic represents, in addition to an epidemiological historical event, an exceptional economic shock. Data from the OECD (2020) suggest that in many countries the loss of GDP due to the presence of the virus and the consequent containment measures will be at least 10%. For this reason, together with the obvious upsurge in medical scientific production on the subject, the phenomenon has had great echo in the economic literature with a strong pressure to merge economic and epidemiological models.

A specific effort has been made to integrate epidemiological compartmental models (SIR, SEIR, SEI...) into a macroeconomic dynamic context, see for example the contributions of Alvarez et al., (2020), Eichenbaum et al., (2020), Jones et al., (2020) and Krueger et al.(2020).

These articles focus on a series of questions essential to health and economic policy and they look, often numerically, at the trade-off between measures capable of containing contagion and those capable of avoiding economic collapse. However, they model the spread of the epidemic with age homogeneous epidemiological compartmental models so they cannot take into account one of the characteristic traits

of the current epidemic, i.e. the great difference in the effects of the disease among people of different ages¹.

In order to address this limitation Acemoglu et al. (2020), Gollier (2020) and Favero et al. (2020) introduce models where the population is divided into a finite number of homogeneous “risk groups” and they study joint economic and epidemiological effects of introducing group-specific policies. Nonetheless in their formulations there is no possibility to move from one group to another and then this kind of approach can take into account the different effects of the disease on different age groups only if it is assumed that the duration of the epidemic is negligible compared to the age range contained in each group. However, this hypothesis is not very likely in the case of an epidemic lasting several years and it is inadequate in the case of diseases that become endemic in the population².

Instead of using age-homogeneous epidemiological compartmental models or epidemiological compartmental models with closed risk groups, it is possible, as we do in the present work, to describe more accurately the joint dynamics of the epidemic and of the age structure of the population by using explicit age-structured compartmental models, i.e. age-specific epidemiological models with ageing process modeled *à la* Mc Kendrick (1925). This type of models was initially introduced by Anderson and May (1985) and Dietz and Schenzle (1985) and later adapted to numerous contexts and applications, see the books by Iannelli (1995), Iannelli and Milner (2017) and Martcheva (2015) for a structured and modern description of the matter.

The more accurate description of the ageing-epidemics diffusion dynamics comes at a price and, indeed, one of the features of the continuous time compartmental age-structured models is to describe the epidemiological dynamics through transport type partial differential equations (PDEs). This means that, if one wants to study an associated optimal control problem through dynamic programming, its dynamics needs to be formulated in an infinite-dimensional set-up.

In this, paper whose main aim is methodological, we initially present a class of macro-dynamic models (partly already introduced in the literature) that incorporates an epidemiological dynamics which generalizes the benchmark age-structured SIR model. Then we provide a general framework to study such optimal control

¹In fact, the probability of aggravation of COVID-19 infection and mortality varies very significantly with age. Salje et al. (2020) find for example that less than 1% of people under 40 years of age who contract the disease need hospital care against more than 10% of people over 70 years of age and that mortality in the two groups is respectively less than 0.02% and more than 2%.

²These limitations are obviously justified by the need to produce policy indications in a short time in order to contrast the spread of the current pandemic.

problems through the dynamic programming approach; finally we present verification type results that hold for the whole class of problems.

The class of models that we study in the abstract form is rather general and is able, in the context of the epidemiological dynamic described by an age-structured SIR, to reproduce as special cases several of the settings proposed by the recent articles mentioned above (more details can be found in Section 2). Specific traits of the model are:

- The epidemiological model is general and can be set with a wide variety of age-dependent parameters: mortality rate (due to epidemics and also to other causes), chance of being hospitalized if infected, birth rate and probability of contagion among cohorts. Moreover, the age-specific mortality rate can take into account the saturation of hospitals and health systems, a phenomenon that has been repeatedly observed in the areas most affected by COVID-19 (see for instance Moghadas, 2020)
- The planner has two different policy levers: on the one hand, as in most of the models mentioned, she can reduce the mobility of people and partially stop the economic activity (lock-down), on the other hand she can implement some costly action to reduce the diffusion of the virus, for instance by testing the population extensively to try to quarantine individuals fast once they have contracted the virus. Both policies can be age-specific (in particular targeted lock-downs suggested by Acemoglu et al., 2020 are among possible policies)
- The time horizon can be either infinite (as in benchmark growth models) or finite if it is considered (e.g. Gollier, 2020) that the spread of the virus stops at some point due to the discovery of a vaccine or a cure
- Labor productivity is age-specific (this fact is important for policies: targeted lock-downs for less productive people impact less the production). The production function (as a function of aggregated labor and capital) is general as well as the optimal social function that can be specified to take into account cost-benefit analysis, strictly humanitarian or economic targets, standard (Benthamite and Millian for instance) social welfare functionals.

Since our results are proven for the abstract model they hold for any possible specification. Somehow related papers are Richard et al. (2020) and Colombo et al. (2020) (which also contains a spatial spread modeling of the virus diffusion) where the authors present models with a complete age structure but with a simple cost structure and no factor accumulation.

We also mention some works in the mathematical economic literature which study optimal control problems where the state equations are, as in the present work, of age-structured type: Barucci and Gozzi (1998 and 2001), Anita (2000), Ainseba et

al (2002), Feichtinger et al (2003, 2004 and 2006), Hritonenko and Yatsenko (2005), Veliov (2008), Boucekkine et al (2013), Skritek and Veliov (2015).

The contribution of this work is (i) to propose a general fully age-structured macro-dynamic set-up in continuous time for analysis of epidemics and economic dynamic (Section 2); (ii) to provide a suitable Hilbert space environment where one can rewrite the problem and perform dynamic programming (Section 3); (iii) to prove verification type results (Section 5), see in particular Theorem 5.2 and Corollaries 5.3 and 5.4.

We must be clear on the fact that we do not solve the problem explicitly, nor numerically. Here our main goal is to provide a general ground which can be the departure point to attack special cases of our general model. In particular our main contribution is the proof of the verification type results of Section 5. These are nontrivial to obtain in our general infinite dimensional setting and they are crucial to find the optimal policies in a closed-loop form depending on the derivatives of the value function. These type of theorems are the object of various papers (see e.g. Faggian and Gozzi (2010), Fabbri et al. (2010)) or of book chapters (e.g. Chapter 5 of Yong and Zhou, 1999 or Chapter 4 of Li and Yong, 1995) but none of them applies to our case. The main reasons are the following. First, due to the age-structured nature of the problem and the presence of the mortality forces, we have to work with semigroups in weighted infinite dimensional spaces which do not have regularizing properties (which are very useful and which are usually true when the state equation is of (nondegenerate) second order). Second, the presence of the nonlinear equation for capital rules out the standard regularity assumptions that are used e.g. in Chapter 4 of Li and Yong (1995) and that we treat using *ad-hoc* arguments. Third, in our case we have state constraints, which makes much more difficult to deal with the problem. We use the approach of weakening the constraints which has been used, up to now, only in case when explicit solutions of the HJB equation are available (see e.g. Fabbri and Gozzi, 2008, Boucekkine et al., 2019).

The paper is organized as follows. In Section 2 we introduce the structure of the model: epidemiological dynamics, policies, structure of the economy and welfare functional. In Section 3 we show how to reformulate the model and the related optimal control problem in a suitable Hilbert space setting. Section 4 is devoted to dynamic programming while in Section 5 we provide the verification results. Section 6 concludes.

2. THE MODEL

2.1. Epidemics dynamics. We denote by $s(a, t)$ the density of susceptible individuals of age $a \in [0, \bar{a}]$ (being $\bar{a} > 0$ the maximum age) at time $t \geq 0$. Similarly

$i(a, t)$ (respectively $r(a, t)$) denotes the density of infected/infectious individuals (respectively recovered individuals) of age a at time t . Hence the total numbers of susceptible, infected, and recovered individuals at time t are

$$S(t) = \int_0^{\bar{a}} s(a, t) da, \quad I(t) = \int_0^{\bar{a}} i(a, t) da, \quad R(t) = \int_0^{\bar{a}} r(a, t) da.$$

The age-dependent density of the total population $n(a, t)$ is then given by

$$n(a, t) = s(a, t) + i(a, t) + r(a, t)$$

and the total population at time t is

$$N(t) = \int_0^{\bar{a}} n(a, t) da = S(t) + I(t) + R(t).$$

In modeling the mortality we generalize the standard age-structure SIR framework (see Martcheva, 2015, Chapter 12). First we define

$$\Xi(t) := \int_0^{\bar{a}} i(a, t) \xi(a) da$$

the number of people at time t in “critical conditions” i.e. people who have to use the services of hospital/healthcare facilities to treat themselves at the risk of saturating them. In the case of the COVID-19 epidemic, the emphasis is, for example, on people needing to be hospitalized in an ICU (see for instance Moghadas, 2020). $\Xi(t)$ depends on the number of sick people per cohort multiplied by the prevalence $\xi(a)$ of people in need of specific care for each age group³. In a context of saturation of hospital services, the mortality of the infected will be increased. Let us therefore assume that the mortality rate for infected individual μ_I is not only a function of the age of the individuals but it also an increasing function of Ξ . We use then the notation $\mu_I(a, \Xi(t))$. We suppose, for simplicity, that the (age-specific) mortality rates of susceptible and recovered individuals, respectively $\mu_S(a)$ and $\mu_R(a)$ do not depend⁴ on $\Xi(t)$. Finally $\gamma(a)$ and $\beta(a)$ denote respectively the (age-specific) recovery and birth rates.

The age-specific force of infection $\lambda(a, t)$ depends on the distribution of infected individuals as follows

$$(1) \quad \lambda(a, t) = \frac{1}{N(t)} \int_0^{\bar{a}} m(a, \tau) i(\tau, t) d\tau.$$

In this expression the joint-distribution $m(a, \tau)$ measures the different probability of contagion between cohorts (for instance virus diffusion can be easier among children

³In the case of COVID-19 for example, in the data of Salje (2020), 2.9% of infected individuals are hospitalized ranging from 0.1% in people under 20 years to approximately 30% in individuals with 80 years of age or older.

⁴Indeed it is possible to incorporate this dependence in the model without big problems.

for childhood diseases). It is the continuous version of the social contact matrix across age classes used by Gollier (2020).

All in all, the *laissez faire* benchmark population dynamics, that is the epidemics dynamics without policy intervention (omitting the initial conditions at time $t = 0$) is the following:

$$(2) \quad \begin{cases} \frac{\partial s(a,t)}{\partial t} + \frac{\partial s(a,t)}{\partial a} = -\lambda(a,t)s(a,t) - \mu_S(a)s(a,t), \\ \frac{\partial i(a,t)}{\partial t} + \frac{\partial i(a,t)}{\partial a} = \lambda(a,t)s(a,t) - (\mu_I(a, \Xi(t)) + \gamma(a))i(a,t), \\ \frac{\partial r(a,t)}{\partial t} + \frac{\partial r(a,t)}{\partial a} = \gamma(a)i(a,t) - \mu_R(a)r(a,t) \\ s(0,t) = \int_0^{\bar{a}} \beta(a)n(a,t)da \\ i(0,t) = 0 \\ r(0,t) = 0. \end{cases}$$

This system is the standard age-structure SIR model (see Martcheva, 2015, Chapter 12) except for the fact that μ_I depends on Ξ . In the particular case where $\mu_I(a, \Xi(t)) = \tilde{\mu}_I(a)$ we are exactly in the standard setting. Note that, since λ and Ξ depend linearly on i , system (2) is non linear in the variables (s, i, r) .

We now introduce two of the three policy levers that the planner has in our model (the third is the choice of consumption and will be described in the next subsection). We suppose that the planner can deal with the epidemic in two ways:

- (i) partially stopping economic activity and people mobility then reducing the contagion frequency among individual (lockdown);
- (ii) implementing some costly action to reduce the diffusion of the virus, for instance by testing the population extensively to try to quarantine individuals faster once they have contracted the virus.

More precisely

- (i) We suppose that the planner can reduce mobility, thus the probability of infecting and being infected of cohort a at time t , by a factor $\theta(a, t) \in [0, 1]$ at the cost of reducing the contribution of the concerned individuals to work or of reducing their work productivity (for example resorting to teleworking). This is the type of intervention which is modeled in almost all the macro-dynamic models we mentioned in the introduction, for instance in Alvarez et al., (2020) and Eichenbaum et al., (2020) where, by the way, age-structure policies are not possible since there is no age structure of the population. Taking different values of θ for different a correspond to targeted lock-downs of Acemoglu et al. (2020).
- (ii) We suppose, as in some of the mentioned papers, that the planner can reduce by a factor $\eta(a, t) \in [0, 1]$ the probability that infected individuals of cohort

a at time t contaminate other people. This is done at the cost

$$(3) \quad D_\eta(t) := D \left(\int_0^{\bar{a}} (1 - \eta(a, t)) i(a, t) e(a) da \right).$$

where $e(a)$ is an age-specific relative cost and D is a concave (as, for instance in Piguillem and Shi, 2020) or linear (as in Gollier, 2020) function which represents some form of congestion (e.g. shortage of tests on the international market or shortage of suitable medical personnel to administer the tests).

The evolution of the epidemics is then again described by (2) but, instead of $\lambda(a, t)$ written in (1) we have now the following age-specific force

$$(4) \quad \lambda^{\theta, \eta}(a, t) = \frac{1}{N(t)} \int_0^{\bar{a}} m(a, \tau) \theta(\tau, t) \eta(\tau, t) i(\tau, t) d\tau.$$

Hence we get the following state equation (still omitting the initial conditions at time $t = 0$):

$$(5) \quad \begin{cases} \frac{\partial s(a, t)}{\partial t} + \frac{\partial s(a, t)}{\partial a} = -\lambda^{\theta, \eta}(a, t) s(a, t) - \mu_S(a) s(a, t), \\ \frac{\partial i(a, t)}{\partial t} + \frac{\partial i(a, t)}{\partial a} = \lambda^{\theta, \eta}(a, t) s(a, t) - (\mu_I(a, \Xi(t)) + \gamma(a)) i(a, t), \\ \frac{\partial r(a, t)}{\partial t} + \frac{\partial r(a, t)}{\partial a} = \gamma(a) i(a, t) - \mu_R(a) r(a, t) \\ s(0, t) = \int_0^{\bar{a}} \beta(a) n(a, t) da \\ i(0, t) = 0 \\ r(0, t) = 0. \end{cases}$$

This is the dynamics we will consider in the paper. Of course if the authority fixes $\theta(a, t) \equiv 1$, $\eta(a, t) \equiv 1$ we find again the free diffusion dynamics (2).

2.2. Production and capital accumulation. We suppose that labor supply is perfectly inelastic to wage, that infected people do not work and that labor productivity is age-specific⁵ and proportional to a certain parameter $\alpha(a)$ (we can specify for instance $\alpha(a) = 0$ for children or for individuals older than a fixed retirement age). Total labor supply in efficiency units in the *laissez faire* benchmark is then given by $\int_0^{\bar{a}} (s(a, t) + r(a, t)) \alpha(a) da$. In the controlled case we suppose that getting a factor $\theta(a, t) \in [0, 1]$ in the expression of the age-specific force of diffusion impacts the productivity of cohort a reducing the productivity to $\varphi(\theta(a, t))$ so that total

⁵We abstract from other reasons of productivity heterogeneity among population and from heterogeneity of tasks.

labor supply in efficiency units⁶ is now

$$(6) \quad L(t) = \int_0^{\bar{a}} (s(a, t) + r(a, t))\alpha(a)\varphi(\theta(a, t))da.$$

We suppose that $\varphi: [0, 1] \rightarrow [0, 1]$ is an increasing function with $\varphi(1) = 1$.

As for the production we stick to the standard structure of neoclassical growth models and we suppose that the total production at time t is described by an aggregated production function F of the two factors: labor $L(t)$ and capital $K(t)$:

$$Y(t) = F(K(t), L(t)).$$

This formulation is more general than the one used by other macro-dynamic papers we mentioned. Indeed in all of them except Favero et al. (2020) which uses a Cobb-Douglas production function, the authors use production functions which are linear function of labor (or effective labor) or they even do not model production.

We abstract from international trade (closed economy) and from governmental expenditure so the planner can choose at any time $t \geq 0$ how to allocate the national total production $Y(t)$ among total investment $M(t)$, consumption of various cohorts and costs for testing people, which is defined in (3) above. If we denote by $c(a, t)$ the per-capita consumption of individuals of age a at time t we get the following budget constraint:

$$Y(t) = M(t) + C(t) + D_\eta(t) := M(t) + \int_0^{\bar{a}} c(a, t)n(a, t)da + D \left(\int_0^{\bar{a}} (1 - \eta(a, t))i(a, t)e(a)da \right).$$

Supposing to have an exponential capital depreciation *à la* Jorgenson we get the dynamic accumulation law for capital:

$$(7) \quad \dot{K}(t) = F(K(t), L(t)) - \int_0^{\bar{a}} c(a, t)n(a, t)da - \delta K(t) - D \left(\int_0^{\bar{a}} (1 - \eta(a, t))i(a, t)e(a)da \right).$$

where $\delta > 0$ is the constant depreciation rate.

2.3. Choosing the target. In the literature there are several interesting choices for the functional to maximize. It is not easy to include all of them in an abstract form that leaves the problem tractable, so in this section we introduce several functionals that will be discussed later in the article.

The first functional we introduce is a standard welfare functional. Observe that, even if the model we study here is not directly an endogenous fertility model, the fact of having an endogenous mortality (depending on the choice of θ) makes it *de facto* an endogenous population size model. Therefore we have to choose carefully

⁶A similar approach is considered for instance by Jones et al. (2020) which introduce an “effective labor supply”.

the structure of the social utility that we describe. We implicitly fix the utility of dead people (and non-born people through the initial condition $\int_0^{\bar{a}} \beta(a)n(a,t)da$) equal to 0 and we consider the following social utility functional:

$$(8) \quad \int_0^\infty \int_0^{\bar{a}} e^{-\rho t} n^\nu(a,t) u(c(a,t), \theta(a,t)) da dt.$$

To assure, for the same per capita (age-dependent) consumption, the instantaneous utility to be increasing in the number of living people and therefore the planner being averse to death of agents, the per-capita utility function u needs to be positive. Still observe that in this model formulation there is room for a dilution effect: the larger the population the lower the percapita consumption so the instantaneous utility does not need to always be increasing in the population size.

The per-capita utility function u depends both on the individual consumption and on the mobility freedom θ . We suppose that u is (positive and) an increasing function in both the variables. The dependence of utility on θ is not standard but the relevance of this choice can easily be argued by looking at the various side effects of lock down (see for example Clemens, 2020). In any case, as a special case, of course one can specify u so that it does not depend on theta.

The form of this first functional is the age-structured version of a standard functional often appearing in the optimal population literature. The parameter ν which appears in its expression measures the degree of altruism towards individuals of future cohorts (see Palivos and Yip, 1993). The case $\nu = 1$ corresponds to the classical total utilitarianistic (or ‘‘Benthamite’’) case where the planner target is to maximize the sum of individuals’ utility.

The functional (8) is infinite horizon and implicitly suggests that no exogenous element impedes the spread of the virus. Another possibility, as suggested by Gollier (2020), is to consider a final time T at which an event (a cure or more probably the discovery of a vaccine) stops the epidemics. We describe some possible targets in this context.

The trade-off of virus containment policies is: reducing the number of deaths VS economic losses. Some of the functional aspects can be dwelt on only one of these aspects. For instance one can decide to focus on economic activity and to maximize the final production capacity:

$$(9) \quad F(K(T), \tilde{L}(T))$$

where $\tilde{L}(T)$ is defined as

$$(10) \quad \tilde{L}(T) = \int_0^{\bar{a}} n(a,T)\alpha(a)da$$

(once the vaccine is in and the outbreak is over, everyone is cured and everyone is productive) or even more simply, to maximize final capital level

$$(11) \quad K(T)$$

or to maximize the flow of production

$$(12) \quad \int_0^{+\infty} e^{-\rho t} Y(t) dt.$$

or its finite-horizon counterpart

$$(13) \quad \int_0^T e^{-\rho t} Y(t) dt.$$

Conversely, in the spirit of Acemoglu et al. (2020), one can include humanitarian aspects by minimizing the weighted sum of number of deaths due to the virus and of the opposite of the total discounted output:

$$(14) \quad \chi \int_0^T \int_0^{\bar{a}} \mu_I(a, \Xi(t)) i(a, t) da - \int_0^T e^{-\rho t} Y(t) dt.$$

where $\chi > 0$ is the relative weight of deaths.

3. INFINITE DIMENSIONAL FORMULATION OF THE MODEL

In this section we introduce a convenient infinite-dimensional formulation for system (5) coupled with equation (7) and for the control problem of maximizing the target given in (8) (or in (12), (9), (11), (13)). Since this section is somehow technical we add, where possible, explanations on the technical notation we use.

We make the following set of assumptions, which also includes those already stated in the previous sections. These assumptions will always be in force in the remainder of the paper, without recalling them.

Hypothesis 3.1

(i) μ_S and μ_R are positive, belong to $L^1_{loc}(0, \bar{a})$ ⁷ and

$$\int_0^{\bar{a}} \mu_S(a) da = \int_0^{\bar{a}} \mu_R(a) da = +\infty;$$

(ii) $\mu_I: [0, \bar{a}] \times \mathbb{R} \rightarrow \mathbb{R}_+$ is measurable. Moreover it is Lipschitz continuous in the second variable, uniformly with respect to the first one. Finally it is increasing in the second variable and

$$\int_0^{\bar{a}} \mu_I(a, \kappa) da = +\infty, \quad \forall \kappa \in \mathbb{R};$$

⁷ $L^1_{loc}(a, b)$ is the vector space of functions f that are Lebesgue-integrable on every $[a', b'] \subset (a, b)$.

- (iii) $F(\cdot, L)$ is Lipschitz for every $L \in \mathbb{R}_+$, with Lipschitz constants uniformly bounded in L ⁸;
- (iv) $\varphi: [0, 1] \rightarrow [0, 1]$ is increasing and $\varphi(1) = 1$;
- (v) $\alpha, \beta, \gamma, e, \xi: [0, \bar{a}] \rightarrow \mathbb{R}_+$ are in $L^2(0, \bar{a})$ ⁹;
- (vi) $D: \mathbb{R} \rightarrow \mathbb{R}$ is positive and concave;
- (vii) $\delta > 0, \nu \in [0, 1]$;
- (viii) $u: \mathbb{R} \times [0, 1] \rightarrow \mathbb{R}$ is positive, continuous and increasing in both variables;

We now start rewriting the system (5) but first we introduce an important notational standard.

Notation 3.2 In system (5) the three state trajectories, $s(\cdot, \cdot)$, $i(\cdot, \cdot)$, $r(\cdot, \cdot)$ are seen as functions of two variables, i.e.

$$(s, i, r)(\cdot, \cdot): [0, +\infty) \times [0, \bar{a}] \rightarrow \mathbb{R}^3, \quad (t, a) \rightarrow (s(t, a), i(t, a), r(t, a))$$

However now it is convenient to see such trajectories as functions from $t \in \mathbb{R}_+$ to a suitable infinite dimensional Hilbert space H of functions in the variable $a \in [0, \bar{a}]$ with values in \mathbb{R}^3 . H can be seen also as the product of three Hilbert spaces of functions with values in \mathbb{R} and its generic element will be denoted by $h = (h_1, h_2, h_3)$ or, if no confusion arises, by (s, i, r) . To avoid misunderstandings we will denote the H -valued state trajectories putting a hat over the original name, i.e.

$$(\hat{s}, \hat{i}, \hat{r}): \mathbb{R}_+ \rightarrow H \quad t \rightarrow (\hat{s}(t), \hat{i}(t), \hat{r}(t))$$

Sometimes we will write \hat{h} for $(\hat{s}, \hat{i}, \hat{r})$ and, when we want to underline that they are functions, we write $\hat{h}(\cdot)$ or $(\hat{s}(\cdot), \hat{i}(\cdot), \hat{r}(\cdot))$. Now observe that, for every $t \geq 0$, $\hat{s}(t), \hat{i}(t), \hat{r}(t)$ are functions of a . We will denote their value at any given $a \in [0, \bar{a}]$ by $\hat{s}(t)[a], \hat{i}(t)[a], \hat{r}(t)[a]$ so to emphasize the different role of the two variables. Clearly we have

$$\hat{s}(t)[a] = s(t, a), \quad \hat{i}(t)[a] = i(t, a), \quad \hat{r}(t)[a] = r(t, a).$$

The same will be done for the controls strategies $c(\cdot, \cdot), \theta(\cdot, \cdot), \eta(\cdot, \cdot)$. More precisely, we will fix a control space Z of functions in the variable $a \in [0, \bar{a}]$ with values in \mathbb{R}^3 . Also Z can be seen as the product of three Hilbert spaces of functions with values in \mathbb{R} and its generic element will be denoted by $z = (z_1, z_2, z_3)$ or, if no confusion is possible, by (c, θ, η) . Also here, to avoid misunderstandings, we will denote the control trajectories putting a hat over the original name, i.e. we call $\hat{c}, \hat{\theta}, \hat{\eta}$ the

⁸meaning that for every $L \in \mathbb{R}_+$ there exists $\kappa_L \geq 0$ such that for every $K_1, K_2 \in \mathbb{R}_+$ $|F(K_1, L) - F(K_2, L)| \leq \kappa_L |K_1 - K_2|$ and $\sup_{L \in \mathbb{R}_+} \kappa_L < +\infty$.

⁹ $L^2(a, b)$ is the Hilbert space of square-Lebesgue-integrable functions on (a, b) , endowed with the inner product $\langle f, g \rangle_{L^2} = \int_a^b f(x)g(x)dx$.

functions

$$(\hat{c}, \hat{\theta}, \hat{\eta}) : \mathbb{R}_+ \rightarrow Z \quad t \rightarrow (\hat{c}(t), \hat{\theta}(t), \hat{\eta}(t))$$

such that

$$\hat{c}(t)[a] = c(t, a), \quad \hat{\theta}(t)[a] = \theta(t, a), \quad \hat{\eta}(t)[a] = \eta(t, a).$$

Sometimes we write \hat{z} for $(\hat{c}, \hat{\theta}, \hat{\eta})$ and, when we want to underline that they are functions we write $\hat{z}(\cdot)$ or $(\hat{c}(\cdot), \hat{\theta}(\cdot), \hat{\eta}(\cdot))$.

We are now ready to introduce the spaces H , Z , and the space \mathcal{Z}_0 of basic control strategies (this is not the space of admissible control strategies which will have to take into account the state constraints that we will introduce below).

Define the probability of surviving to age a for a susceptible individual as

$$\pi_S(a) = \exp\left(-\int_0^a \mu_S(\tau) d\tau\right)$$

and, similarly, define the probability of surviving to age a for a recovered individual as

$$\pi_R(a) = \exp\left(-\int_0^a \mu_R(\tau) d\tau\right).$$

Consider the set

$$H = \left\{ h \in L^2(0, \bar{a}; \mathbb{R}^3) : \frac{h_1}{\pi_S} \in L^2(0, \bar{a}), h_2 \in L^2(0, \bar{a}), \frac{h_3}{\pi_R} \in L^2(0, \bar{a}) \right\}.$$

H is a Hilbert space when endowed with the inner product

$$\begin{aligned} \langle h, g \rangle_H &= \left\langle \frac{h_1}{\pi_S}, \frac{g_1}{\pi_S} \right\rangle_{L^2} + \langle h_2, g_2 \rangle_{L^2} + \left\langle \frac{h_3}{\pi_R}, \frac{g_3}{\pi_R} \right\rangle_{L^2} \\ &=: \langle h_1, g_1 \rangle_{\pi_S} + \langle h_2, g_2 \rangle_{L^2} + \langle h_3, g_3 \rangle_{\pi_R}. \end{aligned}$$

Remark 3.3 The choice of the space H is different from the standard one made, e.g., in Iannelli and Martcheva, 2003. Indeed it would be standard to put the weight also on the second component. However this is not possible since, in our model, we have the new and important feature that the mortality force μ_I is state dependent. With our choice the space H is bigger than the usual one. We finally observe that this choice will reflect also in the form of the adjoint operator in Proposition 3.4.

It is useful for later purposes to introduce the positive cone in H as follows

$$H_+ = \{h \in H : h_i \geq 0 \text{ a.e. in } [0, \bar{a}]\} \subset H.$$

The control space Z is given as:

$$\begin{aligned} Z = \{z = (z_1, z_2, z_3) : z_i \in L^2(0, \bar{a}), i = 1, 2, 3; \\ z_1(a) \geq 0, z_2(a), z_3(a) \in [0, 1], \forall a \in [0, \bar{a}]\}. \end{aligned}$$

Z is a Hilbert space when endowed with the inner product

$$\langle z, w \rangle_Z = \langle z_1, w_1 \rangle_{L^2} + \langle z_2, w_2 \rangle_{L^2} + \langle z_3, w_3 \rangle_{L^2}.$$

Finally the space \mathcal{Z}_0 of all basic control strategies is, coherently with the requirements of Section 2, a space of functions from \mathbb{R}_+ to Z and is chosen as

$$\mathcal{Z}_0 := L^2(\mathbb{R}_+; Z);$$

For coherence with Notation 3.2 we will denote by $z = (c, \theta, \eta)$ the points of Z and by $\hat{z} = (\hat{c}, \hat{\theta}, \hat{\eta})$ the points of \mathcal{Z}_0 .

Now we reformulate system (5) providing also existence and uniqueness of the solution. We need to introduce some operators which originate from the various term of the system.

First we introduce the unbounded linear operator $A : D(A) \subset H \rightarrow H$ defined as

$$A = \begin{pmatrix} -\frac{\partial}{\partial a} - \mu_S & 0 & 0 \\ 0 & -\frac{\partial}{\partial a} - \gamma & 0 \\ 0 & \gamma & -\frac{\partial}{\partial a} - \mu_R \end{pmatrix}$$

with domain

$$D(A) = \left\{ h \in H : \frac{h_1}{\pi_S}, h_2, \frac{h_3}{\pi_R} \in W^{1,2}(0, \bar{a}), \right. \\ \left. h_1(0) = \int_0^{\bar{a}} \beta(a)(h_1 + h_2 + h_3)(a) da, h_2(0) = h_3(0) = 0 \right\}^{10},$$

corresponding to the linear part of system (5). It can be shown as in Iannelli and Martcheva (2003) that A generates a strongly continuous semigroup¹¹ $T(t)$ on H such that $T(t)(H_+) \subset (H_+)$ for every $t \geq 0$.

Second we define linear functional $\bar{\Xi}$ which reformulates the function Ξ given in Section 2.

$$(15) \quad \bar{\Xi} : H \rightarrow \mathbb{R}$$

$$(16) \quad \bar{\Xi}(h) = \int_0^{\bar{a}} h_2(a) \xi(a) da.$$

¹¹A semigroup of operators on H is a family of operators $T(t)$ on H indexed by $t \geq 0$ such that $T(t) + T(s) = T(t + s)$ and $T(0)$ is the identity; a semigroup $T(t)$ is strongly continuous if $\|T(t)h - h\|_H \rightarrow 0$ as $t \rightarrow 0$ for every $h \in H$.

For every control point $z = (c, \theta, \eta) \in Z$ we define the nonlinear operators (depending only on the components θ and η of the control point)

$$\Lambda^{\theta, \eta}: H_+ \setminus \{0\} \rightarrow L^\infty(0, \bar{a})^{12},$$

$$\Lambda^{\theta, \eta}(h)(a) = \frac{1}{\int_0^{\bar{a}} (h_1 + h_2 + h_3)(\tau) d\tau} \int_0^{\bar{a}} m(a, \tau) \theta(\tau) \eta(\tau) h_2(\tau) d\tau,$$

and

$$B^{\theta, \eta}: H_+ \setminus \{0\} \rightarrow H,$$

$$B^{\theta, \eta}(h)(a) = \begin{pmatrix} -\Lambda^{\theta, \eta}(h)(a) h_1(a) \\ \Lambda^{\theta, \eta}(h) h_1(a) - \mu_I(a, \Xi(h)) h_2(a) \\ 0 \end{pmatrix}.$$

Now by Hypothesis 3.1(ii) and by the fact that we have $\theta, \eta \in L^\infty(0, \bar{a})$ (see the definition of Z), the operator $B^{\theta, \eta}$ is Lipschitz continuous on $H_+ \setminus \{0\}$ and there exists a positive constant α such that $\alpha B^{\theta, \eta}(h) + h \in H_+$ for every $h \in H_+$ (see Iannelli and Martcheva, 2003).

We can consequently write system (5) as the evolution equation for the unknown $\hat{h}: [0, +\infty) \rightarrow D(A)$ with control strategies $\hat{\theta}$ and $\hat{\eta}$:

$$(17) \quad \frac{d}{dt} \hat{h}(t) = A \hat{h}(t) + B^{\hat{\theta}(t), \hat{\eta}(t)}(\hat{h}(t)).$$

Given control strategies $\hat{\theta}(\cdot), \hat{\eta}(\cdot)$ and an initial condition $h_0 \in H_+$ we look for mild solutions of the above systems in H_+ , i.e., for functions $[0, +\infty) \ni t \mapsto \hat{h}(t) \in H_+$ that satisfy

$$\hat{h}(t) = T(t) h_0 + \int_0^t T(t-s) B^{\hat{\theta}(s), \hat{\eta}(s)}(\hat{h}(s)) ds$$

Thanks to the fact that A generates a strongly continuous semigroup that leaves H_+ invariant and thanks to the properties of B , there exists (see e.g. Bensoussan et al. (2007)) a unique function $\hat{h}(\cdot)$ that satisfies (17) and such that $\hat{h}(0) = h_0$ and $\hat{h}(t) \in H_+$ for every $t \in [0, +\infty)$. Such solution will be denoted by $\hat{h}^{\hat{\theta}, \hat{\eta}; h_0}$ or by $\hat{h}^{\hat{z}; h_0}$.

We now add the equation for K to the system, see (7). For control points $z = (c, \theta, \eta) \in Z$ we define the functionals on H

$$L^\theta(h) = \int_0^{\bar{a}} (h_1(a) + h_3(a)) \alpha(a) \varphi(\theta(a)) da,$$

$$C^c(h) = \int_0^{\bar{a}} c(a) (h_1 + h_2 + h_3)(a) da,$$

$$D^\eta(h) = D \left(\int_0^{\bar{a}} (1 - \eta(a)) h_2(a) e(a) da \right).$$

For any control strategy $\hat{z}(\cdot) \in \mathcal{Z}_0$, any $h_0 \in H_+$ and any $K_0 \in \mathbb{R}$ we are then considering the Cauchy problem

$$(18) \quad \begin{cases} \hat{h}'(t) &= A\hat{h}(t) + B^{\hat{\theta}(t), \hat{\eta}(t)}(\hat{h}(t)) \\ K'(t) &= -\delta K + F\left(K(t), L^{\hat{\theta}(t)}(\hat{h}(t))\right) - C^{\hat{c}(t)}(\hat{h}(t)) - D^{\hat{\eta}(t)}(\hat{h}(t)), \\ \hat{h}(0) &= h_0, \\ K(0) &= K_0. \end{cases}$$

Observe that the first equation does not depend on K . Hence, once we know the mild solution $h^{\hat{z}; h_0}$ of the first equation we can plug it into the second one. Since F is Lipschitz in K , uniformly in the second variable¹³, we know that the second equation has a solution $K^{\hat{z}; K_0, h_0}$ (that depends also on h_0 since the trajectory \hat{h} appears in the second equation). We then conclude that, for every $\hat{z} = (\hat{c}, \hat{\theta}, \hat{\eta}) \in \mathcal{Z}_0$ the above system admits a unique solution $(h^{\hat{z}; h_0}, K^{\hat{z}; K_0, h_0})$ such that $h^{\hat{z}; h_0}$ is the mild solution of the first equation with initial datum h_0 and $h(t) \in H_+$ for every $t \geq 0$. We can write system (18) in a more compact way as

$$(19) \quad \begin{cases} \frac{d}{dt}(\hat{h}, K)(t) &= \tilde{A}(\hat{h}(t), K(t)) + \tilde{B}^{\hat{z}(t)}(\hat{h}(t), K(t)), \quad t \geq 0 \\ (\hat{h}, K)(0) &= (h_0, K_0) \in H_+ \times \mathbb{R} \end{cases}$$

where $\tilde{A}: D(A) \times \mathbb{R} \rightarrow H \times \mathbb{R}$ is the linear operator defined by

$$\tilde{A}(h, K) = (Ah, -\delta K)$$

and, for $z = (c, \theta, \eta) \in Z$, $\tilde{B}^z = \tilde{B}^{c, \theta, \eta}: H \times \mathbb{R} \rightarrow H \times \mathbb{R}$ is given by

$$\tilde{B}^z(h, K) = \left(B^{\theta, \eta}(h), F(K, L^\theta(h)) - C^c(h) - D^\eta(h) \right).$$

The following proposition follows from basic material in Iannelli and Martcheva (2003), Iannelli (1995), Bensoussan et al. (2007).

Proposition 3.4 *The linear operator \tilde{A} generates a strongly continuous semigroup $\tilde{T}(t)$ on $H \times \mathbb{R}$ that leaves $H_+ \times \mathbb{R}$ invariant, while the operator B^z is Lipschitz. The Cauchy problem (19) admits a unique mild solution, that coincides with that of (18).*

The adjoint operator of \tilde{A} with respect to the inner product

$$\langle (h, K), (p, Q) \rangle_{H \times \mathbb{R}} := \langle h, p \rangle_H + KQ,$$

is the linear operator $\tilde{A}^: (D(A^*) \times \mathbb{R}) \rightarrow H \times \mathbb{R}$ given by*

$$\tilde{A}^*(p, Q) = (A^*p, -\delta Q),$$

¹³The fact the two equations are not fully coupled can be exploited to cover also the case when F is a Cobb-Douglas function by using a Bernoulli-type change of variable.

where

$$A^* = \begin{pmatrix} \frac{\partial}{\partial a} + \mu_s & 0 & 0 \\ 0 & \frac{\partial}{\partial a} - \gamma & \frac{\gamma}{\pi_R} \\ 0 & 0 & \frac{\partial}{\partial a} + \mu_R \end{pmatrix}$$

and

$$D(A^*) = \left\{ p = (p_1, p_2, p_3) : \frac{p_1}{\pi_S}, p_2, \frac{p_3}{\pi_R^2} \in W^{1,2}(0, \bar{a}), \right. \\ \left. \frac{p_1}{\pi_S}(\bar{a}) = p_2(\bar{a}) = \frac{p_3}{\pi_R}(\bar{a}) = p_1(0) = 0 \right\}.$$

Now we are in position to define precisely the set of admissible control strategies and to rewrite the target functionals. First of all, due to the presence of positivity constraints both on \hat{h} and K the set of admissible control strategies depends on the initial data and is the set

$$(20) \quad \mathcal{Z}_{ad}(h_0, K_0) = \left\{ \hat{z}(\cdot) \in \mathcal{Z} : \left(h^{\hat{z}; h_0}, K^{\hat{z}; K_0, h_0} \right) (t) \in H_+ \times \mathbb{R}_+ \text{ for a.e. } t \in [0, +\infty) \right\}.$$

We now rewrite the target functionals beginning with (8). Define the function

$$J_1 : H_+ \times \mathbb{R} \times \mathcal{Z}_0 \rightarrow \mathbb{R},$$

$$(21) \quad J_1(h_0, K_0; \hat{z}) \\ = \int_0^\infty e^{-\rho t} \int_0^{\bar{a}} \left(\hat{h}_1^{\hat{z}; h_0}(t)[a] + \hat{h}_2^{\hat{z}; h_0}(t)[a] + \hat{h}_2^{\hat{z}; h_0}(t)[a] \right)^\gamma u \left(\hat{c}(t)[a], \hat{\theta}(t)[a] \right) da dt.$$

Then for every given initial datum $(h_0, K_0) \in \mathbb{H}_+ \times \mathbb{R}_+$ the problem of maximizing (8) in Section 2 translates precisely in maximization of the function $J_1(h_0, K_0; \cdot)$ given by (21) over the set $\mathcal{Z}_{ad}(h_0, K_0)$. It will be useful, as a shorthand, to define the function $U : H_+ \times Z \rightarrow \mathbb{R}$,

$$(22) \quad U_1(h; z) = U_1(h; c, \theta) = \int_0^{\bar{a}} (h_1(a) + h_2(a) + h_3(a))^\gamma u(c(a), \theta(a)) da,$$

so that

$$(23) \quad J_1(h_0, K_0; \hat{z}) = \int_0^\infty e^{-\rho t} U_1 \left(h^{\hat{z}; h_0}(t); \hat{c}(t), \hat{\theta}(t) \right) dt.$$

The other infinite horizon problem of Section 2 have the same set of admissible strategies, hence to define it precisely it is enough the rewrite the corresponding target functional. To do this we simply have to change the function J_1 . The target

(12) can be rewritten defining the functional J_2 in the form (23) simply substituting U_1 from (22) with U_2 defined as (recall the definition of $L(\cdot)$ given in (6))

$$(24) \quad U_2(h, K; \theta) = F \left(K, \int_0^{\bar{a}} (h_1(a) + h_3(a)) \alpha(a) \varphi(\theta(a)) da \right).$$

The *value functions* of the two maximization problems described above are defined as

$$(25) \quad V_i(h, K) := \sup_{\hat{z} \in \mathcal{Z}_{ad}(h, K)} J_i(h, K; \hat{z}), \quad i = 1, 2.$$

The other four functionals of Section 2.3 are taken with finite horizon $T > 0$. It is useful, to apply the dynamic programming approach, to let also the initial time vary. Hence, when studying these targets the initial condition of the state equation (19) is taken at a generic time $t_0 \in [0, T]$ and the state equation itself takes the form

$$(26) \quad \begin{cases} \frac{d}{dt}(\hat{h}, K)(t) &= \tilde{A}(\hat{h}(t), K(t)) + \tilde{B}^{\hat{z}(t)}(\hat{h}(t), K(t)), \quad t \in [t_0, T], \\ (\hat{h}, K)(t_0) &= (h_0, K_0) \in H_+ \times \mathbb{R}; \end{cases}$$

its solution (which exists and is unique thanks to Proposition 3.4) is denoted by $(h^{\hat{z}; t_0, h_0}, K^{\hat{z}; t_0, K_0, h_0})$. The set of admissible control consequently becomes

$$(27) \quad \begin{aligned} &\mathcal{Z}_{ad}(t_0, h_0, K_0) \\ &= \left\{ \hat{z}(\cdot) \in \mathcal{Z} : \left(h^{\hat{z}; t_0, h_0}, K^{\hat{z}; t_0, K_0, h_0} \right)(t) \in H_+ \times \mathbb{R}_+ \text{ for a.e. } t \in [t_0, T] \right\}. \end{aligned}$$

For the same reason also the lower limit of the integral in the target is set at a generic time $t_0 \in [0, T]$. Therefore both the the target functional and the value function depend also on t_0 .

To rewrite target (9) we define

$$(28) \quad \begin{aligned} &J_3(t_0, h_0, K_0; \hat{z}) = \\ &F \left(K^{\hat{z}; t_0, h_0, K_0}(T), \int_0^{\bar{a}} \left(\hat{h}_1^{\hat{z}; t_0, h_0}(T)[a] + \hat{h}_2^{\hat{z}; t_0, h_0}(T)[a] + \hat{h}_3^{\hat{z}; t_0, h_0}(T)[a] \right) \alpha(a) da \right) \end{aligned}$$

Target (11) can be rewritten as

$$(29) \quad \hat{J}_4(t_0, h_0, K_0; \hat{z}) = K^{\hat{z}; t_0, h_0, K_0}(T).$$

While the above two targets only contain a final reward, the next two contain only a current reward. To rewrite target (13) we set

$$(30) \quad J_5(h_0, K_0; \hat{z}) = \int_0^T e^{-\rho t} U_2 \left(h^{\hat{z}; h_0}(t), K^{\hat{z}; t_0, K_0, h_0}(t); \hat{\theta}(t) \right) dt.$$

where U_2 is defined in (24). Eventually to rewrite target (14) we define

$$(31) \quad J_6(t_0, h_0, K_0; \hat{z}) = \chi \int_{t_0}^T \int_0^{\bar{a}} \mu_I \left(a, \Xi(\hat{h}_2^{\hat{z}; t_0, h_0}(t)) \right) \hat{h}_2^{\hat{z}; t_0, h_0}(t)[a] da dt \\ - \int_0^T e^{-\rho t} U_2 \left(h^{\hat{z}; h_0}(t), K^{\hat{z}; t_0, K_0, h_0}(t); \hat{\theta}(t) \right) dt.$$

or simply

$$J_6(t_0, h_0, K_0; \hat{z}) = \int_{t_0}^T U_3 \left(t, h^{\hat{z}; h_0}(t), K^{\hat{z}; t_0, K_0, h_0}(t); \hat{\theta}(t) \right) dt,$$

where

$$(32) \quad U_3(t, h, K; \theta) = \chi \int_0^{\bar{a}} \mu_I(a, \Xi(h_2)) h_2(a) da - e^{-\rho t} U_2(h, K; \theta).$$

Note that here above $\Xi : L^2(0, \bar{a}) \rightarrow \mathbb{R}$ is the linear functional given by $\Xi(h_2) = \int_0^{\bar{a}} h_2(a) \xi(a) da$, as in (15). The *value functions* in the above four finite horizon cases are defined as:

$$V_i(t, h, K) := \sup_{\hat{z} \in \mathcal{Z}_{ad}(t, h, K)} J_i(t, h, K; \hat{z}), \quad i = 3, 4, 5, 6.$$

4. DYNAMIC PROGRAMMING AND HJB EQUATIONS

The starting point of the dynamic programming approach to the problems of this paper is the Dynamic Programming Principle, which we call DPP from now on, (see e.g. Theorem 1.1, p. 224 of Li and Yong, 1995, for a statement and a proof which apply to the present case) which is a functional equation for the value function. Once DPP is established the standard path is to write the differential form of DPP, the HJB equation, find a solution v of it, and prove a Verification Theorem i.e. a sufficient condition for optimality in terms of the function v (which can be then proved to be the value function) and its derivatives. Both steps may be very complicated, depending on the features of the problem; this is particularly true when one deals with problems in infinite dimension. Indeed, while for finite dimensional problems the theory of HJB equations and of the corresponding verification results is quite well established with many regularity results, this is not the case for infinite dimensional problems. Indeed only few results are available and each case must be treated ad hoc. One can see, for example, Theorem 5.5, p.263 of Li and Yong (1995) and the papers Faggian and Gozzi (2010), Fabbri et al. (2010).

Here we abstract away from the existence and uniqueness of regular solutions of the HJB equation (which is a challenging subject and which will be next step of our work) and we concentrate on Verification Theorems and their consequences.

Since we formulated various different problems with different targets, here we concentrate on targets (8) and (14); the results can be easily extended to the other cases.

Consider first the problem, given any initial datum $(h_0, K_0) \in \mathbb{H}_+ \times \mathbb{R}_+$, of maximizing target (8) over all $\hat{z} \in \mathcal{Z}_{ad}(h_0, K_0)$. Formally, the Hamilton-Jacobi-Bellman (HJB) equation associated to such control problem is (the unknown here is $v : H_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$)

$$(33) \quad \rho v(h, K) = \sup_{z \in Z} \mathbb{H}_{CV}(h, K, D_h v(h, K), D_K v(h, K); z)$$

where the so-called Current Value Hamiltonian is defined as

$$\mathbb{H}_{CV} : ((D(A) \cap H_+) \times \mathbb{R}) \times (H \times \mathbb{R}) \times Z \rightarrow \mathbb{R}.$$

$$(34) \quad \mathbb{H}_{CV}(h, K, p, Q; z) = \langle \tilde{A}(h, K), (p, Q) \rangle_{H \times \mathbb{R}} + \langle \tilde{B}^z(h, K), (p, Q) \rangle_{H \times \mathbb{R}} + U_1(h; z)$$

However this form of the HJB equation is not very convenient for two main reasons.

- First of all the unknown is defined only in $H_+ \times \mathbb{R}_+$, because U_1 is defined in H_+ . This is a serious problem since the set H_+ has empty interior in H and this yields issues when defining properly the Fréchet derivative $D_h v$. To overcome this difficulty we observe that the formula defining U_1 makes sense as long as $h_1 + h_2 + h_3 \geq 0$. Therefore we can extend U_1 to the half space

$$(35) \quad H_+^1 := \{h \in H : \langle h, \mathbf{1} \rangle \geq 0\}$$

(here $\mathbf{1}$ is the function with constant value $(1, 1, 1)$ on H). Indeed the interior part of this set in H is simply

$$\text{Int } H_+^1 := \{h \in H : \langle h, \mathbf{1} \rangle > 0\}.$$

Note that in this way we are enlarging the positivity constraint on the variables (s, i, r) so the resulting equation is associated to a different problem with a possibly larger value function which we call V_1^1 . However, as explained, e.g., in the appendix of Boucekkine et al. (2019), this would allow to solve also the original one if the resulting optimal strategy satisfies the original constraints (i.e. the corresponding state trajectory \hat{h} stays in H_+).

- Second, the term $\tilde{A}(h, K)$ has the problem of requiring $h \in D(A)$, a property in general not satisfied by the mild solution \hat{h} of equation (17). Thus it is convenient to bring the operator \tilde{A} on the other side of the inner product. The drawback is that we need to require additional regularity for the solution v , that $D_h v$ belong to $D(A^*)$ (see Definition 4.1 below).

We then consider the unknown v defined on $H_+^1 \times \mathbb{R}_+$ and modify the Current Value Hamiltonian as follows (we let it keep the same name since we will be using only the following one from now on)

$$\begin{aligned} \mathbb{H}_{CV} &: (H_+^1 \times \mathbb{R}) \times ((D(A^*) \times \mathbb{R}) \times Z) \rightarrow \mathbb{R}, \\ \mathbb{H}_{CV}(h, K, p, Q; z) &= \langle (h, K), \tilde{A}^*(p, Q) \rangle_{H \times \mathbb{R}} + \langle \tilde{B}^z(h, K), (p, Q) \rangle_{H \times \mathbb{R}} + U_1(h; z) \\ &= \langle h_1, \frac{\partial p_1}{\partial a} + \mu_S p_1 \rangle_{L_{\pi_S}^2} + \langle h_2, \frac{\partial p_2}{\partial a} - \gamma p_2 + \frac{\gamma}{\pi_R^2} p_3 \rangle_{L^2} + \langle h_3, \frac{\partial p_3}{\partial a} + \mu_R \pi_3 \rangle_{L_{\pi_R}^2} \\ &\quad - \delta K Q - \langle \Lambda^{\theta, \eta}(h) h_1, p_1 \rangle_{L_{\pi_S}^2} + \langle \Lambda^{\theta, \eta}(h) h_1, p_2 \rangle_{L^2} - \langle \mu_I(\cdot, \Xi(h)) h_2, p_2 \rangle_{L^2} \\ &\quad + F(K, L^\theta(h)) Q - C^c(h) Q - D^\eta(h) Q + \langle (h_1 + h_2 + h_3)^\gamma(\cdot), u(z(\cdot)) \rangle_{L^2}. \end{aligned}$$

We denote by \mathbb{H}_{CV}^1 the part of the Hamiltonian that depends on the controls, i.e.

$$\begin{aligned} \mathbb{H}_{CV}^1(h, K, p, Q; z) &= -\langle \Lambda^{\theta, \eta}(h) h_1, p_1 \rangle_{L_{\pi_S}^2} + \langle \Lambda^{\theta, \eta}(h) h_1, p_2 \rangle_{L^2} + F(K, L^\theta(h)) Q \\ &\quad - C^c(h) Q - D^\eta(h) Q + \langle (h_1 + h_2 + h_3)^\gamma(\cdot), u(z(\cdot)) \rangle_{L^2}, \end{aligned}$$

and set

$$\begin{aligned} (36) \quad \mathbb{H}^0 &= \mathbb{H}_{CV} - \mathbb{H}_{CV}^1 \\ &= \langle h_1, \frac{\partial p_1}{\partial a} + \mu_S p_1 \rangle_{L_{\pi_S}^2} + \langle h_2, \frac{\partial p_2}{\partial a} - \gamma p_2 + \frac{\gamma}{\pi_R^2} p_3 \rangle_{L^2} + \langle h_3, \frac{\partial p_3}{\partial a} + \mu_R \pi_3 \rangle_{L_{\pi_R}^2} \\ &\quad - \delta K Q - \langle \mu_I(\cdot, \Xi(h)) h_2, p_2 \rangle_{L^2} \end{aligned}$$

so that

$$\sup_{z \in Z} \mathbb{H}_{CV}(h, K, p, Q; z) = \mathbb{H}^0(h, K, p, Q) + \sup_{z \in Z} \mathbb{H}_{CV}^1(h, K, p, Q; z).$$

Finally we set

$$\mathbb{H}^1(h, K, p, Q) := \sup_{z \in Z} \mathbb{H}_{CV}^1(h, K, p, Q; z)$$

so that the HJB equation (33) writes

$$(37) \quad \rho v(h, K) = \mathbb{H}^0(h, K, p, Q) + \mathbb{H}^1(h, K, p, Q).$$

Now we give the definition of classical solution of (37) in the interior of our enlarged state space $H_+^1 \times \mathbb{R}$. Here we abstract away from the boundary conditions as they will not be crucial for our purposes. Clearly they will become a key point when we want to prove results on existence/uniqueness/regularity of solutions of (37).

Definition 4.1 *We say that a function*

$$v: \text{Int } H_+^1 \times (0, +\infty) \longrightarrow \mathbb{R}$$

is a classical solution of the HJB equation (33) if

- (i) *v is continuously Fréchet differentiable in $\text{Int } H_+^1 \times (0, +\infty)$;*

- (ii) the derivative $D_h v(h, K)$ belongs to $D(A^*)$ for every $(h, K) \in \text{Int } H_+^1 \times (0, +\infty)$ and $A^* D_h v$ is continuous in $\text{Int } H_+^1 \times (0, +\infty)$;
- (iii) v satisfies equation (33) for every $(H, k) \in \text{Int } H_+^1 \times (0, +\infty)$.

We have the following result, which generalizes, e.g., Proposition 1.2, p. 225 of Li and Yong (1995).

Theorem 4.2 *Consider the problem of optimizing the target functional (21) over the set of control strategies*

(38)

$$\mathcal{Z}_{ad}^1(h_0, K_0) = \left\{ \hat{z}(\cdot) \in \mathcal{Z} : \left(\hat{h}^{\hat{z}; h_0}, K^{\hat{z}; K_0, h_0} \right) (t) \in H_+^1 \times \mathbb{R}_+ \text{ for a.e. } t \in [0, +\infty) \right\},$$

where H_+^1 is defined as in (35). Suppose that the value function V_1^1 of this “enlarged” problem is continuously Fréchet differentiable in $\text{Int } H_+^1 \times (0, +\infty)$ and that $D_h V(h, K) \in D(A^*)$ for every $(h, K) \in \text{Int } H_+^1 \times (0, +\infty)$. Then V is a classical solution of the Hamilton-Jacobi-Bellman equation (33) in $\text{Int } H_+^1 \times (0, +\infty)$.

Proof. For simplicity, in this proof, we will write V for V_1^1 and ∇V for the vector $(D_h V_1^1, D_K V_1^1)$. By Theorem 1.1, p. 224 of Li and Yong (1995) V satisfies the dynamic programming principle, that is, for every $(h_0, K_0) \in \text{Int } H_+^1 \times (0, +\infty)$ and every $t \geq 0$

$$(39) \quad V(h_0, K_0)$$

$$= \sup_{\hat{z} \in \mathcal{Z}_{ad}^1(h_0, K_0)} \left\{ \int_0^t e^{-\rho s} U_1 \left(\hat{h}^{\hat{z}; h_0}(s), \hat{z}(s) \right) ds + e^{-\rho t} V \left(\hat{h}^{\hat{z}; h_0}(t), K^{\hat{z}; h_0, K_0}(t) \right) \right\}$$

From now on for simplicity we will write $\underline{h}^{\hat{z}}(t)$ for $\left(\hat{h}^{\hat{z}; h_0}(t), K^{\hat{z}; h_0, K_0}(t) \right)$. Using the chain rule for mild solutions (see for example Proposition 5.5 Li and Yong, 1995) we have, for every $(h_0, K_0) \in \text{Int } H_+^1 \times (0, +\infty)$, every $\hat{z} = (\hat{c}, \hat{\theta}, \hat{\eta}) \in \mathcal{Z}_{ad}^1(h_0, K_0)$ and every $t \geq 0$,

$$\begin{aligned} V \left(\underline{h}^{\hat{z}}(t) \right) - V(h_0, K_0) &= \int_0^t \langle \underline{h}^{\hat{z}}(s), \tilde{A}^* \nabla V(\underline{h}^{\hat{z}}(s)) \rangle_{H \times \mathbb{R}} ds \\ &\quad + \int_0^t \langle \tilde{B}^{\hat{z}}(\underline{h}^{\hat{z}}(s)), \nabla V(\underline{h}^{\hat{z}}(s)) \rangle_{H \times \mathbb{R}} ds. \end{aligned}$$

Therefore, using also (39),

$$\begin{aligned}
0 &\geq \int_0^t e^{-\rho s} U_1(h^{\hat{z}; h_0}(s); \hat{c}(s), \hat{\theta}(s)) ds + e^{-\rho t} V(\underline{h}^{\hat{z}}(t)) - V(h_0, K_0) \\
&= \int_0^t e^{-\rho s} U_1(h^{\hat{z}; h_0}(s); \hat{c}(s), \hat{\theta}(s)) ds + e^{-\rho t} V(\underline{h}^{\hat{z}}(t)) - e^{-\rho t} V(h_0, K_0) \\
&\quad + (e^{-\rho t} - 1)V(h_0, K_0) \\
&= \int_0^t e^{-\rho s} U_1(h^{\hat{z}; h_0}(s); \hat{c}(s), \hat{\theta}(s)) ds + e^{-\rho t} \int_0^t \langle \underline{h}^{\hat{z}}(s), \tilde{A}^* \nabla V(\underline{h}^{\hat{z}}(s)) \rangle_{H \times \mathbb{R}} ds \\
&\quad + \int_0^t \langle \tilde{B}^{\hat{z}}(\underline{h}^{\hat{z}}(s)), \nabla V(\underline{h}^{\hat{z}}(s)) \rangle_{H \times \mathbb{R}} ds + (e^{-\rho t} - 1)V(h_0, K_0).
\end{aligned}$$

Now, since we are in an open set we know that the control strategies can be taken constant (so $\hat{z}(t) = \hat{z}(0)$ for $t \geq 0$) for a while. We divide both sides of the inequality by t and take the limit as $t \rightarrow 0$; finding

$$\begin{aligned}
(40) \quad 0 &\geq U(h_0; \hat{c}(0), \hat{\theta}(0)) + \langle (h_0, K_0), \tilde{A}^* \nabla V(h_0, K_0) \rangle_{H \times \mathbb{R}} \\
&\quad + \langle \tilde{B}^{\hat{z}}(h_0, K_0), \nabla V(h_0, K_0) \rangle_{H \times \mathbb{R}} - \rho V(h_0, K_0).
\end{aligned}$$

Therefore we obtain

$$0 \geq \sup_{z \in Z} \mathbb{H}_{CV}(h_0, K_0, D_h V(h_0, K_0), D_K V(h_0, K_0); z) - \rho V(h_0, K_0).$$

To prove the reverse inequality we fix again $(h_0, K_0) \in \text{Int } H_+^1 \times (0, +\infty)$; by definition of the value function, for any positive ϵ and any positive t we can find an admissible control $\hat{z}^\epsilon(\cdot) \in \mathcal{Z}_{ad}^1(h_0, K_0)$ such that

$$- \epsilon t \leq \int_0^t e^{-\rho s} U_1(\underline{h}^{\hat{z}^\epsilon}(s); \hat{c}^\epsilon(s), \hat{\theta}^\epsilon(s)) ds + e^{-\rho t} V(\underline{h}^{\hat{z}^\epsilon}(t)) - V(h_0, K_0) .$$

Using the equation satisfied by $\underline{h}^{\hat{z}^\epsilon}$ we get

$$\begin{aligned}
- \epsilon t &\leq e^{-\rho t} \left(V(\underline{h}^{\hat{z}^\epsilon}(t)) - V(h_0, K_0) \right) \\
&\quad + \int_0^t e^{-\rho s} U_1(h^{\hat{z}^\epsilon; h_0}(s); \hat{c}^\epsilon(s), \hat{\theta}^\epsilon(s)) ds + (e^{-\rho t} - 1)V(h_0, K_0) \\
&= e^{-\rho t} \langle \tilde{T}(t)(h_0, K_0) - (h_0, K_0), \nabla V(h_0, K_0) \rangle_{H \times \mathbb{R}} \\
&\quad + e^{-\rho t} \left\langle \int_0^t \tilde{T}(t-s) \tilde{B}^{\hat{z}^\epsilon}(\underline{h}^{\hat{z}^\epsilon}(s)), \nabla V(h_0, K_0) \right\rangle_{H \times \mathbb{R}} ds \\
&\quad + \int_0^t e^{-\rho s} U_1(h^{\hat{z}^\epsilon; h_0}(s); \hat{c}^\epsilon(s), \hat{\theta}^\epsilon(s)) ds + o(t) + (e^{-\rho t} - 1)V(h_0, K_0)
\end{aligned}$$

We can then find a continuous function $\sigma : [0, +\infty] \rightarrow [0, +\infty]$ such that $\sigma(0) = 0$ and

$$\begin{aligned} \sigma(\epsilon) &\leq \frac{1}{t} e^{-\rho t} \langle (\tilde{T}(t) - \text{Id})(h_0, K_0), \nabla V(h_0, K_0), \rangle_{H \times \mathbb{R}} \\ &\quad + \frac{1}{t} \int_0^t \langle \tilde{B}^{\pi^\epsilon}(h_0, K_0), \nabla V(h_0, K_0) \rangle_{H \times \mathbb{R}} ds \\ &\quad + \frac{1}{t} \int_0^t e^{-\rho a} U_1 \left(h_0; \hat{c}^\epsilon(s), \hat{\theta}^\epsilon(s) \right) ds + o(1) + \frac{e^{-\rho t} - 1}{t} V(h_0, K_0) \\ &\leq \sup_{z \in Z} \mathbb{H}_{CV}(h_0, K_0, D_h V(h_0, K_0), D_K V(h_0, K_0); z) + \frac{e^{-\rho t} - 1}{t} V(h_0, K_0) + o(1), \end{aligned}$$

which implies, taking the limit as $t \rightarrow 0$,

$$\sigma(\epsilon) \leq \sup_{z \in Z} \mathbb{H}_{CV}(h_0, K_0, D_h V(h_0, K_0), D_K V(h_0, K_0); z) - \rho V(h_0, K_0).$$

Letting now ϵ go to 0 we get the result. \square

Remark 4.3 The above Theorem (4.2) holds in a completely analogous way for the other problems where the target is changed. Of course, in case of finite horizon problems the HJB equation is different; for example, in the case of target (29), it is

$$-\frac{\partial v(t, h, K)}{\partial t} = \sup_{z \in Z} \bar{\mathbb{H}}_{CV}(t, h, K, D_h v(t, h, K), D_K v(t, h, K); z)$$

for $t \in [0, T]$, $(h, K) \in \text{Int } H_+^1 \times (0, +\infty)$ and with final condition $v(T, h, K) = K$. Note that here $\bar{\mathbb{H}}_{CV}$ is defined exactly as \mathbb{H}_{CV} in (34), substituting U_1 with U_3 as given in (32). Note finally that in all such cases we would take the enlarged constraint $h \in H_+^1$ instead of $h \in H_+$.

5. VERIFICATION THEOREMS

We first recall the definition of optimal strategy for our starting problem and for the ‘‘enlarged’’ one.

Definition 5.1 For $(h_0, K_0) \in H_+ \times \mathbb{R}_+$ (respectively $(h_0, K_0) \in \text{Int } H_+^1 \times (0, +\infty)$), an admissible control strategy $\hat{z}^* \in \mathcal{Z}_{ad}(h_0, K_0)$ (respectively $\hat{z}^* \in \mathcal{Z}_{ad}^1(h_0, K_0)$) is called optimal at (h_0, K_0) if

$$V_1(h_0, K_0) = J_1(h_0, K_0; \hat{z}^*(\cdot)) \quad (\text{respectively } V_1^1(h_0, K_0) = J_1(h_0, K_0; \hat{z}^*(\cdot))),$$

that is, if it is a maximizer for J . The corresponding solution $(h^{\pi^*; h_0}, K^{\pi^*; K_0})$ of (19) is called an optimal state trajectory.

The following result is the so-called Verification Theorem which provides sufficient optimality conditions.

Theorem 5.2 *Let v be a classical solution of the HJB equation (33), with the additional property that for every $\hat{z} \in \mathcal{Z}_{ad}^1(h_0, K_0)$*

$$(41) \quad \lim_{T \rightarrow +\infty} e^{-\rho T} v \left(\hat{h}^{\hat{z}; h_0}(T), K^{\hat{z}; h_0, K_0}(T) \right) = 0;$$

then $V_1^1(h_0, K_0) \leq v(h_0, K_0)$ for every $(h_0, K_0) \in \text{Int } H_+^1 \times (0, +\infty)$. Moreover, if an admissible control $\hat{z}^ \in \mathcal{Z}_{ad}^1(h_0, K_0)$ is such that*

$$(42) \quad \sup_{z \in Z} \mathbb{H}_{CV}(\underline{h}^{\hat{z}^*}(t), \nabla v(\underline{h}^{\hat{z}^*}(t)), z) = \mathbb{H}_{CV}(\underline{h}^{\hat{z}^*}(t), \nabla v(\underline{h}^{\hat{z}^*}(t)), \hat{z}^*(t))$$

then \hat{z}^ is optimal at (h_0, K_0) and $V_1^1(h_0, K_0) = v(h_0, K_0)$.*

Proof. We write ∇v for $(D_h v, D_K v)$. Moreover, as in the proof of Theorem 4.2 we write for simplicity $\underline{h}^{\hat{z}}(t)$ in place of $(h^{\hat{z}; h_0}(t), K^{\hat{z}; h_0, K_0}(t))$. We first prove that, for every $\hat{z} \in \mathcal{Z}_{ad}^1(h_0, K_0)$ we have the fundamental identity

$$(43) \quad v(h_0, K_0) = J_1(h_0, K_0; \hat{z}(\cdot)) + \int_0^\infty e^{-\rho t} \left[\sup_{z \in Z} \mathbb{H}_{CV}(\underline{h}^{\hat{z}}(t), \nabla v(\underline{h}^{\hat{z}}(t)), z) - \mathbb{H}_{CV}(\underline{h}^{\hat{z}}(t), \nabla v(\underline{h}^{\hat{z}}(t)), \hat{z}(t)) \right] dt.$$

Indeed, differentiating the function $t \mapsto e^{-\rho t} v(\underline{h}^{\hat{z}}(t))$ and integrating on $[0, T]$ we find

$$\begin{aligned} v(h_0, K_0) &= e^{-\rho T} v(\underline{h}^{\hat{z}}(T)) + \int_0^T e^{-\rho t} \rho v(\underline{h}^{\hat{z}}(t)) dt \\ &\quad - \int_0^T e^{-\rho t} \langle (\underline{h}^{\hat{z}}(t), \tilde{A}^* \nabla v(\underline{h}^{\hat{z}}(t))) \rangle_{H \times \mathbb{R}} dt \\ &\quad - \int_0^T e^{-\rho t} \langle \tilde{B}^{\hat{z}(t)}(\underline{h}^{\hat{z}}(t), \nabla v(\underline{h}^{\hat{z}}(t))) \rangle_{H \times \mathbb{R}} dt. \end{aligned}$$

We can then add and subtract the term $\int_0^T e^{-\rho t} U_2(h^{\hat{z}; h_0}(t), \hat{z}(t)) dt$ on the right hand side and use the fact that v solves the HJB equation (33) to obtain

$$\begin{aligned} v(h_0, K_0) &= e^{\rho T} v(\underline{h}^{\hat{z}}(T)) + \int_0^T U_2(h^{\hat{z}; h_0}(t), \hat{z}(t)) dt \\ &\quad + \int_0^T e^{-\rho t} \left[\sup_{z \in Z} \mathbb{H}_{CV}(\underline{h}^{\hat{z}}(t), \nabla v(\underline{h}^{\hat{z}}(t)); \hat{z}(t)) - \mathbb{H}_{CV}(\underline{h}^{\hat{z}}(t), \nabla v(\underline{h}^{\hat{z}}(t)); \hat{z}(t)) \right] dt. \end{aligned}$$

The fundamental identity then follows taking the limit as $T \rightarrow \infty$ and using (41). Since the last integral is always non-negative, and eventually taking the supremum

over all admissible controls on the right hand side, we get the first claim. The second one follows observing that for such \hat{z}^* we have, thanks to (43) and the first claim,

$$V_1^1(h_0, K_0) \leq v(h_0, K_0) = J_1(h_0, K_0; \hat{z}^*)$$

which implies that $V_1^1(h_0, K_0) = v(h_0, K_0) = J_1(h_0, K_0; \hat{z}^*)$, thus the claim. \square

Corollary 5.3 *Let v be a classical solution of the HJB equation (33) and assume that the set valued map*

$$(h, K) \rightarrow \arg \max_{z \in Z} \mathbb{H}_{CV}(h, K, \nabla v(h, K); z)$$

admits a measurable selection $G : \text{Int } H_+^1 \times (0, +\infty) \rightarrow Z$. Let $(h_0, K_0) \in \text{Int } H_+^1 \times (0, +\infty)$ and assume that the closed loop equation

$$(44) \quad \begin{cases} \frac{d}{dt}(\hat{h}, K)(t) &= \tilde{A}(\hat{h}(t), K(t)) + \tilde{B}^{G(\hat{h}(t), K(t))}(\hat{h}(t), K(t)), \quad t \geq 0 \\ (\hat{h}, K)(0) &= (h_0, K_0), \end{cases}$$

admits a solution $(\hat{h}^{G;h_0}, K^{G;h_0, K_0})$ such that the control strategy

$$\hat{z}^*(t) = G(\hat{h}^{G;h_0}(t), K^{G;h_0, K_0}(t))$$

belongs to $\mathcal{Z}_{ad}^1(h_0, K_0)$. Then \hat{z}^ is optimal.*

Proof. The result immediately follows from the previous Theorem 5.2 and from the fundamental identity (43). \square

Corollary 5.4 *Suppose that the value function V_1^1 is a classical solution of the Hamilton-Jacobi-Bellman equation (33), that $\hat{z}^* \in \mathcal{Z}_{ad}^1(h_0, K_0)$ is optimal at (h_0, K_0) and that*

$$\lim_{T \rightarrow +\infty} e^{-\rho T} V_1^1(h^{\pi^*;h_0}(t), K^{\pi^*;K_0}(t)) = 0.$$

Then \hat{z}^ satisfies (42).*

Proof. Proceeding as in the proof of Theorem 5.2 we can show that the value function V satisfies the fundamental identity (43). Since $\hat{z}^*(\cdot)$ is optimal the integral term on the right hand side of (43) must be 0, and the claim follows. \square

Remark 5.5 The above results allow, if we can find V_1^1 , at least numerically, to solve the problems with the enlarged constraints. To pass to our initial control problem we have to show that, for some $(h_0, K_0) \in H_+ \times \mathbb{R}_+$, the optimal control of the enlarged problem is admissible for the initial problem as well. This has been done e.g. in Boucekine et al (2019) and the same idea may work in some special cases of our set-up.

6. CONCLUSION

Given the strong differences in the effects of some epidemics (and particularly that of COVID-19) as individuals vary in age, it is important, in trying to understand the economic impact of the contagion and in evaluating the policies to combat it, to model it as precisely as possible.

In the previous contributions which integrate the epidemiological dynamics in macro-dynamic models, the stratification by age of population is often absent and, when introduced, it is modeled using a finite number of groups with no possibility to move from one group to another.

In this paper we propose a general fully age-structured time continuous set-up for the macro analysis of epidemics and economic dynamics.

After rewriting the problem using a suitable Hilbert space reformulation of the associated infinite dimensional optimal control problem, we develop the dynamic programming approach providing verification type results which, given our general infinite dimensional setting, cannot be derived from previous results in the literature.

REFERENCES

- Acemoglu, D., Chernozhukov, V., Werning, I. and Whinston, M.D., (2020). Optimal targeted lockdowns in a multi-group SIR model. NBER Working Paper w27102.
- Alvarez, F.E., Argente, D. and Lippi, F., (2020). A simple planning problem for covid-19 lockdown. NBER Working Paper w26981.
- Anderson, R. M. and May, R. M., (1985). Age-related changes in the rate of disease transmission: implications for the design of vaccination programmes. *Epidemiology and Infection*, 94(3):365-436.
- Ainseba, B., Anita, S., and Langlais, M. (2002). Optimal control for a nonlinear age-structured population dynamics model. *Electronic Journal of Differential Equations*, 28, 1-9.
- Anita, S. (2000). *Analysis and control of age-dependent population dynamics*. Springer Science and Business Media.
- Bensoussan, A., Da Prato, G. Delfour, C. and Mitter, K. M. (2007). *Representation and control of infinite dimensional systems*. Birkhauser, Berlin.
- Boucekkine R., Fabbri G., Federico S., Gozzi F. (2019). Growth and agglomeration in the heterogeneous space: a generalized AK approach. *Journal of Economic Geography*, 19 (6), 1287-1318.
- Boucekkine, R., Hritonenko, N., and Yatsenko, Y. (Eds.). (2013). *Optimal control of age-structured populations in economy, demography, and the environment*. Routledge.
- Clemens, V., Deschamps, P., Fegert, J.M., Anagnostopoulos, D., Bailey, S., Doyle, M., Eliez, S., Hansen, A.S., Hebebrand, J., Hillegers, M. and Jacobs, B., 2020. Potential effects of “social” distancing measures and school lockdown on child and adolescent mental health. *European Child and Adolescent Psychiatry*.
- Colombo, R. M., Garavello, M., Marcellini, F., and Rossi, E. (2020). An Age and Space Structured SIR Model Describing the Covid-19 Pandemic. medRxiv.
- Dietz, K. and Schenzle, D., (1985). Proportionate mixing models for age-dependent infection transmission. *Journal of Mathematical Biology*, 22(1):117-120.

- Eichenbaum, M.S., Rebelo, S. and Trabandt, M., (2020). The macroeconomics of epidemics. NBER Working Paper w26882.
- Fabbri G., Gozzi F. (2008). Solving optimal growth models with vintage capital: The dynamic programming approach. *Journal of Economic Theory*, 143(1), 331-373.
- Fabbri, G., Gozzi, F. and Swiech, A., (2010). Verification theorem and construction of ε -optimal controls for control of abstract evolution equations. *Journal of Convex Analysis*, 17(2):611-642.
- Faggian, S. and Gozzi, F. (2010). Optimal investment models with vintage capital: Dynamic programming approach. *Journal of Mathematical Economics*, 46(2010):416-437.
- Favero, C. A., Ichino, A. and Rustichini, A., (2020). Restarting the economy while saving lives under Covid-19. CEPR Discussion Papers 14664.
- Feichtinger, G., Hartl, R.F., Kort, P.M. and Veliov, V.M. (2006) Anticipation Effects of Technological Progress on Capital Accumulation: a Vintage Capital Approach, *J. Econom. Theory* 126(1), 143–164.
- Feichtinger, G., Prskawetz, A., Veliov, V. M. (2004). Age-structured optimal control in population economics. *Theoretical Population Biology*, 65(4), 373-387.
- Feichtinger, G., Tragler, G., and Veliov, V. M. (2003). Optimality conditions for age-structured control systems. *Journal of Mathematical Analysis and Applications*, 288(1), 47-68.
- Gollier, C., (2020). Cost-benefit analysis of age-specific deconfinement strategies. *Covid Economics*, 24:1-31.
- Hritonenko, N., Yatsenko, Y., (2005). Optimization of harvesting age in an integral age-dependent model of population dynamics. *Mathematical Biosciences*, 195(2), 154-167.
- Iannelli, M., (1995). *Mathematical theory of age-structured population dynamics*. Giardini editori e stampatori, Pisa.
- Iannelli, M., Marthceva M., (2003). *Homogeneous dynamical systems and the age-structured SIR model with proportionate mixing incidence*, in “Evolution Equations: Applications to Physics, Industry, Life Sciences and Economics”, Iannelli, M. and Lumer, G., eds.; pp. 227-251. Birkhäuser, Basel.
- Iannelli, M. and Milner, F., (2017). *The basic approach to age-structured population dynamics*. Models Methods and Numerics, 10, Springer, Berlin.
- Jones, C.J., Philippon, T. and Venkateswaran, V., (2020). Optimal mitigation policies in a pandemic: Social distancing and working from home. NBER Working Paper w26984.
- Krueger, D., Uhlig, H. and Xie, T., (2020). Macroeconomic dynamics and reallocation in an epidemic. NBER Working Paper w27047.
- Li, X., Yong, J., (1995). *Optimal Control Theory for Infinite Dimensional Systems*, Systems and Control: Foundations and Applications. Birkhäuser, Boston.
- Martcheva, M., (2015). *An introduction to mathematical epidemiology*, Vol. 61 of “Texts in Applied Mathematics”. New York: Springer.
- McKendrick, A., (1925). Applications of Mathematics to Medical Problems. *Proceedings of the Edinburgh Mathematical Society*, 44, 98-130.
- Moghadas, S. M., Shoukat, A., Fitzpatrick, M. C., Wells, C. R., Sah, P., Pandey, A., Sachs, J. D., Wang, Z., Meyers, L. A., Singer, B. H. and Galvani, A. P., (2020). Projecting hospital utilization during the COVID-19 outbreaks in the United States. *Proceedings of the National Academy of Sciences*, 117(16):9122-9126.
- OECD, (2020). Economic Outlook, June 2020.
- Palivos, T. and Yip, C.K., (1993). *Optimal population size and endogenous growth*. *Economics letters*, 41(1), pp.107-110.

- Piguillem, F. and Shi, L., (2020). The optimal COVID-19 quarantine and testing policies. EIEF Working Paper 20/04.
- Richard, Q., Alizon, S., Choisy, M., Sofonea, M. T., and Djidjou-Demasse, R. (2020). Age-structured non-pharmaceutical interventions for optimal control of COVID-19 epidemic. medRxiv.
- Salje, H., Kiem, C.T., Lefrancq, N., Courtejoie, N., Bosetti, P., Paireau, J., Andronico, A., Hozé, N., Richet, J., Dubost, C.L. and Le Strat, Y., (2020). Estimating the burden of SARS-CoV-2 in France. *Science*, 369(6500):208-211.
- Skritek, B., and Veliov, V. M. (2015). On the infinite-horizon optimal control of age-structured systems. *Journal of Optimization Theory and Applications*, 167(1), 243-271.
- Veliov, V. M. (2008). Optimal control of heterogeneous systems: basic theory. *Journal of Mathematical Analysis and Applications*, 346(1), 227-242.
- Yong, J. and Zhou, X.Y., (1999). *Stochastic controls: Hamiltonian systems and HJB equations*. Springer, Berlin.

ACKNOWLEDGEMENTS

The work of Giorgio Fabbri is supported by the French National Research Agency in the framework of the “Investissements d’avenir” program (ANR-15-IDEX-02) and in that of the center of excellence LABEX MME-DII (ANR-11-LABX-0023-01).

The work of Fausto Gozzi and Giovanni Zanco is supported by the Italian Ministry of University and Research (MIUR), in the framework of PRIN projects 2015233N54_006 (“Deterministic and stochastic evolution equations”) and 2017FKHBA8_001 (“The Time-Space Evolution of Economic Activities: Mathematical Models and Empirical Applications”)

UNIV. GRENOBLE ALPES, CNRS, INRA, GRENOBLE INP, GAEL, GRENOBLE, FRANCE.
Email address: `giorgio.fabbri@univ-grenoble-alpes.fr`

DIPARTMENT OF ECONOMICS AND FINANCE, LUISS UNIVERSITY, ROME, ITALY.
Email address: `fgozzi@luiss.it`

DIPARTMENT OF ECONOMICS AND FINANCE, LUISS UNIVERSITY, ROME, ITALY.
Email address: `gzanco@luiss.it`

To professors Boucekkine, Chakraborty and Goenka
editors of the special issue on the Economics of Epidemics
of the Journal of Mathematical Economics

November 12, 2020

Object: Revision of the paper JME-D-20-00380 : *Verification results for age-structured models of economic-epidemics dynamics* by Giorgio Fabbri, Fausto Gozzi and Giovanni Zanco

Dear professors,

Please find enclosed the second revision of our paper *Verification results for age-structured models of economic-epidemics dynamics* (G. Fabbri, F. Gozzi and G. Zanco), JME-D-20-00380.

First of all we would like to thank you for the fast and competent review process and for the opportunity of submitting a revision of the paper. We tried to answer the questions and to implement the suggestions (yours and the ones of the referees) at our best. We hope you and the referees will appreciate the result of this revision job.

In the following we briefly explain how we modified the paper in order to address the comments contained in your letter and in Reviewer 1's report. Further details are contained in the letter addressed to Reviewer 1. In *italics* we quote some parts of your letter.

- *We have received two referee reports on your paper. The two referees are top experts in the analysis of age structured populations. Reviewer 2, a pure mathematician, recommends accepting the current version. Reviewer 1, who is also an expert in mathematical biology and epidemiology, requires a major revision. Her/his detailed report outlines the heterogeneity of the contributions, the top quality of the methodological one but also a specification mistake. She/he ends up recommending a major revision principally on the basis that your verification theorem is a highly significant contribu-*

tion to the literature of age structured models. We concur with the views expressed by Reviewer 1. We therefore recommend to the Editor-in-chief of the JME to give you the opportunity to revise and resubmit the paper.

As outlined in the first referee report, the main contribution of your work is methodological. The paper has a very clear and innovative contribution with respect to all the related Covid-related papers (Acemoglu et al. or Gollier, for example, both cited) attempting to account for age structures. As correctly pointed out in your introduction, the treatment in Acemoglu et al. is partial. Gollier only runs scenarios (no optimization). It would be nice to be more accurate in your introduction to make even clearer your methodological contributions. The literature à la Kermac-McKendrick is not that old and disseminated in economic theory, it's important to show precisely what you add to the recent wave of related papers.

We agree with this point. The introduction has been focused on the methodological aspects of our contribution. We added some references on the literature à la Kermac-McKendrick and we pointed out in a precise way what is our addition to such literature: the development of a general setting for the Dynamic programming approach and the proof of the verification theorem in such general setting.

- *Section 3 is the essential part of the paper. It's also the most difficult part (formulation of the problem in an abstract infinite-dimensional setting). It's therefore of utmost importance that you make it clear enough and accessible to the casual reader of the JME (which is not granted for this current version). In particular, all the functional spaces should be carefully defined, even the most elementary (for example in separate footnotes when needed). Also please avoid digressions not supported by tangible arguments and/or references. For example, your footnote 7, page 11, is confusing. The target is to have a Section 3 rewritten in such way that the paper is self-contained and clear enough to the casual reader. The appendix can be used for further clarifications if needed.*

We have modified Section 3 to make it more clear to the casual reader. In particular we added definitions for all the functional spaces we use and we cut the footnote 7 at p.11 since it was confusing.

- *Last but not least, we urge you to address the remark of Reviewer 1 on your quadratic specification of the incidence rates. The related discussion will anyway strengthen the microfoundations of your model. Moving from quadratic to linear would not be a big deal for you, we do not foresee any crucial degeneracy problem.*

Following your indication we modified the formula for lambda (which is now linear) and corrected consistently the whole article.

To conclude we would like to thank again for inputs and suggestions and for the opportunity to submit a revised version of the paper. We hope we met all your requests.

Kind regards,

Giorgio Fabbri

Fausto Gozzi

Giovanni Zanco

To Reviewer 1
of the manuscript JME-D-20-00380

November 12, 2020

Object: Revision of the paper JME-D-20-00380 : *Verification results for age-structured models of economic-epidemics dynamics* by Giorgio Fabbri, Fausto Gozzi and Giovanni Zanco

Dear Reviewer,

Please find enclosed a revised version of our paper *Verification results for age-structured models of economic-epidemics dynamics* (G. Fabbri, F. Gozzi and G. Zanco).

Firstly, we would like to thank you for your work and for your comments. You actually found some true “bugs” in the first version of the paper, so your suggestions significantly helped us improving the quality of the manuscript. We implemented all of them in the attached revised version. We hope you will like the result of our work.

In this letter we describe how we revised the manuscript in order to respond to your comments and suggestions. In the following we present answers to specific points (in italics).

Comments that may be taken into account in a potential revision

- *Optimal control of age-structured (or age- and duration-structured) epidemiological models is an established area, although the authors ignore*

mentioning the existing literature.

Thank you for point out this lack. In the revised version we include some important references on control of age-structured (and in particular epidemiological) models. Most of the 10 new references included in the appendix are on this subject.

- *One can include social distancing and testing policies in various ways, out of which the authors have made their choice. The economic component of the model is standard. Also combining the two components of the model is straightforward, although various choices are possible. The authors say that as far as they know “this is the first paper ... on COVID-19... where capital accumulation is explicitly taken into account”. This might be true, but this reviewer is aware of submitted papers that propose age-structured models including macroeconomic dynamics of similar kinds (also in a more or less straightforward way). Therefore, I think the model is not a major contribution of this paper.*

Thank you for the indication. In the revised version we have changed the sentence. We have searched again with attention, we found the paper by Richard et al., (2020) that we added in the reference list and in the text (if you have in mind some more other reference we will gladly add it).

- ***Important:*** *The incidence rate $\lambda^{\theta,\eta}$ in (4) depends “quadratically” on θ . This does not seem to be true. If the authors look more carefully at the micro-foundation of (5), they will see that the dependence on θ should be linear. This mistake should be corrected with all the consequences all over the paper.*

Following your suggestion, we modified the expression for lambda and changed consistently the whole article.

Some additional remarks

- *The cost of testing is involved in the model, although this cost is less*

than 1% of the total economic losses due to COVID-19 in most countries. On the other hand, the cost of lost labor hours due the epidemic (even if no contact restrictions are undertaken) is not included (since $\phi(1) = 1$). The latter cost is larger than the cost of testing (at least \$ 130 billion in the USA due to influenza; for COVID-19 it will be higher).

It is true that $\varphi(1) = 1$ (so that when measures of type “lockdown” are not put into effect there is not a decrease of productivity of the active persons). It is also true however that the infected persons are not counted in the integral that defines the offer of job (in efficiency units) and therefore it seems to us that the cost of lost labor hours due the epidemic is included in the model.

Conversely it is true that the cost of testing people has not been so far extremely high but some arguments can mitigate this evidence. First, the cost it is not completely negligible. For instance in France, at the current level, the cost of testing is around 5 billions of euros¹ per year but the number of tests remains probably insufficient (knowing that the positivity rate is, this week, around 12%²) and this data do not take into account preventive quarantine social costs. Second, other, much more massive testing policies could be implemented, and choosing the best testing policy is among the questions that one can try to answer using models. For instance UK apparently studied a possible plan to massively increase the number of tests expanding testing to 10 million a day at a cost of 100bn of pounds³.

- More about the cost of testing: no testing means $\eta = 1$, full testing means $\eta = 0$. Maybe I misunderstand something, but if the integral in (3) represents cost, then $1 - \eta$ should appear there instead of η .

¹In France things are simpler than in US since all the tests are completely paid by the public wealth service. A larger test campaign only started some weeks ago, and the total projected cost for 2020 is around 2.2 billions of euros (<https://www.leparisien.fr/economie/depistage-du-covid-19-la-lourde-facture-des-tests-francais-16-09-2020-8386251.php>), meaning that at the current level of testing the yearly cost could be around 5 billions of euros.

²<https://www.santepubliquefrance.fr/maladies-et-traumatismes/maladies-et-infections-respiratoires/infection-a-coronavirus/documents/bulletin-national/covid-19-point-epidemiologique-du-15-octobre-2020>

³<https://www.bmj.com/content/370/bmj.m3520>.

We thank you very much for pointing out this mistake. We corrected it in the integral you mentioned and everywhere in the sequel.

- *One of the two objective functions considered in Section 4 is (14). If I do not overlook something, this objective function does not depend on the economic component at all. Moreover, the solution with respect to θ seems to be obvious: $\theta \equiv 0$.*

You are completely right. We corrected it.

- *The authors prove a verification result for the solution of the HJB equation, but do not give any consequences of this result that provide qualitative or quantitative information about the original problem. The paper seems to be fully theoretical.*

We agree with you. Indeed the purpose of the paper, as we tried to emphasize more in this revised version, is to provide a precise theoretical framework for applying dynamic programming to the family of infinite dimensional problems mentioned in Section 2. This is already a nontrivial task which, to our knowledge, has not yet been done in the literature.

Additional statement

- *I cannot judge how understandable the paper will be for the majority of the JME readers, but the dissemination of the used methodology should be encouraged.*

In the revised version we have taken into account your concerns (and those of the editors) about the readability of the paper for casual readers of JME.

Indeed:

- we modified the introduction making more clear the contribution of the paper and adding more references;
- we modified Section 3 to make it more readable and self-contained,

adding explanations for the abstract notations we use.

We hope that we have addressed all your requests in a satisfactory way, and that you appreciate the revised version of the manuscript.

Kind regards,

Giorgio Fabbri

Fausto Gozzi

Giovanni Zanco

Giorgio Fabbri
CNRS - Grenoble Applied Economics Lab
address: GAEL - CS 40700
38058 Grenoble CEDEX 9, France
email: giorgio.fabbri@univ-grenoble-alpes.fr
Tel.: +33 4 76 92 54 23
Web: sites.google.com/site/giorgiofabbri1979

Professor Carvajal
University of California Davis, US
FGV, Rio de Janeiro, Brazil
Editor In Chief of the *Journal of
Mathematical Economics*

Grenoble, August 17, 2020

Object: Declaration of interest for the article *Verification results for age-structured models of economic-epidemics dynamics*

Dear Professor Carvajal,

We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

We confirm that the manuscript has been read and approved by all named authors and that there are no other persons who satisfied the criteria for authorship but are not listed.

We further confirm that the order of authors listed in the manuscript has been approved by all of us.

We confirm that we have given due consideration to the protection of intellectual property associated with this work and that there are no impediments to publication, including the timing of publication, with respect to intellectual property. In so doing we confirm that we have followed the regulations of our institutions concerning intellectual property.

We understand that the Corresponding Author is the sole contact for the Editorial process (including Editorial Manager and direct communications with the office). He/she is responsible for communicating with the other authors about progress, submissions of revisions and final approval of proofs.

We confirm that we have provided a current, correct email address which is accessible by the Corresponding Author.

Sincerely,

Giorgio Fabbri
Fausto Gozzi
Giovanni Zanco