

Bruxism: an orthodontist's perspective

This is the peer reviewed version of the following article:

Original:

Colonna, A., Manfredini, D. (2024). Bruxism: an orthodontist's perspective. SEMINARS IN ORTHODONTICS, 30(3), 318-324 [10.1053/j.sodo.2023.12.010].

Availability:

This version is available <http://hdl.handle.net/11365/1253980> since 2024-01-16T17:49:38Z

Published:

DOI: <http://doi.org/10.1053/j.sodo.2023.12.010>

Terms of use:

Open Access

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.


(Article begins on next page)

Bruxism: an orthodontist's perspective

Anna Colonna , Daniele Manfredini

PII: S1073-8746(23)00125-1
DOI: <https://doi.org/10.1053/j.sodo.2023.12.010>
Reference: YSODO 770

To appear in: *Seminars in Orthodontics*



Please cite this article as: Anna Colonna , Daniele Manfredini , Bruxism: an orthodontist's perspective, *Seminars in Orthodontics* (2023), doi: <https://doi.org/10.1053/j.sodo.2023.12.010>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Bruxism: an orthodontist's perspective

Anna Colonna,¹ Daniele Manfredini¹

¹ School of Dentistry, Department of Biomedical Technologies, University of Siena, Viale Bracci c/o Policlinico "Le Scotte", 53100 Siena, Italy

Corresponding Author: Anna Colonna

School of Dentistry, Department of Biomedical Technologies, University of Siena, Viale Bracci c/o Policlinico "Le Scotte", 53100 Siena, Italy.

E-mail: annachiara.colonna@gmail.com

Journal Pre-proof

ABSTRACT

The aim of the present manuscript is to give a narrative overview for clinicians summarizing the knowledge concerning a phenomenon that they may frequently come across in both children and adults: sleep (SB) and awake (AB) bruxism. Indeed, prevalence rates in adults range from 8% to 16% for SB and 22% to 30% for AB, whilst in children they raise up to 40%.

The complex relationship between bruxism, the most common underlying primary and/or comorbid conditions, and the potential clinical consequences for both AB and SB in children and adults will be considered.

In addition, in view of the fact that over the last few decades the demand for orthodontic treatment at all ages has increased, the correlation between bruxism-related masseter muscle activity (MMA) and the use of orthodontic appliances (i.e., fixed appliances and clear aligners) is also discussed, along with the proposal of some clinical recommendations.

KEYWORDS: bruxism, orthodontics, aligners, awake bruxism, sleep bruxism, masseter muscle activity

BRUXSIM: THE STATE OF THE ART

Definition

Bruxism is a much-debated oral condition that interests several disciplines, such as dentistry, psychology, neurology, and sleep medicine, in both research and clinical settings. In this scenario, due to the constantly evolving knowledge and the different specialties involved in the study of bruxism, several definitions have been proposed over the past decades, until the point that the need to find a common language emerged.¹⁻⁵

Two expert consensus papers contributed substantially to define the construct of bruxism as an umbrella term to embrace a multifaceted spectrum of masticatory muscle activities (MMA), not limited to the act of grinding teeth while asleep.¹⁻²

The bruxism activities that were classically considered in the dental literature (viz., teeth grinding and clenching) were included in the consensus definitions together with an enlarged spectrum of analogue jaw muscle activities without tooth contact, such as mandible bracing (that is, clenching without teeth contact) or thrusting (that is, grinding without teeth contact). According to the proponents, this was a needed addition to achieve a better comprehension of the etiology and clinical consequences. In addition, bruxism activities were defined separately according to the circadian rhythm:

- *“Sleep bruxism is a masticatory muscle activity during sleep that is characterized as rhythmic (phasic) or non-rhythmic (tonic) and is not a movement disorder or a sleep disorder in otherwise healthy individuals.”*
- *“Awake bruxism is a masticatory muscle activity during wakefulness that is characterized by repetitive or sustained tooth contact and/or by bracing or thrusting of the mandible and is not a movement disorder in otherwise healthy individuals.”²*

Note that both definitions begin with ‘masticatory muscle activity’ (MMA), a wording intended to emphasise that focus is put on motor phenomena and the role of the masticatory muscles during sleep and wakefulness as the source of potential clinical consequences, independently on any specific neurological correlates. This suggests an ongoing paradigm shift with respect to the early concept of bruxism as a sleep-time motor activity featuring a specific pattern of masseter contraction (i.e., rhythmic masseter muscle activity [RMMA]) and being accompanied by teeth grinding sounds.⁵⁻⁷

This wording also intends to point out that bruxism should not be considered a disorder but rather a muscle behaviour that can have different aetiologies and that can be harmless,^{8,9} harmful (e.g., intrinsic mechanical tooth wear,⁸⁻¹² masticatory muscle and temporomandibular joint pain,^{13,14} temporomandibular disc displacements¹⁵ and fractures of teeth, restorations, and implants)^{12,16} or even potentially part of protective mechanisms with respect to several health outcomes (e.g., increasing upper airway patency which would aid in the prevention of airway collapse leading to obstructive sleep apnea¹⁷⁻¹⁹ or reducing the risk of detrimental chemical tooth wear by increasing salivation in case of gastro-oesophageal reflux²⁰).

Aetiology and pathophysiology

The concepts about the aetiology of bruxism mirrors the ongoing paradigm shift from peripheral to central regulation that is interesting several stomatognathic phenomena.

Increasing evidence suggests that a certain amount of bruxism activity can be considered physiological in all individuals, and that additive bruxism may be due to a combination of several psychosocial (e.g. stress sensitivity and anxiety)²¹⁻²⁵ biological, genetic, and exogenous factors (e.g., alcohol, smoking, caffeine, recreational substances and some drugs).²⁶⁻²⁸ On the contrary, morphological factors (e.g., features of the facial skeleton and dental occlusion) are no longer considered important.^{4,29}

In this view, whilst a multifactorial model is generically involved in the aetiology of bruxism, it must be underlined that each specific factor may have a different relationship with the different types of MMA.^{1-3,5}

It is also important to remark that AB is commonly characterised by teeth contacting habits or mandible bracing, whilst SB features a combination of all bruxism activities.^{3,5,28} Such different patterns of motor activity as a function of the circadian rhythm may also reflect different etiologies. On one hand, AB is mainly related to psychosocial factors, with a close association with certain personality traits,^{3,5,22,30,32} in other words, AB may be viewed as a stress-coping habit that is particularly frequent in patients with musculoskeletal symptoms.³³ On the other hand, SB is centrally mediated, with a complex interaction of all factors influencing autonomic system function during sleep.^{3,34-36}

Epidemiology

Data on the prevalence of bruxism are mainly related to SB, and are almost exclusively referred to questionnaire reports. Reported prevalence in adults range from 8% to 15% while the reported AB prevalence rates drawn from questionnaire studies range from 22% to 30%. No consistent data about time related fluctuations exists.^{3,37} In the recent years the development of Ecological Momentary Assessment (EMA) strategies made it possible to overcome some of the previous limitations concerning AB reports^{33,38-49} and allowed evaluating the frequency of reported AB behaviors compared to the self-reported activities identified in questionnaires. The results of the most recent smartphone-based EMA studies have suggested that the frequency of AB behaviors in young adults is within the 28.3–40% range of alerts received over one week of EMA monitoring.⁴⁹ In detail, teeth contact is the most common finding, whilst the report of teeth grinding is almost absent.⁴⁹ These preliminary findings could be seen as a reference point for future investigations on the epidemiological features of AB as well as for comparison with selected populations with comorbid conditions (e.g., sleep disorders, psychological and social features, muscle fatigue and muscle pain) in order to contribute a better understanding of this complex topic.^{30,49}

Assessment

The 2013 consensus paper¹ suggested a diagnostic grading system for the operationalisation of bruxism diagnosis (i.e., possible, probable, and definite sleep/awake bruxism) with the aim of defining the advantages and limitations of the available diagnostic approaches, but it is now clear that current knowledge and strategies are not sufficient to pursue such a hierarchical, stackable approach to bruxism evaluation.^{1-3,5,30,50}

In view of this, a Standardized Tool for the Assessment of Bruxism (STAB) has been recently proposed by an international expert panel. The main challenge was to answer the question “how to assess an individual’s bruxism in a reliable, valid, and relevant way?”. In addition to the presence or absence of MMA, clinicians should be able to screen for the potential risk factors, and comorbid conditions, as well as to determine the point at which bruxism becomes more likely associated with clinical consequences.^{51,52}

In general terms, approaches for assessing bruxism can be distinguished as non-instrumental or instrumental, and a combination of both approaches will likely emerge as the best available choice.^{1-3,51,52}

Non-instrumental approaches

Non-instrumental approaches for assessing bruxism include clinical examination and self-report (questionnaires, oral history) which can be used both for SB and AB.¹⁻³

As suggested in the STAB, the clinical examination should include an extraoral evaluation and an intraoral inspection (i.e., domains on joints and muscles, intra-and extra-oral tissues, and teeth and restorations) to identify the signs and symptoms that possibly related with bruxism.^{1-3,51,52}

Concerning the clinical evaluation of joints and muscles, the examiner may evaluate the masseter muscle for the presence of hypertrophy, assess the presence of one or multiple Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) diagnoses,⁵³ and evaluate functional symptoms (e.g., difficulty opening the mouth wide).

Regarding the Intra-oral tissues, the recommendation is to evaluate the presence of the following different signs: tongue scalloping, linea alba, lip impression, alveolar bone exostosis, tongue traumatic lesion. Concerning teeth and restorations wear and/or fracture, the evaluation is based on the Tooth Wear Evaluation System (TWES).¹⁰ It is also suggested to assess the presence of perforations and/or marks on an oral appliance/mouthguard and to perform a periodontal screening. In addition, clinical assessment may include optional items for selected research purposes, such as the skeletal class (e.g., Class I, II, III), facial pattern (e.g., hypo-, normo- and hyper-divergent) and tongue position based on the modified Friedman.^{51,52}

As a general note, it must be remembered that clinical assessment is actually deemed to evaluate the presence of the supposed clinical consequences of bruxism, rather than the actual bruxism activity (i.e., bruxism status) and that it is impossible to differentiate between consequences of SB and AB.³⁰

As for self-reported information, it can be obtained from questionnaires and history taking (i.e., oral interviews), allowing to gather information on perceived bruxism activities and the possible associated factors.^{54,55}

As pointed out in an early paper on the STAB project, there are currently no universally adopted questionnaires specifically designated for the assessment of AB.^{51,52} Examples of instruments that address AB as part of a more global report are the Oral Behaviors Checklist (OBC)⁵⁶ and the BRUX scale.⁵⁷ For this reason, timespan and frequency in which the report of AB is referred varies among the different questionnaires.

Considering SB, not only the patients themselves but also multiple informants can be interviewed (i.e., bed partner or, in the case of children, their parents). The patient and partner are asked to monitor behaviour concerning teeth grinding, teeth clenching and/or jaw bracing preferably using a diary. It is important to underline that, in the case of SB, the validity of findings is questionable due to the unconsciousness of patients during sleep.⁵⁸

Whilst the advantage of this approach is intuitively represented by the possibility to recruit large samples and to screen for the possible presence of bruxism at the individual level, the limitations are also well known.

First, it is important to underline that the intensity and duration of specific MMA cannot be exactly quantified via self-report. In addition, it cannot be excluded that the bruxism-psyche relationship could affect self-reporting, reflecting distress rather than masticatory muscle activity per se.^{59,60} Factors such as cognitive awareness, discriminatory ability and memory may also influence the report and, subsequently, the derived evaluation of bruxism status.

Therefore, the derived 'diagnosis' risks of having incomplete value because of its subjectivity, but it is nevertheless a basis for getting deeper into the assessment process and to gather data for epidemiological purposes and to screen for the possible presence of bruxism at an individual level.^{54,55,61}

Instrumental approaches

Instrumental approaches for assessment are currently available for both circadian forms of bruxism, although with differences related to the different characteristics presented by AB and SB.

Several devices have been available for years to record SB activities, where electromyography (EMG) and polysomnography (PSG) findings are fundamental to provide key evidence of motor activity and on the other hand to provide an evaluation of SB within the sleep architecture. For this reason, full PSG is recommended when the presence of other sleep disorders (e.g., apnoea) is suspected;^{2,3,6} on the contrary, its use cannot be justified for routine cases due to the costs and limited availability of the needed technical equipment. To overcome these limits, in recent years some EMG devices for in-home recordings have emerged as a valid option for an easier approach to assess the motor activity.⁶²⁻⁶⁷

EMG recordings during wakefulness may indeed provide measurements of AB but performing an hour-long EMG recording of jaw muscle activity during wakefulness is difficult for technical reasons and for the potentially poor patients' compliance; in addition, it's not easy to perform due to the paucity of dedicated devices on the market. Recent developments in this direction are promising, but on-field studies are yet to be performed on a large scale, also with the aim of refining the discrimination between AB and non-AB activities during wakefulness.³

In this scenario, it is important to underline that PSG/SB criteria that are still adopted in the field of dental sleep medicine are based just on the count of masseter EMG events associated with sleep arousals, thus providing only a partial picture of the complex range of jaw-muscle activities. Long-lasting, prolonged, low activity contractions, that are now incorporated within the bruxism definition, are excluded from that count, thus putting into serious doubt the internal validity of previous PSG/SB literature findings concerning the etiology and consequences of SB. In addition to that, definitive criteria have never been established for awake bruxism measurement. In view of this, the expert consensus panel pointed out the need to evaluate this phenomenon in its continuum.^{3,50, 66, 68}

As recently suggested by several studies^{33, 38-49} and by the expert consensus papers on bruxism definition,^{1,2} these limitations may be overcome by adopting EMA strategies, also called experience sampling methodology (ESM), to report AB behaviors, since it is a simple method to collect real-time data in the natural environment.

The EMA approach may be optimized with the use of smartphone apps, which are easy to use and intuitive;^{69,70} for this reason, this approach also belongs to the category of instrumental approaches.

This data-recording strategy can find applications for both research and clinical purposes: in the research setting it can be used to increase knowledge on several aspects of AB (e.g., natural course and fluctuations of signs, symptoms, and exposure to etiological factors); in the clinical setting, it helps patients to recognize their habits, monitor changes over time, and adopt corrective measures.

In general terms, as suggested by the international expert group, assessing bruxism with a combination of the different approaches may represent the best possible strategy to overcome the limitations of the different stand-alone methods.¹⁻³

Differential diagnosis

From a clinical point of view, it is interesting to underline that there are a series of conditions for which a differential diagnosis with bruxism may be necessary. Amongst these, noteworthy disorders are: Oromandibular dystonia, Huntington's disease, Tourette's syndrome, Hemifacial spasms, Parkinson's disease, Tardive dyskinesia and REM-behaviour disorder.

For more details the readers are referred to the original publications.^{3,71}

Management

In the light of the paradigm shift that emerged over the past quarter of a century (i.e., bruxism should not be considered a disorder but rather a muscle behaviour that can have different aetiologies and that can be harmless, harmful or even protective with respect to several health outcomes) it is strongly suggested to be prudent with the management of bruxism.^{8, 68, 71}

In this scenario, management should be comprehensively based on common sense conservative approaches, referring to the so-called 'Multiple-P' approach as the standard of reference:⁷²

- Pep talk (counselling)
- Psychology (cognitive-behavioural strategies)
- Physiotherapy (exercises of the jaw muscles)
- Plates (oral appliances)
- Pills (drugs)

The most important "P" is nonetheless the Patient, whose expectations and individual features should steer any clinical decisions. Importantly, it must be remarked that bruxism must always be seen as a condition that mirrors some underlying disorders. As such, the actual treatment, viz., the reduction of bruxism activity, must be discriminated from the management of clinical consequences. The former can be achieved just by addressing the potential cause or comorbid conditions, such as any psychological, neurological, or sleep disorder or any habit that is associated with an additive bruxism.

An important issue for clinicians is that performing irreversible occlusal changes with the aim to decrease pain symptoms in the jaw muscles and/or the TMJ or to reduce bruxism activities is not recommendable, since the association of bruxism with occlusal features is negligible, if at all present.⁷³⁻⁷⁵

BRUXISM IN PEDIATRIC AGE

Bruxism has always been considered an important phenomenon in children, especially due to the worries of parents who reports hearing teeth grindings sounds during their child's sleep. Actually, data about the clinical impact and management of such condition are scarce.

Concerning the epidemiology, as in the case of the adult population, the reported prevalence of SB in pediatric age varies among different studies and is based on parental report of teeth grinding. A recent systematic review suggested that the SB prevalence in children is between 3.5 and 40.6%, without gender differences and decreasing with age (e.g., 3.5-8.5% in children aged under 5 years, and less than 6% in children aged between 7 and 11 years).⁷⁶

Regarding the prevalence of bruxism in adolescents, in a sample of Israeli subjects, AB and SB were reported by 19.2% and 9.2% of participants, respectively, with no gender differences;⁷⁷ while a prevalence of 8.7% and 14.8% for AB and SB respectively, has been reported in a sample of Dutch adolescents.⁷⁸ In addition, according to other studies, higher prevalence rates were found in children and adolescents (3–49%) than in adults (1–15%).⁷³

Among the possible factors that are associated to SB in children, several studies assessed the relationship between parental-reported SB with secondhand smoke and concurrent sleep disturbances (e.g., nocturnal breathing symptoms, bedtime problems, night awakening, nocturnal and mooring symptoms).⁷⁹⁻⁸¹

Other studies found an association with behavioral problems, such as poor school performance, hyperactivity, and attention deficit. Also genetics, anxiety, psychological reactions and some personality features (e.g., high sense of responsibility), concurrent headaches, peer problems, emotional symptoms, mental health problems, aggressive behavior, ADHD and unsteady family environment seem to present a possible association with bruxism in pediatric age.⁸²⁻⁸⁵

Comparable findings have been found in teenagers (11 to 19 years): headache, sleep disturbances (snoring in particular), tooth wear, and jaw muscle fatigue, seem to be related to SB.⁸⁶ In addition, in young teenagers SB seem to be also associated with a history of episodes of verbal school bullying.⁸⁷

For assessment purposes, parental-reported tooth grinding remains the most common choice to perform investigations on a large scale, especially due to the difficulty of performing sleep studies with PSG in the pediatric age due to the important cost and the needed technical equipment. Consequently, the correlation of PSG findings with parental-reported tooth grinding is currently under investigation.⁸⁸

In children, efforts have always been mainly directed to the study of SB and its correlates, whilst information on AB frequency and its associated factors is missing; in view of this, as for adults, the use of EMA approaches is recommended as a strategy to get deeper into this issue at least in the adolescence age groups.

In view of the fact that bruxism in the pediatric age is a symptom of underlying health, lifestyle and biopsychosocial disorders, it is recommended that bruxism management is based on the identification of the underlying condition,

whenever possible.⁸⁹⁻⁹¹ In this scenario considering that the data on the natural course suggest that SB decreases progressively after the age of 9-10 years, relaxation techniques may be the best option for young children, whilst oral appliances are not suggested due to the ever-changing occlusal conditions.⁸⁹⁻⁹¹

Regarding pharmacological treatments, preliminary evidence indicates that hydroxyzine could be effective for parent-reported bruxism in children,⁹² but its routine use is not recommendable for risk-to-benefit ratio. Consequently, considering bruxism management in children and adolescences the construction of a multidisciplinary framework is recommended.⁹³

BRUXISM AND ORTHODONTICS

Over the last few decades, the demand for orthodontic treatment, both in children and adults, has increased; on the other hand, the effects of orthodontic appliances seem to have no clinically relevant effects at a population level as far as TMDs are concerned,⁹⁴ while possible changes in MMA have remained uncertain.

The neutrality of orthodontics on a large scale is due to the fact that, with the exception of some facial morphology features (e.g., skeletal Class II profiles hyperdivergent growth patterns and lower condylar volume) of which some occlusal traits may be proxy markers,^{95,96} the anatomy of dental occlusion contributes minimally to the multifactorial biopsychosocial model (i.e., biological, psychological, and social factors) of TMDs.⁹⁷⁻⁹⁸ Thus, it is understandable that the occlusal changes induced by orthodontics do not prevent or increase the risk of TMD development.^{94,99}

As far as bruxism is concerned, viz., the impact of orthodontics on MMA, the literature seems to report conflicting results.

In order to understand the possible different scenarios for bruxism-ortho relationships, it is important to underline that, as suggested by the consensus papers, for both circadian manifestations, there is a need to move on from the adoption of cut-off points to discriminate between bruxers and non-bruxers and embrace an evaluation based on the amount of activities that is related with the bruxism spectrum in their continuum, thereby not only focusing on the raw number of bruxism events.^{2,3,5,50-52} This is because the current line of thinking suggests that it is not the number of bruxism events per se to represents a risk factor for possible clinical consequences, but rather the general level of EMG activity, which was for instance found to be higher in TMD patients.^{3,68}

To this aim, the few devices available on the market for the assessment of AB should be enhanced to provide a better picture of the situation.^{66, 100}

In recent years, the demand for orthodontic treatments has grown, not only in children and adolescents, but also among adults. Another interesting point is the significant increase in cases treated with invisible aligners, instead of fixed traditional orthodontics, due to the possibility of achieving an aesthetic improvement with a reduced impact on social life, fewer emergencies, greater compliance, and better periodontal health compared to traditional orthodontics,¹⁰¹⁻¹⁰³ both in adults and adolescents.^{104,105}

Within the clinical community, however, there is concern about the potential effects of orthodontics, in particular in the case of aligners, on the homeostasis of the stomatognathic system, due to the potential effect on bruxism activity and the potential onset of muscle symptoms, which remain a debated issue both in clinical and research settings.¹⁰⁶⁻¹⁰⁹

In general, the literature on SB and AB seems to suggest a neutral effect of aligners, and more generally of orthodontics, in most cases.¹⁰⁶⁻¹⁰⁸

Considering AB assessed by EMG, Nota et al.,¹⁰⁶ and Lou et al.,¹¹⁰ reported respectively a decrease in masseter activity during the first month of treatment with aligners and a subsequent return to basal levels and, on the contrary, a transient increase in masticatory muscle activity during the first two weeks of aligner treatment. Pereira et al.¹⁰⁷ adopted a different evaluation strategy (i.e., ecological momentary assessment approach) and concluded that both in the case of fixed appliances or clear aligners the orthodontic treatment did not influence the frequency of AB report. In detail, while the fixed therapy resulted in a decrease at the start of treatment and a return to basal levels after 6 months, the clear aligners were associated with a constant frequency of AB report throughout the observation period.

With regard to SB, Manfredini et al.¹⁰⁸ investigated the effects of invisible orthodontic retainers with an EMG device and suggested an absence of relevant effects on sleep MMA in healthy individuals, in line with Castroflorio et al.¹¹¹ who analyzed the possible effects of clear aligners in six consecutive months in three different groups (i.e., clear aligners, occlusal splint and placebo splint).

An ongoing study evaluated the effects of aligners on sleep and awake MMA in healthy subjects over 24-hour timespans in the home environment with an EMG device during three different sessions: before starting the orthodontic treatment, with passive aligners and with active aligners. In this case, findings suggest that orthodontic treatment with aligners has an influence only in a minority of individuals.¹⁰⁹

In this context, dental professionals and orthodontists can find interesting support to the hypothesis that the use of orthodontic devices does not affect the sleep and awake bruxism activity, with minor exceptions related with inter-individual differences. For this reason, it is recommended to consider the patients' individual adaptability to prevent maladaptive behavior that is not necessarily related with the biomechanics of orthodontics.

In particular, the clinician must be able to evaluate the psychological and psychosocial aspects that play a decisive role in the etiology of most bruxism activities that, in turn, may be a crucial factor for the onset of clinical symptoms. Within this premise, patients' selection and the creation of a healthy doctor-to-patient relationship are actions that must not be underestimated for their value of good prognostic markers. When required, an orthodontist should not hesitate in looking for the involvement of other health and/or dental care professionals to provide multidisciplinary management of bruxism.

Therefore, clinicians in addition to perform a routine TMD-related examination before starting an orthodontic treatment, can use validated instruments (e.g., questionnaires) to collect information about psychosocial assessment of their patients. This would help a lot to identify those subjects who may experience an increase in bruxism-related behaviors.

CONCLUSIONS

- Knowledge on bruxism is rapidly evolving. Sleep and awake bruxism are considered masticatory muscle activities that, as an umbrella, group together several different jaw muscle activities that are not necessarily related to specific clinical consequences.
- Aetiological factors are mainly related to central rather than peripheral factors and may be different with respect to the circadian manifestations of bruxism. AB is mainly related with psychosocial factors, while SB has a more complex central mediation, with an interaction of all factors influencing autonomic system function during sleep.
- To assess bruxism, both non-instrumental and instrumental approaches can be recommended. A combination of both will likely emerge as the best available choice based on the concept of a multidimensional assessment of the bruxism status.
- Bruxism management is centered on the possible presence of clinical consequences and is based on “Multiple-P” approach (i.e., pep talk, psychology, physiotherapy, plates, pills) centered around the most important “P”: the patient. Therefore, performing irreversible occlusal changes (e.g., orthodontic treatment) is not recommendable.
- As for children and adolescents, in case of severe tooth wear and/or high frequency of parental-reported tooth grinding, special focus should be put on the search for comorbid conditions that may represent the actual health concern (e.g., respiratory disturbances, neurological disorders).
- In general, the literature suggests a neutral effect of orthodontic oral appliances (i.e., fixed appliance and clear aligners) on MMA both during wakefulness and sleep, with minor exceptions related with inter-individual differences.
- Before starting orthodontic treatment, clinicians are encouraged to evaluate the patients' individual psychological adaptability to avoid the occurrence of maladaptive behavior.

Acknowledgements

///

Funding

No founding or grant support.

Declaration of competing interest

The authors reported no competing financial interests or personal relationships that could appear to influence the work reported in this paper.

REFERENCES

1. Lobbezoo F, Ahlberg J, Glaros AG, et al. Bruxism defined and graded: An international consensus. *J Oral Rehabil.* 2013; 40:2-4.
2. Lobbezoo F, Ahlberg J, Raphael KG, Wetselaar P, Glaros AG, Kato T, Santiago V, Winocur E, De Laat A, De Leeuw R, Koyano K, Lavigne GJ, Svensson P, Manfredini D. International consensus on the assessment of bruxism: Report of a work in progress. *J Oral Rehabil.* 2018; 45:837–844.
3. Manfredini D, Colonna A, Bracci A, Lobbezoo F (2019) Bruxism: a summary of current knowledge on aetiology, assessment and management. *Oral surgery.* doi: 10.1111/ors.12454.
4. Lobbezoo F, Ahlberg J, Manfredini D, Winocur E. Are bruxism and the bite causally related? *J Oral Rehabil.* 2012;39:489-501.
5. Bracci A, Lobbezoo F, Colonna A, Bender S, Conti PCR, Emodi-Perlman A, Häggman-Henrikson B, Klasser GD, Michelotti A, Lavigne GJ, Svensson P, Ahlberg J, Manfredini D. Research routes on awake bruxism metrics: Implications of the updated bruxism definition and evaluation strategies. *J Oral Rehabil.* 2023;16. doi: 10.1111/joor.13514. Epub ahead of print. PMID: 37191494.
6. Lavigne GJ, Rompré PH, Montplaisir JY. Sleep bruxism: validity of clinical research diagnostic criteria in a controlled polysomnographic study. *J Dent Res.* 1996; 75:546-52.
7. Rompré PH, Daigle-Landry D, Guitard F, Montplaisir JY, Lavigne GJ. Identification of a sleep bruxism subgroup with a higher risk of pain. *J Dent Res.* 2007; 86:837-42.
8. Raphael KG, Santiago V, Lobbezoo F. Is bruxism a disorder or a behavior? Rethinking the international consensus on defining and grading of bruxism. *J Oral Rehabil.* 2016; 43:791-798.
9. Manfredini D, De Laat A, Winocur E, Ahlberg J. Why not stop looking at bruxism as a black/white condition? Aetiology could be unrelated to clinical consequences. *J Oral Rehabil.* 2016; 43:799-801.
10. Wetselaar P, Lobbezoo F. The tooth Wear evaluation system (TWES): a modular clinical guideline for the diagnosis and management planning of worn dentitions. *J Oral Rehabil.* 2016; 43:69-80.
11. Manfredini D, Lombardo L, Visentin A, Arreghini A, Siciliani G. Correlation between sleep-time masseter muscle activity and tooth wear: an electromyographic study. *J Oral Facial Pain Headache.* 2019; 33:199-204.
12. Abe S, Yamaguchi T, Rompré PH, De Grandmont P, Chen YJ, Lavigne GJ. Tooth wear in young subjects: a discriminator between sleep bruxers and controls? *Int J Prosthodont.* 2009; 22:342-50.

13. Manfredini D, Lobbezoo F. Sleep bruxism and temporomandibular disorders: a scoping review of the literature. *J Dent*. 2021; 111:103711.
14. Manfredini D, Lobbezoo F. Relationship between bruxism and temporomandibular disorders: a systematic review of literature from 1998 to 2008. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2010;109:e26-e50.
15. Kalaykova SI, Lobbezoo F, Naeije M. Effect of chewing upon disc reduction in the temporomandibular joint. *J Orofac Pain*. 2011; 25:49-55.
16. Manfredini D, Poggio CE, Lobbezoo F. Is bruxism a risk factor for dental implants? A systematic review of the literature. *Clin Implant Dent Relat Res*. 2014; 16:460-9.
17. Lobbezoo F, Ahlberg J, Manfredini D. The advancement of a discipline: The past, present and future of bruxism research. *J Oral Rehabil*. 2023 Jul 17. doi: 10.1111/joor.13562. Epub ahead of print. PMID: 37458517.
18. Manfredini D, Guarda-Nardini L, Marchese-Ragona R, Lobbezoo F. Theories on possible temporal relationships between sleep bruxism and obstructive sleep apnea events. An expert opinion. *Sleep Breath*. 2015; 19:1459-65.
19. Colonna A, Cerritelli L, Lombardo L, Vicini C, Marchese-Ragona R, Guarda-Nardini L, Manfredini D. Temporal relationship between sleep-time masseter muscle activity and apnea-hypopnea events: A pilot study. *J Oral Rehabil*. 2022; 49:47-53.
20. Ohmure H, Oikawa K, Kanematsu K, Saito Y, Yamamoto T, Nagahama H, Tsubouchi H, Miyawaki S. Influence of experimental esophageal acidification on sleep bruxism: a randomized trial. *J Dent Res*. 2011; 90:665-71.
21. Pierce CJ, Chrisman K, Bennett ME, Close JM. Stress, anticipatory stress, and psychologic measures related to sleep bruxism. *J Orofac Pain*. 1995; 9:51-6.
22. Manfredini D, Landi N, Fantoni F, Segù M, Bosco M. Anxiety symptoms in clinically diagnosed bruxers. *J Oral Rehabil*. 2005;32:584-8.
23. Manfredini D, Lobbezoo F. Role of psychosocial factors in the etiology of bruxism. *J Orofac Pain*. 2009; 23:153-66.
24. Manfredini D, Fabbri A, Peretta R, Guarda-Nardini L, Lobbezoo F. Influence of psychological symptoms on home-recorded sleep-time masticatory muscles activity in healthy subjects. *J Oral Rehabil*. 2011; 38:902-11.
25. Manfredini D, Arreghini A, Lombardo L, Visentin A, Cerea S, Castroflorio T et al. Assessment of coping and anxiety features in bruxers: a portable EMG/ ECG study. *J Oral Fac Pain Head*. 2016; 30:249-54.
26. Lobbezoo F, Soucy JP, Montplaisir JY, Lavigne GJ. Striatal D2 receptor binding in sleep bruxism: a controlled study with iodine-123-iodobenzamide and single photon emission computed tomography. *J Dent Res*. 1996; 75:1804-10.
27. Lavigne GJ, Lobbezoo F, Rompré PH, Nielsen TA, Montplaisir J. Cigarette smoking as a risk or exacerbating factor for restless legs syndrome and sleep bruxism. *Sleep*. 1997; 20:290-3.
28. Lobbezoo F, Aarab G, Van Der Zaag J. Definitions, epidemiology, and etiology of sleep bruxism. In: Lavigne GJ, Cistulli P, Smith M, editors: *Sleep Medicine for Dentists: A Practical Overview*. Chicago: Quintessence Publishing Co, Inc.; 2009:95-100.
29. Manfredini D, Visscher CM, Guarda-Nardini L, Lobbezoo F. Occlusal factors are not related to self-reported bruxism. *J Orofac Pain*. 2012; 26:163-7.

30. Bracci A, Lobbezoo F, Häggman-Henrikson B, Colonna A, Nykänen L, Pollis M, Ahlberg J, Manfredini D, International Network For Orofacial Pain And Related Disorders Methodology INFORM. Current Knowledge and Future Perspectives on Awake Bruxism Assessment: Expert Consensus Recommendations. *J Clin Med.* 2022; 11:5083.
31. Emodi-Perlman A, Manfredini D, Shalev T, Bracci A, Frideman-Rubin P, Eli I. Psychosocial and Behavioral Factors in Awake Bruxism-Self-Report versus Ecological Momentary Assessment. *J Clin Med.* 2021; 27;10:4447.
32. Colonna A, Guarda-Nardini L, Ferrari M, Manfredini D. COVID-19 pandemic and the psyche, bruxism, temporomandibular disorders triangle. *Cranio.* 2021; 15:1-6.
33. Câmara-Souza MB, Bracci A, Colonna A, Ferrari M, Rodrigues Garcia RCM, Manfredini D. Ecological Momentary Assessment of Awake Bruxism Frequency in Patients with Different Temporomandibular Disorders. *J Clin Med.* 2023; 7;12:501.
34. Lavigne GJ, Khoury S, Abe S, Yamaguchi T, Raphael K. Bruxism physiology and pathology: an overview for clinicians. *J Oral Rehabil.* 2008; 35:476–94.
35. Macaluso GM, Guerra P, Di Giovanni G, Boselli M, Parrino L, Terzano MG. Sleep bruxism is a disorder related to periodic arousals during sleep. *J Dent Res.* 1998; 77:565–73.
36. Huynh N, Kato T, Rompre PH, Okura K, Saber M, Lanfranchi PA et al. Sleep bruxism is associated To micro-arousals and an increase in cardiac sympathetic activity. *J Sleep Res.* 2006; 15:339–46.
37. Manfredini D, Winocur E, Guarda_Nardini L, Paesani D, Lobbezoo F. Epidemiology of bruxism in adults. A systematic review of literature. *J Orofac Pain.* 2013; 27:99–110.
38. Bracci A, Djukic G, Favero L, Salmaso L, Guarda-Nardini L, Manfredini D. Frequency of awake bruxism behaviors in the natural environment. A seven day, multiple-point observation of real time report in healthy young adults. *J. Oral Rehabil.* 2018; 45:423–429.
39. Colonna A, Lombardo L, Siciliani G, Bracci A, Guarda-Nardini L, Djukic G, Manfredini D. Smartphone-based application for EMA assessment of awake bruxism: compliance evaluation in a sample of healthy young adults. *Clin Oral Investig.* 2020; 24:1395-1400.
40. Nykänen L, Manfredini D, Lobbezoo F, Kämppi A, Colonna A, Zani A, Almeida AM, Emodi-Perlman A, Savolainen A, Bracci A, Ahlberg J. Ecological Momentary Assessment of Awake Bruxism with a Smartphone Application Requires Prior Patient Instruction for Enhanced Terminology Comprehension: A Multi-Center Study. *J Clin Med.* 2022; 15;11:3444.
41. Zani A, Lobbezoo F, Bracci A, Ahlberg J, Manfredini D. Ecological Momentary Assessment and Intervention Principles for the Study of Awake Bruxism Behaviors, Part 1: General Principles and Preliminary Data on Healthy Young Italian Adults. *Front Neurol.* 2019; 1;10:169.
42. Emodi-Perlman A, Manfredini D, Shalev T, Yevdayev I, Frideman-Rubin P, Bracci A, Arnias-Winocur O, Eli I. Awake Bruxism-Single-Point Self-Report versus Ecological Momentary Assessment. *J Clin Med.* 2021; 15;10:1699.
43. Dias R, Vaz R, Rodrigues MJ, Serra-Negra JM, Bracci A, Manfredini D. Utility of Smartphone-based real-time report (Ecological Momentary Assessment) in the assessment and monitoring of awake bruxism: A multiple-week interval study in a Portuguese population of university students. *J Oral Rehabil.* 2021; 48:1307-1313.

44. Câmara-Souza MB, Carvalho AG, Figueredo OMC, Bracci A, Manfredini D, Rodrigues Garcia RCM. Awake bruxism frequency and psychosocial factors in college preparatory students. *Cranio*. 2023; 41:178-184.
45. Zani A, Lobbezoo F, Bracci A, Djukic G, Guarda-Nardini L, Favero R, Ferrari M, Aarab G, Manfredini D. Smartphone-based evaluation of awake bruxism behaviours in a sample of healthy young adults: findings from two University centres. *J Oral Rehabil*. 2021; 48:989-995.
46. Emodi-Perlman A, Manfredini D, Shalev T, Bracci A, Frideman-Rubin P, Eli I. Psychosocial and Behavioral Factors in Awake Bruxism-Self-Report versus Ecological Momentary Assessment. *J Clin Med*. 2021; 27; 10:4447.
47. Bucci R, Manfredini D, Lenci F, Simeon V, Bracci A, Michelotti A. Comparison between Ecological Momentary Assessment and Questionnaire for Assessing the Frequency of Waking-Time Non-Functional Oral Behaviours. *J Clin Med*. 2022 ;5;11:5880.
48. Dias R, Lima R, Prado IM, Colonna A, Ferrari M, Serra-Negra JM, Manfredini D. Impact of Confinement by COVID-19 in Awake and Sleep Bruxism Reported by Portuguese Dental Students. *J Clin Med*. 2022; 18;11:6147.
49. Colonna A, Bracci A, Ahlberg J, Câmara-Souza MB, Bucci R, Conti PCR, Dias R, Emodi-Perlman A, Favero R, Häggmån-Henrikson B, Michelotti A, Nykänen L, Stanisić N, Winocur E, Lobbezoo F, Manfredini D. Ecological Momentary Assessment of Awake Bruxism Behaviors: A Scoping Review of Findings from Smartphone-Based Studies in Healthy Young Adults. *J Clin Med*. 2023; 28;12:1904.
50. Manfredini D, Ahlberg J, Wetselaar P, Svensson P, Lobbezoo F. The bruxism construct: From cut-off points to a continuum spectrum. *J. Oral Rehabil*. 2019; 46, 991–997.
51. Manfredini D, Ahlberg J, Aarab G, Bracci A, Durham J, Ettlin D, et al. Towards a standardised tool for the assessment of bruxism (STAB) - Overview and general remarks of a multidimensional bruxism evaluation system. *J Oral Rehabil*. 2020;47:549-56.
52. Manfredini D, Ahlberg J, Aarab G, Bracci A, Durham J, Emodi-Perlman A, Ettlin D, Gallo LM, Häggman-Henrikson B, Koutris M, Peroz I, Svensson P, Wetselaar P, Lobbezoo F. The development of the Standardised Tool for the Assessment of Bruxism (STAB): An international road map. *J Oral Rehabil*. 2022 Oct 19.
53. Schiffman E, Ohrbach R, Truelove E, et al. Diagnostic criteria for temporomandibular disorders (DC/TMD) for clinical and research applications: recommendations of the international RDC/TMD consortium network and orofacial pain special interest group. *J Oral Facial Pain Headache*. 2014; 28:6-27.
54. Van Selms M, Kroon J, Tuomilehto H, Peltomaa M, Savolainen A, Manfredini D, Lobbezoo F, Ahlberg J. Self-reported sleep bruxism among Finnish symphony orchestra musicians: Associations with perceived sleep-related problems and psychological stress. *Cranio*. 2023; 41:323-330.
55. Serra-Negra JM, Lobbezoo F, Correa-Faria P, Lombardo L, Siciliani G, Stellini E, Manfredini D. Relationship of self-reported sleep bruxism and awake bruxism with chronotype profiles in Italian dental students. *Cranio*. 2019; 37:147-52.
56. Markiewicz MR, Ohrbach R, McCall WD Jr. Oral behaviors checklist: reliability of performance in targeted waking-state behaviors. *J Orofac Pain*. 2006; 20:306-16.
57. Van der Meulen MJ, Lobbezoo F, Aartman IH, Naeije M. Self-reported oral parafunctions and pain intensity in temporomandibular disorder patients. *J Orofac Pain*. 2006; 20:31-5.

58. Raphael KG, Janal MN, Sirois DA, Dubrovsky B, Klausner JJ, Krieger AC, Lavigne GJ. Validity of self-reported sleep bruxism among myofascial temporomandibular disorder patients and controls. *J Oral Rehabil.* 2015; 42:751-8.
59. Van Selms MKA, Visscher CM, Knibbe W, Thymi M, Lobbezoo F. The association between self-reported awake oral behaviours and orofacial pain depends on the belief that these behaviours are harmful to the jaw *J Oral Facial Pain Headache.* 2020; 34:273-80.
60. Manfredini D, Ciapparelli A, Dell'Osso L, Bosco M. Mood disorders in subjects with bruxing behavior. *J Dent.* 2005; 33:485-90.
61. Manfredini D, Winocur E, Guarda-Nardini L, Lobbezoo F. Self-reported bruxism and temporomandibular disorders: findings from two specialised centres. *J Oral Rehabil.* 2012; 39:319-25.
62. Castroflorio T, Bargellini A, Rossini G, Cugliari G, Deregibus A, Manfredini D. Agreement between clinical and portable EMG/ECG diagnosis of sleep bruxism. *J Oral Rehabil.* 2015; 42:759–64.
63. Manfredini D, Ahlberg J, Castroflorio T, Poggio CE, Guarda-Nardini L, Lobbezoo F. Diagnostic accuracy of portable instrumental devices to measure sleep bruxism: a systematic literature review of polysomnographic studies. *J Oral Rehabil.* 2014; 41:836–42.
64. Casett E, Réus JC, Stuginski-Barbosa J, Porporatti AL, Carra MC, Peres MA, de Luca Canto G, Manfredini D. Validity of different tools to assess sleep bruxism: a meta-analysis. *J Oral Rehabil.* 2017; 44:722-34.
65. Castroflorio T, Deregibus A, Bargellini A, Debernardi C, Manfredini D. Detection of sleep bruxism: comparison between an electromyographic and electrocardiographic portable holter and polysomnography. *J Oral Rehabil.* 2014; 41:163–9.
66. Colonna A, Noveri L, Ferrari M, Bracci A, Manfredini D. Electromyographic Assessment of Masseter Muscle Activity: A Proposal for a 24 h Recording Device with Preliminary Data. *J Clin Med.* 2022; 12:247.
67. Colonna A, Segù M, Lombardo L, Manfredini D. Frequency of Sleep Bruxism Behaviors in Healthy Young Adults over a Four-Night Recording Span in the Home Environment. *Applied Sciences.* 2021; 11:195.
68. Manfredini D, De Laat A, Winocur E, Ahlberg J. Why not stop looking at bruxism as a black/white condition? Aetiology could be unrelated to clinical consequences. *J Oral Rehabil.* 2016; 43:799-801.
69. Yim SJ, Lui LMW, Lee Y, Rosenblat JD, Ragguett RM, Park C, Subramaniapillai M, Cao B, Zhou A, Rong C, Lin K, Ho RC, Coles AS, Majeed A, Wong ER, Phan L, Nasri F, McIntyre RS. The utility of smartphone-based, ecological momentary assessment for depressive symptoms. *J Affect Disord.* 2020; 1;274:602-09.
70. Raento M, Oulasvirta A, Eagle N. Smartphones: An emerging tool for social scientist. *Sociol. Methods Res.* 2009; 37, 426–454.
71. Manfredini D, Bracci A, Lobbezoo F. The right clinical information, right where it's needed. *BMJ Best Practice.* <https://newbp.bmj.com/topics/en-us/708>
72. Manfredini D, Ahlberg J, Winocur E, Lobbezoo F. Management of sleep bruxism in adults: a qualitative systematic literature review. *J Oral Rehabil.* 2015; 42(11):862–74.
73. Melo G, Duarte J, Pauletto P, Porporatti AL, Stuginski-Barbosa J, Winocur E et al. Bruxism: an umbrella review of systematic reviews. *J Oral Rehabil.* 2019; 46:666–90.
74. Manfredini D, Serra-Negra J, Carboncini F, Lobbezoo F. Current concepts of bruxism. *Int J Prosthodont.* 2017; 30:437–8.

75. Greene CS, Manfredini D. Treating Temporomandibular Disorders in the 21st Century: Can We Finally Eliminate the "Third Pathway"? *J Oral Facial Pain Headache*. 2020; 34:206-216.
76. Manfredini D, Restrepo C, Diaz-Serrano K, Winocur E, Lobbezoo F. Prevalence of sleep bruxism in children: a systematic review of the literature. *J Oral Rehabil*. 2013; 40:631-42.
77. Emodi Perlman A, Lobbezoo F, Zar A, Friedman Rubin P, van Selms MK, Winocur E. Self-Reported bruxism and associated factors in Israeli adolescents. *J Oral Rehabil*. 2016; 43:443-450.
78. Van Selms MK, Visscher CM, Naeije M, Lobbezoo F. Bruxism and associated factors among Dutch adolescents. *Community Dent Oral Epidemiol*. 2013; 41:353-63.
79. Castroflorio T, Bargellini A, Rossini G, Cugliari G, Rainoldi A, Deregibus A. Risk factors related to sleep bruxism in children: A systematic literature review. *Arch Oral Biol*. 2015; 60:1618-24.
80. Sousa HCS, Lima MDM, Dantas Neta NB, Tobias RQ, Moura MS, Moura LFAD. Prevalence and associated factors to sleep bruxism in adolescents from Teresina, Piauí. *Rev Bras Epidemiol*. 2018;21:e180002.
81. Segù M, Pollis M, Santagostini A, Meola F, Manfredini D. Correlation between Parental-Reported Tooth Grinding and Sleep Disorders: Investigation in a Cohort of 741 Consecutive Children. *Pain Res Manag*. 2020; 30;2020:3408928.
82. Guo H, Wang T, Niu X, Wang H, Yang W, Qiu J, Yang L. The risk factors related to bruxism in children: A systematic review and meta-analysis. *Arch Oral Biol*. 2018; 86:18-34.
83. Carra MC, Huynh N, Morton P, Rompré PH, Papadakis A, Remise C, Lavigne GJ. Prevalence and risk factors of sleep bruxism and wake-time tooth clenching in a 7- to 17-yr-old population. *Eur J Oral Sci*. 2011; 119:386-394.
84. Rossi D, Manfredini D. Family and school environmental predictors of sleep bruxism in children. *J Orofac Pain*. 2013; 27:135-141.
85. Restrepo C, Manfredini D, Lobbezoo F. Sleep behaviors in children with different frequencies of parental-reported sleep bruxism. *J Dent*. 2017; 66:83-90.
86. Castroflorio T, Bargellini A, Rossini G, Cugliari G, Deregibus A. Sleep bruxism in adolescents: a systematic literature review of related risk factors. *Eur J Orthod*. 2017; 39:61-68.
87. Serra-Negra JM, Pordeus IA, Corrêa-Faria P, Fulgêncio LB, Paiva SM, Manfredini D. Is there an association between verbal school bullying and possible sleep bruxism in adolescents? *J Oral Rehabil*. 2017; 44:347-353.
88. Restrepo C, Manfredini D, Castrillon E, Svensson P, Santamaria A, Alvarez C, Manrique R, Lobbezoo F. Diagnostic accuracy of the use of parental-reported sleep bruxism in a polysomnographic study in children. *Int J Paediatr Dent*. 2017; 27:318-325.
89. Barbosa Tde S, Miyakoda LS, Pocztaruk Rde L, Rocha CP, Gaviao MB. Temporomandibular disorders and bruxism in childhood and adolescence: review of the literature. *Int J Pediatr Otorhinolaryngol*. 2008; 72:299-314.
90. Restrepo C, Alvarez E, Jaramillo C, Vélez C, Valencia I. Effects of psychological techniques on bruxism in children with primary teeth. *J Oral Rehabil*. 2001; 28:354-360.
91. Restrepo C, Gómez S, Manrique R. Treatment of bruxism in children: a systematic review. *Quintessence Int*. 2009; 40:849-855.
92. Ghanizadeh A, Zare S. A preliminary randomised double-blind placebo-controlled clinical trial of hydroxyzine for treating sleep bruxism in children. *J Oral Rehabil*. 2013; 40:413-17.

93. Saulue P, Carra MC, Laluque JF, d'Incau E. Understanding bruxism in children and adolescents. *Int Orthod*. 2015; 13:489-506.
94. Manfredini D, Stellini E, Gracco A, Lombardo L, Nardini LG, Siciliani G. Orthodontics is temporomandibular disorder-neutral. *Angle Orthod*. 2016; 86:649-54.
95. Colonna A, Manfredini D, Lombardo L, Muscatello L, Marchese-Ragona R, Arveda N, Siciliani G. Comparative analysis of jaw morphology and temporomandibular disorders: A three-dimension imaging study. *Cranio*. 2020; 38:158-67.
96. Manfredini D, Segù M, Arveda N, Lombardo L, Siciliani G, Alessandro Rossi, Guarda-Nardini L. Temporomandibular Joint Disorders in Patients With Different Facial Morphology. A Systematic Review of the Literature. *J Oral Maxillofac Surg*. 2016; 74:29-46.
97. Suvinen TI, Reade PC, Kemppainen P, Könönen M, Dworkin SF. Review of aetiological concepts of temporomandibular pain disorders: towards a biopsychosocial model for integration of physical disorder factors with psychological and psychosocial illness impact factors. *Eur J Pain*. 2005; 9:613-33.
98. Manfredini D. The triangle bruxism, pain, and psychosocial factors. *Phd Thesis*. ACTA Amsterdam, 2011.
99. Michelotti A, Rongo R, D'Antò V, Bucci R. Occlusion, orthodontics, and temporomandibular disorders: Cutting edge of the current evidence. *J World Fed Orthod*. 2020; 9(3S):S15-S18.
100. Ymaguchi T, Mikami S, Okada K. Validity of a newly developed ultraminiature cordless EMG measurement system. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2007;104:e22-7.
101. Wang Q, Ma JB, Wang B, Zhang X, Yin YL, Bai H. Alterations of the oral microbiome in patients treated with the Invisalign system or with fixed appliances. *Am J Orthod Dentofacial Orthop*. 2019;156:633-40.
102. Feu D, Catharino F, Duplat CB, Capelli Junior J. Esthetic perception and economic value of orthodontic appliances by lay Brazilian adults. *Dental Press J Orthod*. 2012;17:102-14.
103. Colonna A, Siciliani G, Lombardo L. Orthodontic emergencies and perspectives during and after the COVID-19 pandemic: The Italian experience. *Pesquisa Brasileira em Odontopediatria e Clínica Integrada*, 2020; 21, pp. 1-8, e0181
104. Arveda N, Colonna A, Palone M, Lombardo L. Aligner hybrid orthodontic approach to treat severe transverse divergence in an adolescent girl: A case report. *Int Orthod*. 2022; 20:100686.
105. Arveda N, Colonna A, Siciliani G, Lombardo L. Class III correction using clear aligners, sectional wires, and miniscrew anchorage. *J Clin Orthod*. 2023; 57:54-62.
106. Nota A, Caruso S, Ehsani S, Ferrazzano GF, Gatto R, Tecco S. Short-Term Effect of Orthodontic Treatment with Clear Aligners on Pain and sEMG Activity of Masticatory Muscles. *Medicina*. 2021; 19:57:178.
107. Pereira NC, Oltramari PVP, Conti PCR, et al. Frequency of awake bruxism behaviour in orthodontic patients: Randomised clinical trial: Awake bruxism behaviour in orthodontic patients. *J Oral Rehabil*. 2021; 48:422-29.
108. Manfredini D, Lombardo L, Vigiani L, Arreghini A, Siciliani G. Effects of invisible orthodontic retainers on masticatory muscles activity during sleep: a controlled trial. *Prog Orthod*. 2018; 19:24.
109. Colonna A, Lobbezoo F, Gravili G, Lombardo L, Ahlberg J, Manfredini D. Effects of orthodontic aligners on 24-hour masseter muscle activity: a multiple-day electromyographic study. *In submission*
110. Lou T, Tran J, Castroflorio T, Tassi A, Cioffi I. Evaluation of masticatory muscle response to clear aligner therapy using ambulatory electromyographic recording. *Am J Orthod Dentofacial Orthop*. 2021; 159(1):e25-e33.

111. Castroflorio T, Bargellini A, Lucchese A, Manuelli M, Casasco F, Cugliari G, Cioffi I, Deregibus A. Effects of clear aligners on sleep bruxism: randomized controlled trial. *J Biol Regul Homeost Agents*. 2018;32 (2 Suppl. 2):21-29.

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: