



A Fungal Perspective on Conservation Biology

This is the peer reviewed version of the following article:

Original:

HEILMANN-CLAUSEN, J., Barron, E.S., Boddy, L., Dahlberg, A., Griffith, G.W., Nord´en, J., et al. (2015). A Fungal Perspective on Conservation Biology. CONSERVATION BIOLOGY, 29(1), 61-68 [10.1111/cobi.12388].

Availability:

This version is available <http://hdl.handle.net/11365/942642> since 2020-07-06T16:17:19Z

Published:

DOI:10.1111/cobi.12388

Terms of use:

Open Access

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.

(Article begins on next page)

1 Essay

2 **A fungal perspective on conservation biology**

3 Running head: A fungal perspective on conservation biology

4 Keywords: Decomposers, ecosystem services, forest ecology, indicator species, lichens, non-
5 timber forest products, pathogens, soil biology

6 Word count: 6050

7

8 Jacob Heilmann-Clausen¹, Elizabeth S. Barron², Lynne Boddy³, Anders Dahlberg⁴, Gareth
9 W. Griffith⁵, Jenni Nordén⁶, Otso Ovaskainen⁷, Claudia Perini⁸, Beatrice Senn-Irlet⁹, Panu
10 Halme^{1,10}

11 ¹Centre for Macroecology, Evolution and Climate, Natural History Museum of Denmark,
12 University of Copenhagen, Denmark (JHeilmann-Clausen@snm.ku.dk)

13 ²Department of Organismic and Evolutionary Biology, 16 Divinity Avenue, Harvard
14 University, Cambridge, MA 02138, USA (barrone@uwosh.edu)

15 ³School of Bioscience, Cardiff University, Museum Avenue, Cardiff CF10 3AX, Wales UK
16 (BoddyL@cardiff.ac.uk)

17 ⁴Swedish Species Information Centre, Swedish University of Agricultural Sciences, P.O. Box
18 7007, S-750 07 Uppsala, Sweden (Anders.Dahlberg@slu.se)

19 ⁵ Institute of Biological, Environmental and Rural Sciences, Cledwyn Building, Aberystwyth
20 University, Penglais, Aberystwyth, Ceredigion SY23 3DD, Wales UK (gwg@aber.ac.uk)

21 ⁶ Natural History Museum, University of Oslo, P.O. Box 1172 Blindern, NO-0318 Oslo,
22 Norway; Microbial Evolution Research Group, Department of Biosciences, University of
23 Oslo, P.O. Box 1066 Blindern, NO-0316 Oslo, Norway (jenni.norden@nhm.uio.no)

24 ⁷Department of Biosciences, P.O. Box 65, FI-00014 University of Helsinki, Finland
25 (otso.ovaskainen@helsinki.fi)

26 ⁸BIOCONNET- BIODiversity and CONservation NETwork, Department of Life Sciences,
27 University of Siena, Italy (claudia.perini@unisi.it)

28 ⁹ Swiss Federal Research Institute WSL, Zürcherstrasse 111, CH- 8903 Birmensdorf,
29 Switzerland (beatrice.senn@wsl.ch)

30 ¹⁰Department of Biological and Environmental Science, University of Jyväskylä, Finland
31 (panu.p.j.halme@jyu.fi)

32

33 Corresponding author: Jacob Heilmann-Clausen, Centre for Macroecology, Evolution and
34 Climate, Natural History Museum of Denmark, University of Copenhagen, Denmark
35 (JHeilmann-Clausen@snm.ku.dk), phone +45 23839997

36

37 **Abstract**

38 Hitherto fungi have rarely been considered in conservation biology, but this is changing as
39 the field moves from addressing single species issues to an integrative ecosystem-based
40 approach. The current emphasis on biodiversity as a provider of ecosystem services throws
41 the spotlight on the vast diversity of fungi, their crucial roles in terrestrial ecosystems and the
42 benefits of considering fungi in concert with animals and plants. But also for other reasons
43 fungal conservation science is growing as an independent field. In this paper we review the
44 role of fungi as actors in ecosystems, and provide an overview of the current state of fungal
45 conservation. On this basis we discuss five areas in which fungi can be readily integrated
46 into, and benefit conservation biology: 1) as providers of habitats and processes important for
47 other organisms, 2) as indicators on desired or undesired trends in ecosystem functioning, 3)
48 in identification of habitats of conservation value, 4) as providers of a powerful links between
49 human societies and the natural world as providers of food, medicine and biotechnological
50 tools, and 5) in the development of novel tools and approaches for conservation in
51 megadiverse organism groups. We hope that the conservation community will value these
52 potentials, and engage in mutualistic connections with mycologists, appreciating fungi as a
53 crucial part of nature

54

55 **Introduction**

56 Since the Rio Convention on Biological Diversity was signed in 1992, the conservation of
57 biological diversity has been an important topic in international politics, and the urgent need
58 for action was reignited at the tenth meeting of the Conference of the Parties to the
59 Convention on Biological Diversity in Nagoya (CBD 2010). Conservation initiatives have
60 evolved since the late 20th century from an initial focus on protection of pristine areas and
61 particular ('charismatic') species of animals and plants to a more holistic ecosystem-based
62 approach (e.g. Salafsky et al. 2002; Rands et al. 2010; Mace et al. 2012). So far fungi have
63 received limited emphasis in conservation biology (Vesterholt 2008; Minter 2010; Griffith
64 2012), except as potential threats to ecosystem health, individual species or species groups
65 (e.g. Fisher et al. 2012). Reasons for this neglect are complex but seem mainly to relate to a
66 general suspicious view on fungi in the Anglo-Saxon world, their hidden lifestyle and
67 challenging diversity, and a historical classification as an odd division of the *Plantae*.
68 (Minter 2010). We are certain that the situation is changing, both due to an ongoing
69 revolution in methods to obtain data on fungal species and communities (e.g. Peay et al.
70 2008; Halme et al. 2012), and because fungi are foundational to a wide variety of ecosystem
71 services.

72 In this essay we aim to indicate directions towards a full and balanced appreciation of fungi
73 in conservation biology. First, we review the critical roles fungi play in ecosystems. Then we
74 give a brief overview of the current state of fungal conservation. We show that fungal
75 conservation is important in its own right, and further stress how inclusion of the fungal
76 component of biodiversity can benefit conservation in general.

77

78 **Fungi as ecosystem actors**

79 Fungi constitute a megadiverse kingdom, with at least 1.5, but probably as many as 3-5
80 million species, of which only about 100,000 are formally described to date (Blackwell 2011;
81 Hawksworth 2012; Scheffers et al. 2012). Some are unicellular, but the majority form
82 mycelia, which range in size from colonies extending a few millimeters to some of the largest
83 organisms on the planet, e.g. honey fungi (*Armillaria* spp.) whose mycelia can occupy many
84 hectares of forest floor. The majority of fungi are hidden for most of their lives in the
85 substrates which they inhabit. Some form fruit bodies periodically or cause visible symptoms
86 in attacked host-plants, but only lichens are generally visible throughout most of their
87 lifecycle. Dispersal is usually passive, and maintained by microscopic, windborne spores, but
88 aquatic dispersal and animal vectors are important for many species. Profuse spore
89 production may easily lead to the view that fungi generally have much wider distribution
90 ranges and face less dispersal limitation than most other multicellular organisms. Evidence
91 for this idea is diminishing, as new research findings on spore dispersal (e.g. Norros et al.
92 2012) and fungal biogeography based on molecular markers (Taylor 2006; Salgado-Salazar et
93 al. 2013) show that fungi tend to be much less well dispersed and ubiquitous than believed in
94 the past.

95 Despite their hidden lifestyle, fungi maintain crucial processes in all terrestrial
96 ecosystems as decomposers of dead plant tissues and biotrophic partners of almost all
97 terrestrial multicellular organisms. As decomposers fungi are especially prominent in forests
98 and other ecosystems where grazing, fire or human harvesting are not dominant in carbon
99 cycling (Boddy et al. 2008). Plants produce between 5-33 t/ha of organic matter in forest
100 ecosystems every year, with an estimated global carbon pool of 73 petagrams in dead wood
101 (Pan et al. 2011). Most of this organic matter is lignocellulose, an intricate mixture of
102 recalcitrant biopolymers, with fungi being the only organisms possessing the requisite
103 enzymatic capability to mediate its efficient catabolism (Boddy et al. 2008). This process is

104 crucial for the release of nutrients and energy stored in plant litter, so fungi form the basis of
105 soil food chains and are grazed upon directly, or indirectly in plant litter, by a wide range of
106 invertebrate and vertebrate taxa (Stokland et al. 2012). In addition, networks of fungal hyphae
107 are stabilising soil particles into macroaggregates (Caesar-Tonthat 2002) and may thereby
108 protect soils against erosion (Tisdall et al. 2012).

109 Fungi are involved in diverse mutualistic associations. Lichenized fungi associated
110 with green algae or cyanobacteria, are highly stress-tolerant and mediate most primary
111 production and nitrogen fixation in desert and polar ecosystems, that covers 6 % of the
112 Worlds surface (e.g. Belnap 2002; Haas & Purvin 2006). They also dominate other
113 microhabitats in other climate zones such as tree trunks, rock surfaces and living leaves of
114 rainforest trees (Scheidegger & Werth 2009). Most plants (ca. 90% of species) are reliant on
115 mycelial networks intimately connected with their roots -mycorrhizas- for the uptake of
116 water, N, P and mineral nutrients from soil (Smith & Read 2008). In return for the water and
117 nutrients, mycorrhizal fungi receive substantial amounts of sugars from their plant partners,
118 typically 15 to 30 % of the net primary production (Chapin et al. 2011).

119 Mycorrhizal fungi are not only important for nutrient cycling, but also for mineral
120 weathering and carbon storage in forest ecosystems (Courty et al. 2010; Clemmensen et al.
121 2013). Further, they are tightly involved in plant competition, and because different groups of
122 fungi have very different enzymatic capacities, changes in plant composition mediated by
123 natural or anthropogenic processes might result in dramatic shifts in ecosystem processes
124 (Averill et al. 2014).

125 More cryptically, the internal tissues of all vascular plants host diverse communities
126 of asymptomatic fungal endophytes, of which some are mutualistic and prevent attacks from
127 pathogens and herbivores, while other are decomposers with a latent invasion strategy (e.g.
128 Rodriguez et al. 2009). Fungal endophytes represent a hyperdiverse group globally, both in

129 terms of unknown species and undiscovered bioactive compounds (Arnold & Lutzoni 2007;
130 Smith et al. 2008). As a functional group, fungal endophytes are not clearly delimited from
131 fungi classified as pathogens. In quite many cases beneficial effects to the host may shift to
132 pathogenic, due to environmental changes or imbalance in co-evolutionary processes. For
133 example, the recent outbreaks of ash-dieback in Europe are caused by the endophytic
134 *Hymenoscyphus pseudoalbidus*, which most likely originates in Eastern Asia where it lives in
135 non-pathogenic association with Manchurian Ash (*Fraxinus mandschurica*) (Zhao et al.
136 2012). In parts of Europe it has now replaced the native *Hymenoscyphus albidus*, that used to
137 be a harmless latent decomposer of dead leaves and petioles of the European Ash (*F.*
138 *excelsior*)(Pautasso et al. 2013). Other biotrophic fungi associate with animals, as mutualists,
139 e.g. in the rumen of herbivorous mammals or as a feeding source for insect larvae in wood, or
140 as parasites.

141 Sadly the public perception, and perhaps that of many conservation biologists, is that
142 fungi are extremely harmful because of the pathogenic ability of a few species (Fisher et al.
143 2012). Well known examples include the apparent extinction of several amphibian species
144 due to chytridiomycosis (Pounds et al. 2006) and the alteration of European and North-
145 American landscapes by chestnut blight, Dutch elm disease, and ash-dieback (Loo 2009;
146 Pautasso et al. 2013). However, natural disturbances are integral to the functioning and
147 continued evolution of ecosystems, and recent studies even suggest that pathogenic fungi are
148 drivers of biodiversity in tropical forest ecosystem, due to their density dependent attacks on
149 species that might otherwise become dominant by competitive exclusion (Bagchi et al. 2014).
150 Interestingly, many outbreaks of pathogenic fungi are caused or strongly reinforced by
151 human manipulations, not least the unintentional movement of fungal species around the
152 globe (e.g. Brasier 2008).

153

154 **Current state of fungal conservation**

155 The factors that threaten susceptible fungal populations are essentially the same as those
156 threatening animals and plants, including the degradation, loss and fragmentation of natural
157 and managed habitats, climate change, deposition of nitrogen and other pollutants (Sala et al.
158 2000; Dahlberg et al. 2010).

159 Fungal conservation is most highly developed in Fennoscandia (Dahlberg et al. 2010)
160 a region of relatively low overall biodiversity. We identify several reasons for this. First of
161 all, the boreal zone consists largely of coniferous forests, which provide a wealth of niches
162 for fungal species, but host relatively few vascular plants and larger animals. Secondly, and
163 perhaps linked to the scarcity of large charismatic animals, the tradition to focus more on
164 habitats than on specific species is deeply rooted in Fennoscandia (Raunio et al. 2008). In
165 practice, species from many species groups are used together to identify and prioritize
166 conservation measures. As discussed in the next section, cryptogams are well suited as
167 indicator species to identify sites, in particular forests, with specific conditions and histories.
168 Thirdly, Fennoscandia has a long tradition in fungal taxonomy and a good community of
169 amateur field biologists, which has resulted in a large and increasing knowledge on the
170 ecology and distribution of macrofungi that has formed the basis for the successful red-list
171 evaluation of more than 5000 species (Rassi et al. 2010).

172 Fungal red-listing is now widely used for management and conservation activities
173 across Europe; according to Dahlberg & Mueller (2011) only two of 35 national red lists for
174 fungi were produced in other parts of the world (New Zealand and Japan). A few countries
175 including Finland, Norway, Sweden and the UK have launched action plans to protect
176 specific fungal habitats and species, and in at least 12 European countries there are examples
177 of considering fungi in selection and prioritization of nature reserves (Senn-Irlet et al. 2007;
178 Dahlberg et al. 2010). Outside of Europe and the Pacific Northwest region of the USA

179 (Molina 2008) initiatives and strategies to conserve fungal biodiversity are more scattered
180 (but see Minter 2001; Buchanan & May 2003; Manoharachary et al. 2005; Abdel-Azeem
181 2010), and only three fungal species are currently globally red-listed. However, the situation
182 is changing, and the five fungal specialist groups of IUCN aim to have several hundred
183 fungal species globally red-listed in the near future (IUCN 2013). Organizations dedicated to
184 fungal conservation are also on the rise. The European Council for the Conservation of Fungi
185 (ECCF) was formed in 1985, and in 1991 a fungal specialist group was established within the
186 International Union for Conservation of Nature (IUCN). Since 2007, fungal conservation
187 committees or groups have also been established in Africa, South America and the US
188 (Barron 2011) and an International Society for Fungal Conservation (ISFC) was founded in
189 2011, suggesting a need for attention to fungal conservation at both the national and
190 international levels.

191

192 **What can fungi offer conservation biology?**

193 Current approaches to conservation acknowledge that human wellbeing and social resilience
194 depend on global biodiversity, a view that is formalized in the concept of ecosystem services.
195 The Millennium Ecosystem Assessment (World Resources Institute 2005) grouped
196 ecosystem services into four categories - regulating, supporting, provisioning and cultural
197 services. Like other multicellular organisms, fungi provide all of these (Pringle et al. 2011),
198 but the fundamental role fungi have as regulators of ecosystem processes in terrestrial
199 ecosystems places them centrally in the development of sustainable land use (Parker 2010;
200 Mace et al. 2012). However, it is just as evident that the majority of threatened fungi do not
201 contribute, and cannot even survive, in areas managed for timber and crop production. Hence
202 the arguments for their conservation should be based on arguments that are related to other
203 ecosystem services, some of which might be impossible to quantify in economic terms. We

204 believe that fungi deserve conservation in their own right, but below we will review how
205 conservation can benefit in general by the inclusion of fungi (Fig. 1).

206

207 Fungi as providers of services for other organisms

208 As described in the previous section, fungi are the drivers of several key processes in natural
209 ecosystems. Most of these are maintained by larger guilds of fungi, like the recycling of
210 nutrients from dead wood, or plant nutrition maintained by mycorrhizal fungi. Within guilds,
211 fungal communities are often very species rich, suggesting high levels of functional
212 redundancy. Both experimental (e.g. Strickland et al. 2009; Fukami et al. 2010) and
213 explorative studies (e.g. Taylor et al 2014) have reported high levels of niche differentiation
214 and less redundancy than expected in fungal communities, indicating that species identities
215 matter in major ecosystem processes where fungi contribute.

216 In other cases specific or smaller set of fungal species play key roles for other biota.
217 Fungi provide a principal food resource for many organisms, including mammals, orchids
218 and insects. In many cases associations are species specific or strongly selective, implying
219 that understanding of the fungal part of the association is crucial for the conservation of the
220 dependent feeders (e.g. Claridge & May 1994; Pyare & Longland 2002; Komonen 2003;
221 Bailarote et al. 2012). Polypores and other long-lived fleshy fruitbodies are particular rich
222 habitats for dependent insects, especially beetles and diptera. For example, the Dryad's
223 Saddle (*Polyporus squamosus* (Huds.) Fr.), hosts over 246 beetle species in Europe (Benich
224 1952). Other fungi are involved in the formation of microhabitats, such as cavities in trees
225 that are critical for hollow breeding birds, mammals, arthropods and epiphytes (e.g. Parsons
226 et al. 2003; Fritz & Heilmann-Clausen 2010; Remm & Löhmus 2011; Cockle et al. 2012). In
227 some cases these associations may be species specific (e.g. Jackson & Jackson 2004).

228

229 Fungi as indicators of ecosystem processes

230 With their narrow and thin-walled hyphae fungi are exposed to chemicals in the environment
231 and highly sensitive to microclimatic gradients, a fact that has been utilized in developing
232 indicator schemes based on fungi. Lichens are among the most sensitive organisms regarding
233 changes in air quality. In fact, the earliest record of biodiversity loss resulting from human
234 industrial activity was made by Thomas Pennant in 1773 who observed the decline of lichens
235 as a result of copper smelting at Parys Mountain, Wales (Pennant 1781). The differential
236 sensitivities of lichens to SO₂ and other airborne pollutants have since been widely used as a
237 proxy measure of air quality in both urban and natural habitats (Conti & Cecchetti 2001;
238 Nimis et al. 2002).

239 Non-lichenized fungi are also affected by SO₂ pollution, but anthropogenic nitrogen
240 pollution is now the most pervasive threat, with the decline of some ectomycorrhizal species,
241 e.g. stipitate hydroids and also *Cortinarius* spp. being particularly dramatic, though more
242 widespread changes in species composition in polluted areas are of equal concern (Arnolds
243 2001; Lilleskov et al. 2011).

244 The effects of global climate change on fungi are difficult to quantify, but it is
245 apparent that the warming climate over recent decades has altered the phenology of fungal
246 fruiting (Kausserud et al. 2012). For example, many fungi previously known to fruit only in
247 the fall now also fruit in spring, and mycorrhizal fungi associated with deciduous trees now
248 fruit later in the year. Changes in fungal community structure provide an early warning of
249 changing ecosystem processes, but so far there have been few efforts to implement this in
250 standardized monitoring schemes. Broadly, fungi constitute the most visible link to the vast
251 biodiversity underground, and are basal to the highly diverse decomposer food chains.

252 Incorporating fungi into ecosystem level indices such as the biodiversity intactness index
253 (Scholes & Biggs 2005) and the living planet index (Loh et al. 2005), which so far neglected
254 decomposers in general, would greatly enhance the value of these indices. Rapid advances in
255 the use of DNA-based methods for monitoring fungal communities (Schoch et al. 2012;
256 Lindahl et al. 2013) and increasing understanding of their functions, will likely facilitate the
257 use of fungi as bio-indicators of soil status and processes.

258

259 Fungi as indicators in conservation planning

260 The very specific habitat requirements of fungi make them well-suited as indicators for
261 selecting conservation areas and monitoring their status. A fungal angle on habitats simply
262 expands our understanding of the biotic space, and puts emphasis on microhabitats and
263 processes that are pivotal for biodiversity, but easily overlooked if fungi are not addressed.
264 For instance, specialized wood-inhabiting fungi may be absent from otherwise valuable
265 woodland habitats due to the lack of veteran trees and dead wood, and may become extinct at
266 the landscape scale if remaining old growth habitats are fragmented (Nordén et al. 2013).
267 Similarly, some ectomycorrhizal and lichenized fungi are highly sensitive to breaks in forest
268 continuity, and may be lost from forest ecosystems if mature trees are not retained through
269 rotations (Coppins & Coppins 2002; Rosenvald & Lõhmus 2008). These processes are also
270 important for many other organisms, including arthropods, molluscs and microfauna, but in
271 practice fungi will often be the easiest group to monitor.

272 Especially in Europe, several indicator schemes based on fungi have been suggested
273 to assess the conservation value of forests and grasslands (e.g. Coppins & Coppins 2002;
274 Heilmann-Clausen & Vesterholt 2008); and in Sweden and the Baltic countries fungi have
275 played a central role in the identification of key forest habitats – smaller areas selected to

276 lifeboat biodiversity in the managed forest landscape (Timonen et al. 2011). While fungal
277 indicator schemes are generally proposed based on field experience rather than hard
278 evidence, several studies have posthoc confirmed the validity of several indicator species
279 (e.g. Penttilä et al 2006; Müller et al. 2007).

280

281 Connections between fungi and humanity

282 The cultural value and public appreciation of fungi varies in different parts of the world, but
283 in the English-speaking world they have traditionally been viewed with great suspicion.
284 While this might be one reason that fungi have been somewhat overlooked in conservation
285 biology, the situation is clearly changing as people become more aware of the wide variety of
286 uses of fungi. In reality links between fungi and people are ancient. Fungi have been used as
287 food-sources, medicine, crafts, arts and tinder for thousands of years. They also feature in
288 religious ceremonies, where fungal statues and images are evident in relicts of ancient
289 civilizations and Stone Age art (Rutter 2010).

290 Wild fungi are a sustainable and renewable resource, which may help to turn public
291 opinion in favor of habitat conservation. Today, more than 1100 wild fungi are collected for
292 food or traditional medicine in over 80 countries worldwide (Boa 2004). Increasing global
293 markets for edible and medicinal mushrooms since the 1980s has led to increased harvesting
294 of many species both for subsistence use and for commercial sale. Over-exploitation by
295 harvesters (Minter 2010), or negative effects of harvesting on habitats (Egli et al. 2006) are
296 rare, and positive effects of increased use, such as increased awareness of fungi and their
297 habitats, yield many benefits for conservation. Their utility provides incentives for
298 conservation, as many prized wild fungi are restricted to relatively undisturbed natural
299 habitats. Indeed, edible wild fungi are increasingly seen as an economic alternative or

300 supplement to timber production in Europe and the United States (e.g. Aldea et al. 2012).
301 Even larger economic interests are associated with fungi as principal sources of enzymes,
302 antibiotics and other chemicals in the biotechnology sector. These interests are expected to
303 increase considerably in the coming century as novel products are discovered from fungi
304 (Erjavec et al. 2012; Rambold et al. 2013). This might help restore links between humanity
305 and nature at a discursive level, even though bioprospecting in general may be overrated as a
306 potential incentive for conservation in practice (Costello & Ward 2006).

307 In times of increasing concern for disconnectedness between growing urban
308 populations and the outdoors, the simple joy of collecting wild edible fungi with minimal or
309 no negative environmental impacts may be exactly the kind of activities that the conservation
310 movement should be encouraging through education and a focus on sustainability. The
311 tradition of public involvement in the scientific discipline of mycology is long. Even today
312 many fungal taxonomists collaborate with amateurs to obtain interesting specimens, and more
313 recently long time-series data from fungal forays have been used in high profile scientific
314 papers of conservation relevance (Gange et al. 2007; Kauserud et al. 2012). The amount and
315 quality of fungal data collected is increasing immensely through the development of internet
316 based platforms for species recording allowing easy storage of metadata, including
317 documentation photos, and facilitating communication between amateurs and professionals
318 (Halme et al. 2012).

319 While this development is very similar to what is happening in citizen science based
320 projects on birds, plants and butterflies, high fungal species richness and relatively poorly
321 resolved taxonomy impose new challenges and innovative solutions (Molina et al. 2011). For
322 instance, Emery and Barron (2010) involved local non-professional experts to investigate the
323 taxonomy and possible reasons for decline of edible morels in the US Mid-Atlantic Region,

324 hence shortcutting the link between amateur field knowledge and taxonomic expertise. Some
325 professional mycologists may see the growth of fungal amateur activity as a threat in a time
326 where funding to do basic taxonomic work is shrinking. However, successful citizen science
327 is only possible if backed by skilled professionals that can support and train the interested
328 amateurs. We fully agree with Korf (2005) and Barron (2011) that the limited environment of
329 professional mycologists could benefit by increasing involvement with the public, even
330 though this might imply a reconsideration of research questions and approaches.

331

332 Development of new tools for biodiversity monitoring

333 Finally, we believe that the current knowledge gap in fungal biodiversity may prove to be an
334 important driver in the development of novel tools with a broad relevance in conservation
335 biology, especially molecular analyses making use of DNA barcodes for species
336 identification. In part due to the rapid developments of high throughput ‘NextGen’ DNA
337 sequencing, remarkable new insights into fungal biodiversity have already emerged which in
338 some cases have direct conservation relevance (e.g. Kubartová et al. 2012; van der Linde et
339 al. 2012; Ovaskainen et al. 2013). A larger challenge is to put such information into an
340 appropriate conservation context and to combine it with other types of ecological knowledge.
341 Designing relevant sampling protocols for fungi, processing massive bioinformatic data sets
342 that include many unknown organisms (Hibbett et al. 2011), and considering relevance for
343 other organismic groups are all aspects of this emerging suite of methods that require
344 significant consideration moving forward.. Hence fungal conservation research strengthened
345 by metagenomics is not happening in isolation, and methodological improvements and
346 subsequent understanding of species distributions, dynamics and contributions to processes
347 are likely to have considerable impact in other fields of conservation biology.

348

349 **Conclusions**

350 Fungal conservation science is maturing as its own field, and has much to offer as
351 conservation biology moves from addressing single species to an integrative ecosystem
352 based approach. Fungi provide the most visible link to the vast biodiversity underground, and
353 are basal to the highly diverse decomposer food chains. In addition they are key mutualist
354 partners of plants and animals, playing fundamental regulating roles in all terrestrial
355 ecosystems. Incorporating mycological knowledge is crucial in the development of
356 sustainable practices in agriculture and forestry, in assessments of the state of natural
357 ecosystems, and in conservation planning that intends to cover all major aspects of
358 biodiversity.

359 Socially, due to their attractive fruit bodies, fungi represent a rich source of
360 wonderment, and are additionally valuable as food, in traditional medicine and as a source of
361 bioactive compounds. In most cases, modest collecting of wild fungi is non-detrimental to
362 ecosystems, and an increasing understanding of fungi may indeed help conservation to gain
363 broader understanding in rural as well as urban settings.

364 With an estimated 1.5 million species worldwide but only 100.000 species named so
365 far, many conservationists might suggest that seriously consideration fungi in conservation is
366 premature. While we agree that the big unknowns in fungal biology are challenging, we also
367 see obvious solutions. Given the magnitude of fungal diversity, the immense variation in life-
368 histories and ecological strategies, and the variety of links between fungi and people, a single
369 approach to fungal conservation is untenable and undesirable. Rather, a variety of case
370 specific strategies should be considered. For example, in the selection of forest patches for a
371 reserve network, polypores might be the most appropriate fungal tool. When considering

372 education and outreach campaigns, a focus on wild edibles and visually striking fungi makes
373 sense. When assessing effects of air pollution in urban setting, epiphytic lichens are the
374 obvious choice. This mirrors the situation in animal conservation, where various taxonomic
375 and functional groups are typically addressed separately, unless interactions or obvious
376 requirements for complementarity call for a complex approach.

377 Fungal conservation initiatives are currently under development within the
378 mycological community, and in different national and international organizations and
379 institutions where mycologists participate. We hope that the conservation community will
380 welcome these initiatives, and engage in mutualistic connections with mycologists,
381 appreciating fungi as a crucial part of nature that needs to be taken into account in our efforts
382 to conserve biodiversity on Earth.

383

384 **Acknowledgements**

385 We are grateful to the European Section of the Society for Conservation Biology for giving
386 us the opportunity to organize a symposium on fungal conservation on the 3rd European
387 Conference of Conservation Biology in Glasgow, Scotland (August 2012), which launched
388 the discussion presented in this article. We also thank M. Ainsworth and two anonymous
389 reviewers for their valuable input to this paper. During the preparation of the manuscript the
390 Aage V. Jensen Foundation supported the first author. The participation of E.S. Barron was
391 supported by the US National Science Foundation under Grant No. 1127269.

392

393 **References**

394 Aldea, J., F. Martínez-Peña, and L. Diaz-Balteiro. 2012. Integration of fungal production in
395 forest management using a multi-criteria method. *European Journal of Forest*
396 *Research* **131**:1991–2003.

397 Arnolds, E. 1992. The analysis and classification of fungal communities with special
398 reference to macrofungi. Pages 7–47 in W. Winterhoff, editor. *Fungi in Vegetation*
399 *Science*. Kluwer Academic Publishers, Dordrecht.

400 Arnolds, E. 2001. The future of fungi in Europe: threats, conservation and management.
401 Pages 64–80 in D. Moore, M. N. Nauta, S. E. Evans, and M. Rotheroe, editors. *Fungal*
402 *conservation, issues and solutions*. Cambridge University Press, Cambridge.

403 Bailarote, B. C., B. Lievens, and H. Jacquemyn. 2012. Does mycorrhizal specificity affect
404 orchid decline and rarity? *American Journal of Botany* **99**:1655–1665.

405 Blaaid, R., T. Carlsen, S. Kumar, R. Halvorsen, K. I. Ugland, G. Fontana, and H. Kauserud.
406 2012. Changes in the root-associated fungal communities along a primary succession
407 gradient analysed by 454 pyrosequencing. *Molecular Ecology* **21**:1897–1908.

408 Blackwell, M. 2011. The Fungi: 1, 2, 3 ... 5.1 million species? *American Journal of Botany*
409 **98**:426–438.

410 Boa, E. 2004. Wild edible fungi: A global overview of their use and importance to people.
411 *Non-Wood Forest Products 17*. FAO, Rome.

412 Boddy, L., J. C. Frankland, and P. van West, editors. 2008. *Ecology of Saprotrophic*
413 *Basidiomycetes*. Elsevier, Amsterdam.

414 Caesar-Tonthat, T. C. 2002. Soil binding properties of mucilage produced by a basidiomycete
415 fungus in a model system. *Mycological Research* **106**:930–937.

416 CBD (Convention on Biological Diversity). 2010. *Convention on Biological Diversity, COP*
417 *10 Decision X/2: Strategic Plan for Biodiversity 2011–2020*. Available from
418 <http://www.cbd.int/decision/cop/?id=12268>> (accessed 11 December 2012).

419 Chapin, F. S., P. A. Matson, and H. A. Mooney 2011. Principles of terrestrial ecosystems
420 ecology. Springer, New York.

421 Claridge, A. W., and T. W. May. 1994. Mycophagy among Australian mammals. Australian
422 Journal of Ecology **19**:251–275.

423 Conti, M. E., and G. Cecchetti. 2001. Biological monitoring: lichens as bioindicators of air
424 pollution assessment - a review. Environmental Pollution **114**:471–492.

425 Coppins, A. M., and B. J. Coppins 2002. Indices of ecological continuity for woodland
426 epiphytic lichen habitats in the British Isles. British Lichen Society, London.

427 Dahlberg, A., D. R. Genney, and J. Heilmann-Clausen. 2010. Developing a comprehensive
428 strategy for fungal conservation in Europe: current status and future needs. Fungal
429 Ecology **3**:50–64.

430 Dahlberg, A., and G. M. Mueller. 2011. Applying IUCN red-listing criteria for assessing and
431 reporting on the conservation status of fungal species. Fungal Ecology **4**:147–162.

432 Egli, S., M. Peter, C. Buser, W. Stahel, and F. Ayer. 2006. Mushroom picking does not
433 impair future harvests - results of a long-term study in Switzerland. Biological
434 Conservation **129**:271–276.

435 Emery, M. R., and E. S. Barron. 2010. Using local ecological knowledge to assess morel
436 decline in the US Mid-Atlantic region. Economic Botany **64**:205–216.

437 Erjavec, J., J. Kos, M. Ravnikar, T. Dreo, and J. Sabotic. 2012. Proteins of higher fungi -
438 from forest to application. Trends in Biotechnology **30**:259–273.

439 Fisher, M. C., D. A. Henk, C. J. Briggs, J. S. Brownstein, L. C. Madoff, S. L. McCraw, and
440 S. J. Gurr. 2012. Emerging fungal threats to animal, plant and ecosystem health.
441 Nature **484**:186–194.

442 Gange, A. C., E. G. Gange, T. H. Sparks, and L. Boddy. 2007. Rapid and recent changes in
443 fungal fruiting patterns. Science **316**:71–71.

444 Griffith, G. W. 2012. Do we need a global strategy for microbial conservation? *Trends in*
445 *Ecology & Evolution* **27**:1–2.

446 Griffith, G. W., et al. 2013. The international conservation importance of Welsh 'waxcap'
447 grasslands. *Mycosphere* **4**:969–984.

448 Halme, P., J. Heilmann-Clausen, T. Rämä, T. Kosonen, and P. Kunttu. 2012. Monitoring
449 fungal biodiversity - towards an integrated approach. *Fungal Ecology* **5**:750–758.

450 Hawksworth, D. L. 2012. Global species numbers of fungi: are tropical studies and molecular
451 approaches contributing to a more robust estimate? *Biodiversity and Conservation* **21**:
452 2425–2433.

453 Heilmann-Clausen, J., and J. Vesterholt. 2008. Conservation: selection criteria and
454 approaches. Pages 325–347 in L. Boddy, J. C. Frankland, and P. van West, editors.
455 *Ecology of Saprotrophic Basidiomycetes*. Elsevier, Amsterdam.

456 Hibbett, D. S., A. Ohman, D. Glotzer, M. Nuhn, P. Kirk, and R. H. Nilsson. 2011. Progress in
457 molecular and morphological taxon discovery in Fungi and options for formal
458 classification of environmental sequences. *Fungal Biology Reviews* **25**:38–47.

459 Ingram, D. S. 1999. Biodiversity, plant pathogens and conservation. *Plant Pathology* **48**:433–
460 442.

461 IUCN (International Union for Conservation of Nature). 2013. Fungi. Available from
462 [http://iucn.org/about/work/programmes/species/who_we_are/ssc_specialist_groups_a](http://iucn.org/about/work/programmes/species/who_we_are/ssc_specialist_groups_and_red_list_authorities_directory/fungi/)
463 [nd_red_list_authorities_directory/fungi/](http://iucn.org/about/work/programmes/species/who_we_are/ssc_specialist_groups_and_red_list_authorities_directory/fungi/) (accessed 25 July 2013).

464 Kauserud, H., et al. 2012. Warming-induced shift in European mushroom fruiting phenology.
465 *Proceedings of the National Academy of Sciences of the United States of America*
466 **109**:14488–14493.

467 Komonen, A. 2003. Hotspots of insect diversity in boreal forests. *Conservation Biology*
468 **17**:976–981.

469 Kubartová, A., E. Ottosson, A. Dahlberg, and J. Stenlid. 2012. Patterns of fungal
470 communities among and within decaying logs, revealed by 454 sequencing.
471 *Molecular Ecology* **21**:4514–4532.

472 Lilleskov, E. A., E. A. Hobbie, and T. R. Horton. 2011. Conservation of ectomycorrhizal
473 fungi: exploring the linkages between functional and taxonomic responses to
474 anthropogenic N deposition. *Fungal Ecology* **4**:174–183.

475 Lindahl, B. D., et al. 2013. Fungal community analysis by high-throughput sequencing of
476 amplified markers - a user's guide. *New Phytologist* **199**:288–299.

477 Loh, J., R. E. Green, T. Ricketts, J. Lamoreux, M. Jenkins, V. Kapos, and J. Randers. 2005.
478 The Living Planet Index: using species population time series to track trends in
479 biodiversity. *Philosophical Transactions of the Royal Society B-Biological Sciences*
480 **360**:289–295.

481 Loo, J. 2009. Ecological impacts of non-indigenous invasive fungi as forest pathogens.
482 *Biological Invasions* **11**:81–96.

483 Mace, G. M., K. Norris, and A. H. Fitter. 2012. Biodiversity and ecosystem services: a
484 multilayered relationship. *Trends in Ecology & Evolution* **27**:19–26.

485 McKenzie, E. H. C., and P. R. Johnston. 2004. *Puccinia embergeriae* sp nov on Chatham
486 Islands sow thistle (*Embergeria grandifolia*) and a note on *Miyagia pseudosphaeria* on
487 sow thistles (*Sonchus* spp.) in New Zealand. *New Zealand Journal of Botany* **42**:657–
488 661.

489 Minter, D. 2010. Safeguarding the future. Pages 143–153 in L. Boddy, and M. Coleman,
490 editors. *From another kingdom: the amazing world of Fungi*. Royal Botanic Garden
491 Edinburgh, Edinburgh.

492 Nordén, J., R. Penttilä, J. Siitonen, E. Tomppo, and O. Ovaskainen. 2013. Specialist species
493 of wood-inhabiting fungi struggle while generalists thrive in fragmented boreal
494 forests. *Journal of Ecology* **101**:701–712.

495 Ovaskainen, O., D. Schigel, H. Ali-Kovero, P. Auvinen, L. Paulin, B. Nordén, and J. Nordén.
496 2013. Combining high-throughput sequencing with fruit body surveys reveals
497 contrasting life-history strategies in fungi. *ISME Journal* **7**:1696–1709.

498 Pan, Y. D., et al. 2011. A large and persistent carbon sink in the world's forests. *Science*
499 **333**:988–993.

500 Parker, S. S. 2010. Buried treasure: soil biodiversity and conservation. *Biodiversity and*
501 *Conservation* **19**:3743–3756.

502 Pautasso, M., G. Aas, V. Queloz, and O. Holdenrieder. 2013. European ash (*Fraxinus*
503 *excelsior*) dieback - A conservation biology challenge. *Biological Conservation*
504 **158**:37–49.

505 Pennant, T. 1781. *Tours in Wales*, Vol. 3. Benjamin White, London.

506 Pounds, J. A., et al. 2006. Widespread amphibian extinctions from epidemic disease driven
507 by global warming. *Nature* **439**:161–167.

508 Pringle, A., E. Barron, K. Sartor, and J. Wares. 2011. Fungi and the Anthropocene:
509 Biodiversity discovery in an epoch of loss. *Fungal Ecology* **4**:121–123.

510 Rambold, G., M. Stadler, and D. Begerow. 2013. Mycology should be recognized as a field
511 in biology at eye level with other major disciplines - a memorandum. *Mycological*
512 *Progress* **12**:455–463.

513 Rassi, P., Hyvärinen, E., Juslén, A., Mannerkoski, I. (eds.) 2010. *The 2010 Red List of*
514 *Finnish species*. Edita, Helsinki. 685 pp.

515 Raunio, A., Schulman, A., Kontula, T. (eds.) 2008. *Suomen luontotyyppien uhanalaisuus*
516 *(Assessment of threatened habitat types in Finland)*. Finnish Environment Institute,
517 Helsinki.

518 Rodriguez, R. J., J. F. White, A. E. Arnold, and R. S. Redman. 2009. Fungal endophytes:
519 diversity and functional roles. *New Phytologist* **182**:314–330.

520 Rosenvald, R., and A. Lõhmus. 2008. For what, when, and where is green-tree retention
521 better than clear-cutting? A review of the biodiversity aspects. *Forest Ecology and*
522 *Management* **255**:1–15.

523 Rutter, G. 2010. Fungi and humanity. Pages 93–103 in L. Boddy, and M. Coleman, editors.
524 *From another kingdom: the amazing world of Fungi*. Royal Botanic Garden
525 Edinburgh, Edinburgh.

526 Sala, O. E., et al. 2000. Global biodiversity scenarios for the year 2100. *Science* **287**:1770–
527 1774.

528 Scheffers, B. R., L. N. Joppa, S. L. Pimm, and W. F. Laurance. 2012. What we know and
529 don't know about Earth's missing biodiversity. *Trends in Ecology & Evolution*
530 **27**:501–510.

531 Scheidegger, C., and S. Werth. 2009. Conservation strategies for lichens: insights from
532 population biology. *Fungal Biology Reviews* **23**:55–66.

533 Schmit, J. P., and G. M. Mueller. 2007. An estimate of the lower limit of global fungal
534 diversity. *Biodiversity and Conservation* **16**:99–111.

535 Schoch, C. L., et al. 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a
536 universal DNA barcode marker for Fungi. *Proceedings of the National Academy of*
537 *Sciences of the United States of America* **109**:6241–6246.

538 Scholes, R. J., and R. Biggs. 2005. A biodiversity intactness index. *Nature* **434**:45–49.

539 Senn-Irlet, B., J. Heilmann-Clausen, D. Genney, and A. Dahlberg. 2007. Guidance for
540 conservation of macrofungi in Europe. A document prepared for the European
541 Council for Conservation of Fungi (ECCF) within the European Mycological

542 Association (EMA) and the Directorate of Culture and Cultural and Natural Heritage,
543 Council of Europe, Strasbourg.

544 Shaver, G. R., and F. S. Chapin. 1991. Production - biomass relationships and element
545 cycling in contrasting Arctic vegetation types. *Ecological Monographs* **61**:1–31.

546 Smith, S. E., and D. J. Read 2008. *Mycorrhizal symbiosis*. Academic Press, Amsterdam.

547 Stokland, J. N., J. Siitonen, and B. G. Jonsson 2012. *Biodiversity in dead wood*. Cambridge
548 University Press, Cambridge, UK.

549 Tisdall, J. M., S. E. Nelson, K. G. Wilkinson, S. E. Smith, and B. M. McKenzie. 2012.
550 Stabilisation of soil against wind erosion by six saprotrophic fungi. *Soil Biology &*
551 *Biochemistry* **50**:134–141.

552 van der Linde, S., E. Holden, P. I. Parkin, I. J. Alexander, and I. C. Anderson. 2012. Now you
553 see it, now you don't: The challenge of detecting, monitoring and conserving
554 ectomycorrhizal fungi. *Fungal Ecology* **5**:633–640.

555 World Resources Institute. 2005. *Millennium Ecosystem Assessment. Ecosystems and human*
556 *well-being: Biodiversity synthesis*. World Resources Institute, Washington, DC.

557

558

559

560 Figure 1. Four examples emphasizing how fungi provide added value in biodiversity
561 conservation: (1) They provide and give direct insight into important supporting ecosystem
562 services including nutrient cycling, and mycorrhizal symbiosis that enhance plant nutrition
563 and resistance to drought, soil pollution and pathogens (A, Three different ectomycorrhizas
564 on European Beech (*Fagus sylvatica* L.)). (2) They are useful as indicators when evaluating
565 the conservation potential of conservation areas or the conservation outcome of conducted
566 management actions (B, *Hygrocybe punicea* (Fr.) P. Kumm., a waxcap species that is
567 commonly used as an indicator of grassland sites with high conservation value). (3) They
568 play an important role in developed countries in providing recreational values and
569 reconnecting urban citizens with nature (C, A family collecting fungi for food and learning
570 about their identification, near Copenhagen, Denmark). (4) They provide a sustainable
571 income from intact forests for the local people in developing countries and can thus play a
572 role in turning local attitudes positive towards conservation areas (D, women selling fruit
573 bodies of native mycorrhizal fungi in a street market in Zambia). Photo courtesy of Jens H.
574 Petersen (A), Nigel Bean (B), Flemming Rune (C), Marja Härkönen (D).