



## **Nano-sized polystyrene affects feeding, behavior and physiology of brine shrimp *Artemia franciscana* larvae**

This is a pre print version of the following article:

*Original:*

Bergami, E., Bocci, E., Vannuccini, M.L., Monopoli, M., Salvati, A., Dawson, K.A., et al. (2015). Nano-sized polystyrene affects feeding, behavior and physiology of brine shrimp *Artemia franciscana* larvae. *ECOTOXICOLOGY AND ENVIRONMENTAL SAFETY*, 123, 18-25 [10.1016/j.ecoenv.2015.09.021].

*Availability:*

This version is available <http://hdl.handle.net/11365/982091> since 2017-05-23T11:49:17Z

*Published:*

DOI:10.1016/j.ecoenv.2015.09.021

*Terms of use:*

Open Access

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.

(Article begins on next page)

Manuscript Number: EES-15-605R1

Title: Nano-sized polystyrene affects feeding, behaviour and physiology of brine shrimp *Artemia franciscana* larvae

Article Type: SI: BECOME Meeting

Section/Category: Ecotoxicology

Keywords: Nanoplastics; polystyrene; marine zooplankton; *Artemia franciscana*; accumulation; molting

Corresponding Author: Dr. Ilaria Corsi, Ph.D.

Corresponding Author's Institution: Università degli Studi di Siena

First Author: Elisa Bergami, Master degree

Order of Authors: Elisa Bergami, Master degree; Elena Bocci, Ph.D.; Maria Luisa Vannuccini, Master degree; Marco Monopoli, Ph.D.; Anna Salvati, Ph.D.; Kenneth A Dawson, Ph.D.; Ilaria Corsi, Ph.D.

Abstract: Nano-sized polymers as polystyrene (PS) constitute one of the main challenges for marine ecosystems, since they can distribute along the whole water column affecting planktonic species and consequently disrupting the energy flow of marine ecosystems. Nowadays very little knowledge is available on the impact of nano-sized plastics on marine organisms. Therefore, the present study aims to evaluate the effects of 40 nm anionic carboxylated (PS-COOH) and 50 nm cationic amino (PS-NH<sub>2</sub>) polystyrene nanoparticles (PS NPs) on brine shrimp *Artemia franciscana* larvae. No signs of mortality were observed at 48 h of exposure for both PS NPs at nauplius stage but several sub-lethal effects were evident. PS-COOH (5-100 µg/ml) resulted massively sequestered inside the gut lumen of larvae (48h) probably limiting food intake. Some of them were lately excreted as fecal pellets but not a full release was observed. Likewise, PS-NH<sub>2</sub> (5-100 µg/ml) accumulated in larvae (48h) but also adsorbed at the surface of sensorial antennules and appendages probably hampering larvae motility. In addition, larvae exposed to PS-NH<sub>2</sub> undergo multiple molting events during 48h of exposure compared to controls. The activation of a defense mechanism based on a physiological process able to release toxic cationic NPs (PS-NH<sub>2</sub>) from the body can be hypothesized. The general observed accumulation of PS NPs within the gut during the 48h of exposure indicates a continuous bioavailability of nano-sized PS for planktonic species as well as a potential transfer along the trophic web. Therefore, nano-sized PS might be able to impair food uptake (feeding), behavior (motility) and physiology (multiple molting) of brine shrimp larvae with consequences not only at organism and population level but on the overall ecosystem based on the key role of zooplankton on marine food webs.



UNIVERSITÀ  
DI SIENA  
1240

*Università degli Studi di Siena*

*Dipartimento di Scienze Fisiche, della Terra e dell'Ambiente*

Siena, 6<sup>th</sup> September 2015

Dear Editor

Please find enclosed the electronic submission of our revised manuscript entitled:

***Nano-sized polystyrene affects feeding, behaviour and physiology of brine shrimp *Artemia franciscana* larvae***

The main subject of the manuscript is to investigate distribution and sub-lethal effects of carboxylated (PS-COOH) and amine (PS-NH<sub>2</sub>) polystyrene nanoparticles as model nanoplastics in brine shrimp *Artemia salina* in order to determine whether these materials have similar effects in marine organisms as what observed in common human and other mammalian cell lines.

Nanoplastic debris, resulted from run-off and weathering breakdown of macro and microplastics, represent an emerging concern for marine ecosystems. While microplastics are quite well studied, fate and impact of nanoscale plastics in the marine environment is almost unknown and this is raising concern due to the increasing abundance in water column and food webs and nanoscale properties which could imply toxicity to marine biota. Polystyrene NPs can be considered as good model for studying both environmental fate of nanoplastics, in terms of interactions with the surrounding media, and toxicity for marine organisms focusing on specific pathways of cellular uptake. In fact the toxicity mechanisms have been quite well described in human cell models, while how PS NPs can interact with their surroundings as for instance with marine waters and enter the cells of marine organisms is still largely unknown.

Nanoparticle stability in natural seawater was measured by DLS, while distribution and toxicity (mortality) were monitored through light and fluorescence microscopy within 48 h of exposure.

The detailed secondary characterization performed in the present study clearly showed that in natural seawater brine shrimp larvae are exposed mainly to nanoscale objects in the case of PS-NH<sub>2</sub> (< 100 nm)



UNIVERSITÀ  
DI SIENA  
1240

*Università degli Studi di Siena*

*Dipartimento di Scienze Fisiche, della Terra e dell'Ambiente*

while PS-COOH NPs originated microscale aggregates (> 1000 nm).

Our findings suggest that the different aggregation of the two tested PS NPs in natural sea water (89 nm for PS-NH<sub>2</sub> and PDI > 0.4 for PS-COOH) and, more importantly, the different surface charges might affect their cellular uptake and distribution and consequent sub-lethal effects in brine shrimp larvae.

Our results showed that brine shrimp larvae might be vulnerable to the amino modified PS NPs, as observed for mammalian cell lines and confirmed also on our recent work on sea urchin embryos (Della Torre et al., 2014). PS-COOH NPs did not showed any relevant effect but a significant accumulation inside the gut (as well as PS NH<sub>2</sub>) was observed and this may also determine transfer through the trophic food web, raising serious concern about the exposure of organisms at higher trophic levels.

Furthermore, our study also show that careful assessment of NP properties and stability in natural sea water media is needed in order to properly determine their characteristics once exposed to marine organisms (e.g. plankton), as different aggregation state may lead to different uptake and distribution routes, and the different surface properties clearly have different impact.

We hope that the revised version of the ms could comply with the standards of Ecotoxicology and Environmental Safety journal and will be considered for publication in the special issue: BECOME meeting. The authors state that there is no conflict of interest.

Many thanks in advance for your consideration.

Your Sincerely

Ilaria Corsi

Ilaria Corsi, PhD  
Adjunct Professor of Ecotoxicology  
Department of Physical Earth and Environmental Sciences  
University of Siena  
via Mattioli, 4  
53100 Siena Italy  
Phone: +39 0577 232830  
Fax: +39 0577 232806  
E-mail: [ilaria.corsi@unisi.it](mailto:ilaria.corsi@unisi.it)

**Reviewer suggestions**

**Melanie Auffan, PhD**

Cerege, France

[auffan@cerege.fr](mailto:auffan@cerege.fr)

**Laura Canesi, PhD**

University of Genova, Italy

[laura.canesi@unige.it](mailto:laura.canesi@unige.it)

**Zikri Arslan, PhD**

Jackson State University, USA

[zikri.arslan@jsums.edu](mailto:zikri.arslan@jsums.edu)



*Università degli Studi di Siena*

*Dipartimento di Scienze Fisiche, della Terra e dell'Ambiente*

Siena, 6<sup>th</sup> September 2015

**Journal: Ecotoxicology and Environmental Safety**

Ms. Ref. No EES-15-605

Title: Nano-sized polystyrene affects feeding, behaviour and physiology of brine shrimp *Artemia franciscana* larvae

Author(s): Elisa Bergami, Elena Bocci, Maria Luisa Vannuccini, Marco P. Monopoli, Anna Salvati, Kenneth A Dawson, Ilaria Corsi

Dear Editor,

We do thank you and the two Reviewers for their comments and criticisms.

The manuscript (ms) had been modified accordingly following the comments.

Reviewer #3: From my point of view this paper can be presented as original research article because overall the obtained results fall to reach the purposes indicated by the authors: i) evaluate the effects of 40 nm anionic carboxylated (PS-COOH) and 50 nm cationic amine (PS-NH<sub>2</sub>) polystyrene nanoparticles on brine shrimp *Artemia* larvae by acute mortality test; ii) evaluate the uptake and distribution of these NPs on the brine shrimp, also observing sub-lethal effects during the mortality test.

In my opinion, the original finding of this paper is the observation of accumulation, excretion, adherence at the surface of sensorial antennulae and appendages, and molting induction.

I evaluate positively the tentative quantitative assessment of molts released by PS-NH<sub>2</sub> exposed larvae, by separation of larvae from molts which were quantified by gravimetry.

In addition, physico-chemical characterization of PS-COOH and PS-NH<sub>2</sub> NPs is also reported

using Dynamic Light Scattering analysis and showing Z-average (nm), polydispersity index (PDI) and <zeta>-potential (mV).

*We thank the reviewer for having appreciated the manuscript and the quality of the data presented.*

The manuscript is well written and I agree to accept it with minor revisions as follow:

- In general: in many parts of the text spaces between words are missing and sometimes the verb is not correctly included in the singular or plural form.

*All spaces have been inserted and verbs have been checked carefully and corrected accordingly.*

- Lines 194-195: Artemia salina species is reported in the text. Do authors want to refer to Artemia franciscana?

*Thanks to advices provided by zoologists experts on Artemia genus, we now are confident in indicating the Artemia larve used in our study as franciscana and not salina as previously stated. So both in the title and in the ms the species has been changed from salina to franciscana. We also decided to include the word "larvae" in order to make more clear the biological model used. We do thanks the reviewer for such criticism since perhaps we did not checked properly the origin and attributes of the larvae used in the study.*

- Lines 195-196 "were purchased from Laboratory ... " It should be "... from the company MicroBioTests (Ghent, Belgium)"

*It has been changed according to reviewer suggestion.*

- Lines 226, 285, Recovery experiment: has it been carried out according to Ates et al (2013a) (as reported at line 227) or to Auffan et al (2013) (as reported in table 2)?

*Thanks, there was a mistake in the citation, it has been now changed as Ates et al 2013a*

- Line 221-222, table 2, Accumulation analysis: has it been carried out according to APAT IRSA CNR 2003 (as reported in Table 2) or by optical fluorescent microscope (as reported at lines 221-222)? I seem that this method is only for the mortality test and it doesn't provide an accumulation outcome. Has accumulation been only observed or quantified?

*The reviewer is right, the cited method is only a mortality test and this is what we meant in Table 2 by including this citation. Based on the strong accumulation we previously observed in another model organism as sea urchin (see Della Torre et al., 2015) we decided to observed brine shrimp larvae under optical fluorescence microscopy. We are now working on a protocol for quantify the amount of fluorescence in order to get a dose-response curve and perhaps to cite in our future ms. In order to clarify which method has been used for mortality and accumulation, we have inserted the following sentence in the text after line 211.*

*"Moreover, at 48 h nauplii were also observed by optical fluorescence microscopy in order to identify any sub-lethal effects (see Table 2) including molting, the presence of PS NPs accumulated in the digestive tract or adhering to the external appendages."*

- Line 322: verify "in press"  
It has been changed as Della Torre et al., 2015.

- Lines 377-380, figure 1 : I don't see image d)  
*The “d” image is inside the “c” as spot in the corner.*

Reviewer #1: The authors intends to show the importance of the effects provoked by the presence of nano-plastics in the environment, by using two differently charged polystyrene NPs. Despite the importance of the observation and the study of these NPs in particular (most probably of high importance in a very near future), is missing the physicochemical characterization of the natural waters samples on the Tuscan archipelago, which will undoubtedly help on the understanding of the effects observed. So, I advise the authors to show this characterization and use it to discuss the effects obtained. Even, because the authors say at least twice that is very important to understand the properties of the particles when dispersed in the medium used for the toxicity tests.

*We thank the reviewer for such criticism, a new Table indicated as “Table 3” has been included in the ms showing physico-chemical parameters of natural sea water (NSW) samples used in the present study including also selected contaminants.*

Beside this main comment some other major comments need to be clarified before publication.

- 1) Numerous type errors can be found through all text; most of them related with the absence of a space between words. Some examples are: 4<sup>th</sup> highlight “PS-NH2 were...”; Abbreviations “... (PS-COOH), amino...”; Line 69 “... 2014). Concerning...”, ...

*All ms has been checked carefully and all type-errors corrected accordingly. We realized that this problem might be probably related to have opened the file with two different version of Microsoft Word program.*

- 2) Line 72. Should read “... sized, since EU...” and not “... sized in this area, since EU...”

*Thanks, changed.*

- 3) Lines 72-73. “Good environmental status”. This is very vague, doesn’t mean a lot.

*The main goal of the European Marine Directive (Directive 2008/56/EC) is to achieve Good Environmental Status of EU marine waters by 2020. The Directive defines Good Environmental Status (GES) as:*

*“The environmental status of marine waters where these provide ecologically diverse and dynamic oceans and seas which are clean, healthy and productive” from Article 3 of Directive 2008/56/EC. In order to better explain such citation, the sentence has been rewritten and the Directive citation included.*

- 4) Lines 80-81. The authors could give the measured/expected quantities and also the intrinsic properties that make these particles important to be studied.

*Unfortunately there are no data available regarding neither measured nor expected quantities of polystyrene nanoparticles in sea water samples. Please refer to the ms of Besseling et al., 2014 (cited) in which a sort of assessment has been done but only in freshwater and not in sea water. Concerning the intrinsic properties please refer to the several papers of co-authors of the ms, Monopoli and Dawson, cited in the introduction from line 111 to 119.*

- 5) Line 86. Should read “... expected that they would be more severely exposed to nano-sized... and not “... expected to be more severely exposed even to nano-sized...”



*Many thanks for the advice, it has been changed*

- 6) Lines 89-90. This sentence is very confuse and I really cannot understand what the authors want to say with this.

*Thanks for the criticism, the sentence has been rewritten in two new sentences better explaining the concept of behavior of PS NPs in sea water column as consequence of aggregation.*

- 7) Line 94. “selected” why not to give the composition of the NPs?

*The composition has been included in the lines 111-114 of the ms.*

- 8) Lines 98-119. Both paragraphs should be after the sentence on line 81.

*Thanks for the criticism, the sentences have been moved after line 81.*

- 9) Line 125. Should read “... embryos, which were the first evidence of toxicity reported...” and not “... embryos, where also first evidence of toxicity were reported...”

*Thanks. Changed.*

- 10) Line 131. What are the standard PS?

*We perhaps used the word “standard” not properly, we referred to unmodified PS NPs, without surface functionalization.*

- 11) Line 132. Should read “... determined yet.” And not “... determined as yet.”

*Thanks. Changed.*

- 12) Lines 132-134. What the authors meant with this? Is necessary to rewritten this sentence.

*Thanks, it was not well written since the verb was in the wrong position. It has been rewritten.*

- 13) Line 147. Should read “The aim of the present...” and not “The aim of present...”

*Thanks. Changed.*

- 14) Lines 147-149. It would make more sense to use the exact same size of the NPs in order to evaluate only the effect of the charge. When thinking in surface area 10 nm in difference is big and may cause very different effects; what is the difference in the quantity of groups at the surface of the particle?

*To our knowledge , the only available labeled PS-NPs are those of 40 nm PS-COOH and 50 nm PS-NH2. Past studies on human models tested the same PS-NPs (see cited papers of co-authors Dawson and Monopoli) and such slight difference in dimension has not been considered so important in term of observed toxicity. Being aggregation quite different between the two in NSW, we assume that even being different in primary dimension, once in NSW this difference may not play a significant role in term of groups at the surface of the aggregates. Nevertheless, we will keep this interesting suggestion for future studies.*

- 15) Lines 163-169. In the way that is written it seems that is important to study these nano-plastics because they are NOT present in the environment! So, why to accept this manuscript, if it doesn't have environmental relevance? This needs to be rewritten.

*Thanks the suggestion, the sentence has been rewritten as follows: “Although negative surface charged nanoplastics as PS-COOH have been suggested as the most widespread in the environment (Besseling et al., 2014), no data are currently on their fate and toxicity to marine biota”.*

- 16) Lines 176-179. As mentioned above, is necessary to characterize these waters in terms of organic and inorganic content and metals quantities; the toxic effects can come from the presence of metals.

*As mentioned above a new Table listing physico-chemical parameters including levels of metals of natural sea water samples used in the study has been included in the ms.*

- 17) Lines 249-250. The TEM images shows that the sizes of the particles are not so similar; so, its important to give the average size and the standard deviations; also, the number of particles counted and in how many pictures.

*The average size value of the particles has been included in legend of Figure S1. Regarding their measure it has been calculated using at least 10 differences pictures from each particles and by calculating the average values.*

- 18) Lines 249-250. Why the TEM analyses were not performed for the NPs dispersed in the natural waters? (so, the exact same conditions where the tox experiments were performed.) Moreover, as the authors surely know that the DLS have a bias through larger particles proportional to  $d^6$ . So, its important that the aggregation obtained by DLS would be confirmed by TEM, and even because it seems that the authors have access to this instrument.

*The TEM images were performed in both MilliQ water for primary characterization as well as in NSW for DLS data confirmation. Unfortunately, the background noise of NSW probably due to high ionic salts and NOM made almost impossible to recognize the shape of the PS-NPs and more important to measure their dimension. Due to this very low quality of TEM images and being useless for our study, we decided not to include them in the ms. Meanwhile, we are currently working to solve this issue but it seems a very difficult task for this specific type of NP. We published other papers on titanium dioxide NPs in which we easily showed good TEM images obtained in NSW samples of the same geographical origin (Della Torre et al., doi10.1016/j.hazmat.2015.04.072; Canesi et al. doi10.1016/j.aquatox.2013.11.002).*

- 19) Lines 260-263. How the authors can affirm this if they don't show the characterization data of the natural waters? Also, this explanation will be dependent on the organic matter content.

*Our hypothesis is based on recent studies in which the NOM present in NSW has been shown to drive heteroaggregation phenomenon of surface charged NPs so potentially affecting also the observed aggregation of PS NPs in NSW observed our study. The total organic carbon content listed now in Table 3 show a moderate level of organic matter in NSW samples used in or study so confirming such hypothesis and in agreement with other studies (please see Wang et al., 2013b, cited in the ms).*

- 20) Lines 268-273. This paragraph is totally redundant since the authors didn't do this full characterization and even don't give the characteristics of the water; also, most of this is repeated in the paragraph above.

*Again we were referring to data showed by Wang et al. 2013b but based on our revision and data included in the new Table 3, a confirmation of a role of such parameters in the behavior of such NPs in NSW has been given. The sentence from 271-273 is just describing the different dimension of PS-COOH and PS-NH2 aggregates observed in NSW.*

21) Table 2 is never called in the main text body.

*It has been called in line 202.*

22) Line 301. During how much time this recording was done?

*The recording was done for the further 24 h (recovery test).*

23) Line 302. Should read "... (see Figure S2 and the video on the Supplementary..." and not (see Supplementary..."

*Thanks. Changed.*

24) Lines 317-318. Why this was not done within this study? It seems feasible.

*This is the aim of the running experiments which are now about to end and will be published hopefully soon.*

25) Lines 319-321. What this has to do with the issue being described and discussed in all this paragraph?

*This sentence aim to address the ecological impact of potential effect of nanoplastics in planktonic species and the consequent impact on marine ecosystem due their key role in trophic web.*

26) Lines 340-341. Please, rephrase this sentence.

*The sentence has been rewritten.*

27) Lines 351-353. This is said in the caption of Figure 3, and in fact it should be only here.

*It has been deleted from the text.*

28) Line 353. Should read "... several times..." and not "... several time..."

*Thanks. Changed.*

29) Line 354. Should read "... and in the three..." and not "... and the three..."

*Thanks. Changed.*

30) Line 358. Should read "... crucial step in the..." and not "... crucial in the..."

*Thanks. Changed.*

31) Line 404. So, why these studies were done in absence of food? Also, if there is already evidences that perhaps is necessary to wait more time, why these studies were done only during 24 h?

*Based on our experience and in agreement with previous studies (see Garaventa et al., 2010), 24 h without food at such developmental stage is the maximum time allowed in order not to affect organism survival. Moreover, in order to maintain the same experimental media used in the acute test (48h) we decided not to add any food in the media which could potentially affect also PS NPs*

*behaviour by for instance increasing organic matter content. A long-term toxicity test with food is currently under investigation and data will be ready soon.*

32) Lines 406-408. Any idea why this would happen? Maybe some heteroagglomeration with organic or inorganic colloids present in the water (which cannot be attested since no detail characterization of the water is given)?

*Based on the new data showed on Table 3 on NSW samples, such behaviour might be due to a better dispersion in NSW of positively charged PS NPS as PS-NH<sub>2</sub> than negatively ones PS-COOH) in accordance with the hypothesis of heteroaggregation as reviewer also suggest.*

We do hope that changes made following the suggestions of the reviewers have improved the quality of the manuscript so as to allow the publication in *Ecotoxicology and Environmental Safety* journal.


The authors state that there is no conflict of interest.

Your Sincerely



Ilaria Corsi

Ilaria Corsi, PhD  
Adjunct Professor of Ecotoxicology  
Department of Physical Earth and Environmental Sciences  
University of Siena  
via Mattioli, 4  
53100 Siena Italy  
Phone: +39 0577 232830  
Fax: +39 0577 232806  
E-mail: [ilaria.corsi@unisi.it](mailto:ilaria.corsi@unisi.it)  
Websites: <http://goo.gl/TD7pFS>  
<http://www.dsfta.unisi.it/en/departament/people/teaching-and-research-staff-eng/corsi-ilaria>



Accumulation and excretion



Adherence and molting induction

## Highlights

- Nowadays very little knowledge is available on the impact of nano-sized plastics on marine organisms.
- Polystyrene NPs (PS-COOH and PS-NH<sub>2</sub>) caused no mortality at 48 h of exposure in larvae of brine shrimp *A. franciscana larvae*, but several sub-lethal effects were observed.
- PS NPs resulted massively sequestered inside the gut lumen of larvae (48 h), probably limiting food intake.
- PS-NH<sub>2</sub> were adsorbed at the surface of sensorial antennulae and appendages.
- Multiple molting events during 48 h of exposure compared to controls were observed upon exposure to PS-NH<sub>2</sub>.

1 **Nano-sized polystyrene affects feeding, behavior and physiology of brine shrimp**  
2 *Artemia franciscana* larvae

3

4 Elisa Bergami<sup>1</sup>, Elena Bocci<sup>1</sup>, Maria Luisa Vannuccini<sup>1</sup>, Marco Monopoli<sup>2</sup>, Anna  
5 Salvati<sup>3</sup>, Kenneth A. Dawson<sup>2</sup>, Iliaria Corsi<sup>1\*</sup>

6

7 <sup>1</sup>Department of Physical, Earth and Environmental Sciences, University of Siena,  
8 Italy;

9 <sup>2</sup>Centre for BioNano Interactions, School of Chemistry and Chemical Biology,  
10 University College Dublin, Ireland;

11 <sup>3</sup>Groningen Research Institute of Pharmacy, Division Pharmacokinetics, Toxicology  
12 and Targeting, University of Groningen, The Netherlands

13

14

15

16

17

18

19

20 \*Corresponding Author

21 \* Phone (+39) 0577 232830; fax (+39) 0577 232806; e-mail: [ilaria.corsi@unisi.it](mailto:ilaria.corsi@unisi.it); mail: Department of  
22 Physical, Earth and Environmental Sciences, University of Siena, via Mattioli, 4, 53100 Siena, Italy

23

## 24 Abstract

25 Nano-sized polymers as polystyrene (PS) constitute one of the main challenges for marine  
26 ecosystems, since they can distribute along the whole water column affecting planktonic species  
27 and consequently disrupting the energy flow of marine ecosystems. Nowadays very little  
28 knowledge is available on the impact of nano-sized plastics on marine organisms. Therefore, the  
29 present study aims to evaluate the effects of 40 nm anionic carboxylated (PS-COOH) and 50 nm  
30 cationic amino (PS-NH<sub>2</sub>) polystyrene nanoparticles (PS NPs) on brine shrimp *Artemia franciscana*  
31 larvae. No signs of mortality were observed at 48 h of exposure for both PS NPs at nauplius stage but  
32 several sub-lethal effects were evident. PS-COOH (5-100µg/ml) resulted massively sequestered  
33 inside the gut lumen of larvae (48h) probably limiting food intake. Some of them were lately  
34 excreted as fecal pellets but not a full release was observed. Likewise, PS-NH<sub>2</sub> (5-100 µg/ml)  
35 accumulated in larvae (48h) but also adsorbed at the surface of sensorial antennules and appendages  
36 probably hampering larvae motility. In addition, larvae exposed to PS-NH<sub>2</sub> undergo multiple  
37 molting events during 48h of exposure compared to controls. The activation of a defense  
38 mechanism based on a physiological process able to release toxic cationic NPs (PS-NH<sub>2</sub>) from the  
39 body can be hypothesized. The general observed accumulation of PS NPs within the gut during the  
40 48h of exposure indicates a continuous bioavailability of nano-sized PS for planktonic species as  
41 well as a potential transfer along the trophic web. Therefore, nano-sized PS might be able to impair  
42 food uptake (feeding), behavior (motility) and physiology (multiple molting) of brine shrimp larvae  
43 with consequences not only at organism and population level but on the overall ecosystem based on  
44 the key role of zooplankton on marine food webs.

45

46

## 47 Keywords

48 Nanoplastics, polystyrene, zooplankton, *Artemia franciscana larvae*, accumulation, molting

49

50

## 51 Abbreviations

52 Polystyrene Nanoparticles (PS NPs), carboxylated (PS-COOH), amino (PS-NH<sub>2</sub>), Natural Sea  
53 Water (NSW), Natural Organic Matter (NOM)

54

55

56

57



## 59 **1. Introduction**

60 Several studies estimate that trillions of plastics are floating all over the oceans (Eriksen et al.,  
61 2014), representing one of the most important threats for marine ecosystems (Cozar et al., 2014).  
62 Micro (< 5 mm) and nanoplastics (< 100 nm) resulting also from weathering and fragmentation  
63 processes of macro-debris have been acknowledged as the most dangerous for marine wildlife since  
64 they might be easily ingested causing chemical and physical effects to marine organisms (Cole et  
65 al., 2013). Smaller plastic pieces can be uptaken and retained by small invertebrates, leading to  
66 bioaccumulation (Browne et al., 2011; Lee et al., 2013; Ward and Kach, 2009; Wright et al.,  
67 2013a), toxicity (Cole et al., 2014), but also trophic transfer to top-predators with potential impact  
68 for marine ecosystems as a whole (Farrell and Nelson, 2013; Setala et al., 2014; Watts et al., 2014).  
69 Concerning the Mediterranean basin, one thousand tons of plastic debris have been recently  
70 reported with a frequency comparable to the five subtropical ocean gyres (Cozar et al., 2015).  
71 Therefore, it is necessary to gain a deeper insight into the impact of small plastics including nano-  
72 sized, since EU member states must develop activities to reach “good environmental status, GES”  
73 by 2020, as main goals of the Marine Strategy Framework Directive ([Directive 2008/56/EC](#))  
74 (Galgani et al., 2010). Evidences of harmful effects of plastics is mostly restricted to observation on  
75 individual specimens and larger debris (micro and macroplastics) but concerns have been raised  
76 about physico-chemical effects of nanoplastics for single species up to ecosystem-wide impacts  
77 (UNEP, 2011). Both micro and nanoplastics are expected to increase consistently with time in the  
78 sea and oceans worldwide and important questions regarding sources, fate and biological effects  
79 need to be answered. While microplastics are quite well studied, the occurrence as well as effects of  
80 nano-sized particles are almost unknown and are raising concern due to expected increasing  
81 abundance in sea water (Cozar et al., 2014) and their intrinsic properties (Matranga and Corsi, 2012;  
82 Wright et al., 2013b). Polystyrene (PS) is one of the most largely used plastics worldwide, with an  
83 annual production of over 23 million tons per year (considering PS, high-impact PS and expanded  
84 PS) (Lithner et al., 2011). This polymer persists for several hundred years in the environment and  
85 undergoes to extremely slow depolymerization in marine waters (Andrady, 2003; Innocenti, 2003),  
86 thus leading to the formation of micro and nano-debris (Bandyopadhyay and Basak, 2007; Hofer,  
87 2008). Therefore, PS might pose a serious hazard to marine organisms due to the properties of the  
88 styrene monomer known as carcinogenic and endocrine disruptor (Lithner et al., 2011). These  
89 findings identified PS debris as potential multiple stressor in marine habitats, especially when  
90 available for ingestion by marine wildlife. Recent studies showed also higher sorption of Polycyclic

91 Aromatic Hydrocarbons (PAHs) and Polychlorinated Biphenyls (PCBs) to nano-sized PS compared  
92 to other plastics found in the marine environment (Rochman et al., 2013; Velzeboer et al., 2014).

93 In the last decades, PS based nanomaterials have been largely synthesized for several applications  
94 including packaging and nanomedicine (Bramini, 2014; Salvati et al., 2011; Silvestre et al., 2011).  
95 PS NPs refer to particles at nanoscale dimension with a PS core and variable functional groups,  
96 which determine their chemical reactivity and particle surface charge (Nowack and Bucheli, 2007).  
97 Common functionalized PS NPs include anionic carboxylated (-COOH) and cationic unsaturated  
98 amino (-NH<sub>2</sub>) (Casado et al., 2013; Loos et al., 2014). Several studies revealed their cellular uptake  
99 using *in vitro* models with human cell lines (Bramini, 2014; Lesniak et al., 2010; Liu et al., 2011;  
100 Lunov, 2011; Rossi et al., 2014), but also cytotoxicity and apoptosis in particular for PS-NH<sub>2</sub>  
101 (Bexiga et al., 2011; Wang et al., 2013a). Based on the current data, PS NPs uptake and toxicity  
102 depend on their intrinsic properties such as size and surface charges which affect their interaction  
103 with exposure media (Della Torre et al., 2014a).

104 Marine invertebrates are among the primary biological targets of nanoplastics, being exposed both  
105 to polymeric beads in suspension, as planktonic larvae, and to the fraction in sediments, as adult  
106 organisms (Manzo et al., 2013; Matranga and Corsi, 2012; Moore, 2006). Microplastics ingestion  
107 have been recently documented in several marine species including zooplankton therefore it is  
108 expected that they would be more severely exposed to nano-sized floating debris (Chua et al., 2014;  
109 Cole et al., 2013; Lee et al., 2013; Murray and Cowie, 2011). Nanoscale materials (1-100 nm range)  
110 may end up to a significant aggregation in sea water due to counterbalance of several parameters as  
111 pH, salts, natural organic matter (NOM) and colloids with size and surface chemistry (i.e. charges)  
112 of the nanoparticle itself. Such aggregates may sink along sea column and reach marine sediment  
113 despite but they still undergo vertical repartition with consequent buoyancy due to strong water  
114 currents (i.e. upwelling) or by ingestion by planktonic organisms (Corsi, 2014). Based on recent  
115 evidences of suspended nanoscale materials in sea water, bottom grazers and filter-feeders species  
116 are expected to be exposed to high concentrations of plastic debris in their natural environment  
117 (Wright et al., 2013b). Our recent findings on selected PS NPs showed that both accumulation and  
118 toxicity affect the early life stages of development of the Mediterranean sea urchin *Paracentrotus*  
119 *lividus* depending on NPs surface charges and their aggregation in sea water (Della Torre et al.,  
120 2014a).

121 Specie-specific sensitivities including toxicity of 55 and 110 nm polyethyleneimine-PS NPs for  
122 aquatic organisms has been recently reported by Casado et al., 2013. Likewise, 70 nm PS-COOH  
123 NPs significantly affect algal growth and reproductive success of *Daphnia magna* through diet

124 (Besseling et al., 2014). The few contributions on marine species show a significant accumulation  
125 in the gut of rotifers (Snell and Hicks, 2011) and bivalves (Ward and Kach, 2009; Wegner et al.,  
126 2012) as well as in sea urchin embryos, which were the first evidence of toxicity reported for the  
127 PS-NH<sub>2</sub> (Della Torre et al., 2014a). Therefore, beside the potential in nanotechnology, PS NPs  
128 represent an important source of primary nanoplastics entering in marine environment and based on  
129 the observed toxicity more studies should investigate their impact on organisms belonging to  
130 different trophic levels (Handy et al., 2008; Klaine, 2008; Moore, 2006). Although commonly  
131 applied as model positively charged NPs in nanosafety studies, it is important to stress that PS-NH<sub>2</sub>  
132 are a special surface functionalised variant of standard PS and the presence of similar positively  
133 charged NPs in plastic degradation products has not been fully determined yet.

134 Within marine model species, microcrustaceans are highly recommended in ecotoxicological  
135 studies being numerous and planktonic so expected to be exposed to nano-sized floating debris  
136 including the low-density PS NPs. Any negative impact on such key trophic level might disrupt the  
137 energy flow in marine ecosystems.

138 Brine shrimp *Artemia franciscana*, a filter-feeding anostracan microcrustacean is typical of inland  
139 salt water bodies but also temperate coastal areas and largely used in ecotoxicology studies for  
140 acute toxicity testing as model marine zooplankton species (EPA, 2002; Nunes et al., 2006;  
141 Persoone and Wells, 1987).

142 Up to date, the few recent studies conducted using this species show some variability in the  
143 observed effects towards NPs exposure (e.g. CeO<sub>2</sub>, carbon black, graphene, Ag and other metal  
144 oxides) likely due to differences in the NPs chemical nature, size, surface properties as well as  
145 aggregation in the exposure water media (Arulvasu et al., 2014; Ates et al., 2013a; Ates et al.,  
146 2013b; Auffan et al., 2013; Gambardella et al., 2014; Pretti et al., 2014; Rodd et al., 2014). Brine  
147 shrimps are non-selective filter feeders that can efficiently graze over a wide range of particles  
148 sizes, thus likely including synthetic nanomaterials and nanoplastics (Makridis and Vadstein, 1999).  
149 The aim of the present study is to evaluate the uptake and distribution of 40 nm anionic  
150 carboxylated (PS-COOH) and 50 nm cationic amino modified (PS-NH<sub>2</sub>) PS NPs on brine shrimp *A.*  
151 *franciscana* using larvae mortality test (Nunes et al., 2006).

152

## 153 2. Materials and methods

### 154 2.1 Physico-chemical characterization of PS-COOH and PS-NH<sub>2</sub> NPs

155 40 nm green carboxylated polystyrene nanoparticles (PS-COOH) (505 nm excitation, 515 nm  
156 emission) were purchased from Invitrogen. Unlabeled and blue fluorescently labeled (358 nm  
157 excitation, 410 nm emission) 50 nm amino modified polystyrene (PS-NH<sub>2</sub>) NPs were purchased  
158 from Bangs Laboratories and Sigma, respectively. Fluorescently labeled PS beads have been  
159 recommended as priority test material to be developed and used for ecotoxicological studies (Stone  
160 et al., 2010) and investigated within the FP7 Research Infrastructure QualityNano  
161 ([www.qualitynano.eu](http://www.qualitynano.eu)) (Wang et al., 2013a; Wang et al., 2013b). Fluorescent PS micro and nano-  
162 beads have been also widely used to investigate the impact of micro and nanoplastics on marine  
163 biota (Cole et al., 2013; Lee et al., 2013; Della Torre et al., 2014a). Nevertheless their utilization has  
164 not always been combined with in-depth secondary characterization in the natural media and the  
165 role of their functionalized groups and thus surface charge have been rarely taken into account. This  
166 could lead to an incomplete comprehension of toxicity results. Although negative surface charged  
167 nanoplastics as PS-COOH have been suggested as the most widespread in the environment  
168 (Besseling et al., 2014), no data are currently on their fate and toxicity to marine biota.

169 In this study, PS-NH<sub>2</sub> were also considered as positively charged nanoplastics, although they are a  
170 special surface functionalised variant of common PS and the presence of similar positively charged  
171 NPs as plastic degradation products has not been determined.

172 TEM was applied for primary particle diameter identification of PS NPs (Philips Morgagni 268D  
173 electronics, at 80 KV and equipped with a MegaView II CCD camera). Dynamic Light Scattering  
174 (Malvern instruments) was used for size (Z-average and polydispersity index, PDI) and zeta ( $\zeta$ -)  
175 potential (mV) (Zetasizer Nano Series software, version 7.02, Particular Sciences, UK).  
176 Measurements have been performed in triplicate, each containing 11 runs of 10 seconds for  
177 determining Z-average, 20 runs for the  $\zeta$ -potential.

178 Natural sea water (NSW) was collected from a pristine area in the Tuscan archipelago and used for  
179 PS NPs suspension preparation and without PS NPs as a control. Physico-chemical parameters  
180 including some aquatic contaminants of NSW samples used in the study have been reported in  
181 Table 3. PS NPs suspensions were prepared in 0.45  $\mu$ m filtered NSW (T 25 $\pm$ 1 $^{\circ}$ C, salinity 38‰, pH  
182 8.3, conductivity 6 S/m) and quickly vortexed prior to use but not sonicated.

183 PS NPs concentrations for toxicity tests were chosen based on those used in previous studies on *in*  
184 *vitro* cell models (Bexiga et al., 2011; Salvati et al., 2011; Wang et al., 2013a). Despite data on  
185 environmental concentrations of similar particles in sea water are not available, these concentrations  
186 may be far above real exposure conditions. PS-NH<sub>2</sub> caused apoptosis in 1321N1 human cells at 50

187  $\mu\text{g/ml}$  (Bexiga et al., 2014), but in our previous study (Della Torre et al., 2014a) we observed  
188 induction of an apoptotic pathway in sea urchin developing embryos already at 3  $\mu\text{g/ml}$ , thus  
189 raising concern regarding the impact of lower PS NPs concentrations in marine organisms.

190

## 191 **2.2 Ecotoxicity tests**

192 Acute toxicity test using *A. franciscana* larvae has been developed as standard methods (CNR,  
193 2003; EPA, 2002; Vanhaecke and Persoone, 1981) for assessing the lethality of contaminants at the  
194 first stages of development (up to Instar III nauplius), since 48 h old specimens is considered the  
195 most sensitive larval end-point (Barahona and Sánchez-Fortún, 1996). Recovery experiments were  
196 also performed by transfer PS NPs exposed brine shrimp larvae after 48h in clean NSW and left  
197 there for 24h. For both acute toxicity tests and recovery experiments, certified dehydrated cysts of  
198 brine shrimp *A. franciscana* were purchased from the company MicroBioTests (Ghent, Belgium).  
199 Hatching of the cysts was obtained following the procedure described by Garaventa et al., 2010, by  
200 incubating 100 mg of cysts in glass Petri dishes containing NSW, for 24 h at  $25\pm 1^\circ\text{C}$  under light  
201 source (3000-4000 lux)

202 Newly hatched brine shrimp larvae (Instar I nauplius stage) were separated from unhatched cysts  
203 and transferred based on phototaxis into new glass Petri dishes with NSW.

204

### 205 **2.2.1 Acute toxicity test**

206 Acute toxicity tests (see Table 2) were performed according to standard APAT IRSA CNR 8060  
207 method (CNR, 2003), by adding 10 nauplii to each well of 24-well plates, containing 2 ml with  
208 suspensions of different concentrations of the PS NPs tested (0, 5, 10, 25, 50, 100  $\mu\text{g/ml}$ ) in NSW.  
209 Control was settled in NSW without PS NPs. The plates were kept at  $25\pm 1^\circ\text{C}$  for 48h in dark  
210 conditions, without providing food according to Garaventa et al., (2010). Potassium dichromate was  
211 tested as reference toxicant.

212 At 24 and 48h, the number of dead nauplii (which were motionless for 10 seconds) was counted  
213 under stereomicroscope, in order to calculate the mortality. The validity of the test was guaranteed  
214 by the control group showing <10% of mortality at 48 h. Moreover, at 48 h nauplii were also  
215 observed by optical fluorescence microscopy in order to identify any potential sub-lethal effects  
216 (see Table 2) (i.e. molting, PS NPs accumulation in the digestive tract or adhesion to the external  
217 appendages). A tentative method for calculating the amount of molts released by developing larvae  
218 was developed as follow: media from experimental groups (control-NSW and larvae exposed to PS-  
219 NPs-NSW) were collected, filtered through a 70  $\mu\text{m}$  Falcon Cell Strainer Nylon, which retained all  
220 larvae and then rinsed several times with NSW. All media were then centrifuged at 12000 rpm for 5

221 minutes and the obtained pellet weight and compared with control exposed media. This method will  
222 allow the separation of larvae from molts which were quantified by gravimetry. A number of 400  
223 larvae were considered for the quantification of the amount of molts.

224 All the experiments have been performed in triplicates and repeated at least three times. In order to  
225 determine the presence of PS NPs in the digestive tract of nauplii brine shrimp larvae were observed  
226 under optical fluorescent microscope Olympus BX51 (filter FITC 470/525 for PS-COOH; filter  
227 DAPI 365/445 for labelled PS-NH<sub>2</sub>). Images were taken with DP50 camera at 10X using Olympus  
228 DP-software.

229

### 230 **2.2.2 Recovery experiment**

231 A recovery experiment was performed following the procedure described by Ates et al., (2013a).  
232 Brine shrimp larvae (Instar II-III nauplius stage) after 48h of exposure were collected by a Pasteur  
233 pipette, rinsed using a 100 µm Falcon cell strainer and transferred to 6-well plate, containing 6 ml  
234 of NSW without PS NPs. In these conditions, nauplii were allowed to depurate at 25±1°C for 24 h  
235 in the dark. No food was provided during the recovery test. After 24h of recovery, brine shrimp  
236 larvae were examined under optical fluorescent microscope to determine gut clearance and removal  
237 of any PS NPs on the external surface and appendages of the earlier exposed larvae, compared to  
238 the control group. Recovery experiments have been performed in triplicates and repeated at least  
239 three times.

240

### 241 **2.3 Data analysis**

242 All statistical analysis were performed using Graphpad Prism5. Analysis of variance (ANOVA) was  
243 performed to compare the various treatments, and  $p < 0.05$  was taken as significant cut-off. Results  
244 of acute and recovery tests are mean of at least three independent experiments. LC<sub>50</sub> values were  
245 calculated by fitting the percentage of alive larvae to a classical sigmoidal dose-response model  
246 according to the equation:  $y = b + (a - b) / 1 + 10^{(\text{Log EC}_{50} - x)}$  where  $y$  is response,  $b$  response  
247 minimum,  $a$  response maximum,  $x$  the logarithm of effect concentration and LC<sub>50</sub> the concentration  
248 of effect giving 50% of maximum effect. Each experiment has been performed 3 to 5 times.

249

### 250 3. Results and discussion

251

#### 252 3.1 PS NPs and behavior in exposure media (natural sea water)

253 Primary particles nominal size of 40 nm PS-COOH and 50 nm PS-NH<sub>2</sub> was confirmed by TEM  
254 imaging (Fig. S1).

255 DLS showed the formation of large PS-COOH aggregates in NSW at 25±1°C with a Z-average  
256 larger than 0.9 µm, while PS-NH<sub>2</sub> resulted far less aggregated with a Z-average of 106 nm (± 2 nm  
257 s.d.) and a PDI of 0.24 (Table 1). ζ-potential measurements confirmed their anionic (-9 mV) and  
258 cationic (+18 mV) surface charges (though the presence of aggregation in the PS-COOH NPs can  
259 affect these values). The low absolute values also indicated low stability in the NSW media. Both  
260 PS NPs showed an increasing aggregation in NSW with time (0 until 48 h): for PS-NH<sub>2</sub>, PDI from  
261 0.24 to 0.4 while for PS-COOH remains > 0.3 (Table 1). The observed low stability might be  
262 related to the high ionic strength of the NSW, which can screen the particle surface charges leading  
263 to the observed aggregation, unless the particles are stabilized by other factors, such as for instance  
264 adsorption of biomolecules on their surface (Corsi, 2014). A confirmation is given by the low  
265 absolute values of ζ-potential measured for both PS NPs in NSW which suggests a screening effect  
266 of surface charges due to the higher salt content but also by proteins or other compounds in the  
267 surroundings as for instance the natural organic matter (NOM) present in NSW (Table 3) (Wang et  
268 al., 2013b). NOM as well as natural mineral remain suspended in seawater as biogenic and geogenic  
269 colloids being able to interact with NPs. The so-called heteroaggregation phenomenon is driven by  
270 the affinity between the high surface energy (e.g. charges) of the NPs and these naturally occurring  
271 colloids in NSW (Corsi, 2014).

272 Our findings underline the need to deeply characterize stability of NPs in complex environmental  
273 media as NSW, which therefore is recommended to be used in standardized ecotoxicity tests. A  
274 combination of parameters such as pH, ionic strength, salt concentrations and the presence of other  
275 biomolecules, similarly to what observed for proteins forming a corona on the NP in human blood,  
276 might strongly affect the behavior of surface charged NPs as PS in the media and more important  
277 their interactions within cells (Wang et al., 2013b).

278 Based on our findings, nanoscale aggregates of PS-NH<sub>2</sub> (~ 100 nm) are still present in NSW media  
279 while PS-COOH NPs originated microscale aggregates (> 900 nm) (Table 1).

280

281

282 **Table 1.** Physico-chemical characterization of PS NPs in Milli-Q water (mQW) and natural sea water (NSW) (0.45  $\mu\text{m}$   
 283 filtered,  $T = 25 \pm 1^\circ\text{C}$ , salinity 38 ‰, pH 8.3, conductivity 6 S/m) using DLS analysis showing Z-average (nm),  
 284 polydispersity index (PDI) and  $\zeta$ -potential (mV). Data are referred to PS NPs concentration of 50  $\mu\text{g}/\text{ml}$  and values  
 285 reported as average  $\pm$  standard deviation.  
 286

	40 nm PS-COOH			50 nm PS-NH <sub>2</sub>		
	Z-Average (nm)	PDI	$\zeta$ -potential (mV)	Z-Average (nm)	PDI	$\zeta$ -potential (mV)
mQW	58 $\pm$ 2	0.129 $\pm$ 0.01	- 34 $\pm$ 1	61 $\pm$ 0.2	0.131 $\pm$ 0.02	+ 24 $\pm$ 1
NSW	> 0.9 $\mu\text{m}$	0.302 $\pm$ 0.08	- 9 $\pm$ 2	106 $\pm$ 2	0.243 $\pm$ 0.01	+ 18 $\pm$ 10

287

288 **Table 2.** Summary of the experimental design and biological effects observed in brine shrimp *A. franciscana* larvae  
 289 after exposure to PS-COOH and PS-NH<sub>2</sub> NPs.

PS NPs	Test	Reference	Concentrations	End-point	Outcome
40 nm PS-COOH	Acute Toxicity Test (48 h)	APAT IRSA CNR 8060 (2003)	0, 5, 25, 50, 100 $\mu\text{g}/\text{ml}$	Mortality	None < 100 $\mu\text{g}/\text{ml}$
				Sub-Lethal Effects	Accumulation
	Recovery (24 h)	<i>Ates et al., 2013a</i>	0, 5, 25, 50 $\mu\text{g}/\text{ml}$	Gut Clearance	Not
50 nm PS-NH <sub>2</sub>	Acute Toxicity Test (48 h)	APAT IRSA CNR 8060 (2003)	0, 5, 25, 50, 100 $\mu\text{g}/\text{ml}$	Mortality	None < 100 $\mu\text{g}/\text{ml}$
				Sub-Lethal Effects	Adherence Molting
	Recovery (24 h)	<i>Ates et al., 2013a</i>	0, 5, 25, 50 $\mu\text{g}/\text{ml}$	Removal from external surface	Yes

290

291

292



293 **Table 3.** Physico-chemical parameters including heavy metals (Cr, As, Cd, Hg, Pb, total polycyclic aromatic  
294 hydrocarbons, PAHs) of natural sea water samples used in the study. Data also available at: SIRA RSS. [www.sira.arpat.toscana.it](http://www.sira.arpat.toscana.it)  
295

296

<b>Parameters</b>	
<b>TOC</b>	1,3 %
<b>Total oxygen</b>	6,6 mg/L
<b>Cr</b>	< 1 µg/L
<b>As</b>	1 µg/L
<b>Cd</b>	0,09 µg/L
<b>Hg</b>	0,02 µg/L
<b>Pb</b>	< 1 µg/L
<b>Total PAH</b>	0,12 mg/Kg

297

### 298 **3.2 Brine shrimp *A. franciscana* larvae acute toxicity test**

299 In order to assess acute toxicity of anionic and cationic PS NPs (40 nm PS-COOH and 50 nm PS-  
300 NH<sub>2</sub> respectively), brine shrimp larvae were exposed to NPs suspension in NSW for 48h and  
301 observed for mortality and NPs accumulation according to the standard APAT IRSA CNR 8060  
302 method (CNR, 2003) (Figure 1a,b). No significant mortality was observed at 48 h (nauplius stage)  
303 for both PS NPs up to 100 µg/ml tested concentration but several sub-lethal effects were evident  
304 (Table 2).

305 Light microscopy images of larvae at 12h of exposure clearly showed uptake of PS-COOH  
306 aggregates at all tested concentration (5-100 µg/ml), absent in controls. A massive sequestration  
307 inside the gut lumen was evident for both PS NPs (5-100 µg/ml) at 48 h (nauplius stage), as shown  
308 in Figure 1 (a, b). A further confirmation of the nature of aggregates was given by fluorescent  
309 microscopy which revealed green fluorescence in the gut of PS-COOH exposed larvae as well as  
310 blue fluorescence of PS-NH<sub>2</sub> NPs exposed ones (Figure 1 c,d).

311 By continuous recording after 48h of exposure, some of them were lately excreted as fecal pellets  
312 but not a full release of aggregates present in the gut lumen was observed (see Figure S2 and the  
313 video on the Supplementary Information). This peculiar uptake and sequestration behavior of NPs  
314 observed in brine shrimps have been recently described for several NPs (Arulvasu et al., 2014; Ates  
315 et al., 2013a; Ates et al., 2013b; Auffan et al., 2013; Gambardella et al., 2014; Pretti et al., 2014;  
316 Rodd et al., 2014) as well as for PS micro-beads in other zooplankton species (Cole et al., 2013; Lee  
317 et al., 2013). Our recent paper on sea urchin embryos was the first contribution showing the  
318 accumulation of PS-COOH NPs in embryos during development (Della Torre et al., 2014a).  
319 Accumulation seems in general not affecting mortality but being more associated with several sub-  
320 lethal effects (i.e. behavioral, physiological and biochemical) which can affect survival of brine  
321 shrimp in prolonged exposure scenarios. As already hypothesized for microplastics, the observed  
322 accumulation of PS NPs aggregates in the digestive tract may limit food intake and significantly  
323 affect growth and development of brine shrimp larvae (Besseling et al., 2014; Cole et al., 2013). In  
324 addition, the documented high ability of PS nano-debris to adsorb hydrophobic toxic contaminants  
325 may increase their bioavailability and consequently toxicity to marine organisms (Rochman et al.,  
326 2013; Velzeboer et al., 2014). More studies regarding this phenomenon are urgently needed based  
327 on the evidence of significant gut accumulation of PS NPs in exposed marine organisms as in  
328 general of other NPs as well. Our recent studies using titanium dioxide NPs provide first insight  
329 concerning the potential interference between NPs and toxic marine pollutants with consequences at  
330 various level from increase accumulation of toxicant (e.g. coupled with dioxin) to decrease toxicity  
331 (e.g. cadmium)(Canesi et al., 2014; Della Torre, 2015; Della Torre, 2014b).

332 In more realistic scenarios as during chronic exposure in the natural environment, bioavailability  
333 and uptake of both anionic and cationic PS NPs by planktonic species could lead to their transfer  
334 along marine trophic web with significant ecological consequences being zooplankton an important  
335 food source for other marine organisms. Moreover, the excretion of nano-sized polymers as PS NPs  
336 in fecal pellets could enhance their removal from the sea surface by increasing their sinking at the  
337 sea bottom level (Cole et al., 2013).

338 Despite PS-NH<sub>2</sub> (5-100 µg/ml) also accumulated alike in brine shrimp swimming larvae at 48 h,  
339 many specimens showed empty digestive traits (Fig. 1d,e). More clearly, these cationic nanoplastics  
340 were stuck to the external surface of the swimming larvae (Figure 2c) and in particular to the  
341 sensorial appendages, as shown in Figure 3 c,d. The presence of PS-NH<sub>2</sub> on the appendages was  
342 found to noticeably hampering brine shrimp larvae natation at 48h and thus probably limiting their  
343 ability of feeding. Brine shrimp *A. franciscana* creates feeding currents while swimming in order to  
344 ingest waterborne particles (Ruppert et al., 2004). Therefore longer exposure scenarios will aim to  
345 evaluate the outcome of these sub-lethal observed effects and predict possible consequences at  
346 organism and population level by disrupting behavior, feeding and growth.

347 Cole et al., (2013) recently highlighted that, depending on the size, microplastics can be ingested (7-  
348 20 µm) or externally adhere (3.8 µm) to zooplankton. In our case, the different aggregation of the  
349 two PS NPs in NSW and not their size seems not fully explain the observed effects. PS-NH<sub>2</sub> NPs  
350 which are quite well dispersed (106 nm) in NSW still accumulate to some extent in the digestive  
351 trait of brine shrimp larvae. Therefore, surface charge seems more responsible of the observed  
352 different interaction and impact to marine organisms, as already described in our previous study  
353 with sea urchin *P. lividus* embryos (Della Torre et al., 2014a).


354 In addition, at the highest PS-NH<sub>2</sub> concentrations (50 and 100 µg/ml), several molts were found in  
355 the NSW media after 48 h of exposure (Fig. 2d) compared to controls and lower concentrations. A  
356 tentative quantitative assessment of molts released by PS-NH<sub>2</sub> exposed larvae was developed. A  
357 significant increase of around 50% of molts respect to controls was observed in PS-NH<sub>2</sub> exposed  
358 larvae. This might represent a good tentative method for quantifying the molts events caused by  
359 NPs exposure since in anostracans molts are quite transparent and were easily broken by other brine  
360 shrimps swimming through. By the way, this effect has been observed several times in all replicates  
361 (10) and in the three parallel experiments performed. An increasing number of molts seemed also to  
362 be present at PS-NH<sub>2</sub> NPs higher concentrations (50 and 100 µg/ml). This is the first observation of  
363 an increase of molting events in zooplankton species exposed to NPs and further investigations  
364 should focus on mechanisms able to disrupt this hormone-controlled physiological phenomenon,  
365 which is considered the most crucial step in the life cycle of microcrustaceans as brine shrimp. The

366 importance of molting in the biodistribution and release of NPs has been described by Auffan et al.,  
367 (2013) in *Daphnia pulex* exposed to CeO<sub>2</sub> NPs. While ingestion can be considered the major route  
368 of NPs uptake in microcrustaceans, ecdysis (molting) has been considered as the main physiological  
369 mechanism of CeO<sub>2</sub> NPs release from *Daphnia* able also to decrease the direct trophic transfer to  
370 predators.

371 Based on our findings, we hypothesize that the increase of molting events in brine shrimp larvae  
372 may represent a defense mechanism regardless the exposure to cationic PS NPs (PS-NH<sub>2</sub>). The  
373 potential link between an increase in molting and the presence of PS-NH<sub>2</sub> NPs aggregates adhering  
374 to the external surface and appendages of larvae might better explain the interaction between NPs  
375 and the exposed larvae and explain the mechanism behind the observed effects.

376 Molting is also energy consuming and according to Cole et al., (2014) copepods receive less energy  
377 for their metabolism and reproduction during this peculiar stage. An increase of molting events  
378 could therefore potentially affect this energy flow with serious consequences on brine shrimp larvae  
379 growth. Long-term study including also low levels of PS-NH<sub>2</sub> exposure will help to define how  
380 cationic nanoplastics may affect brine shrimp *A. franciscana* physiology and consequently survival  
381 and reproduction.

382




383

384

385 **Figure 1.** Effects of 40 nm PS-COOH and 50 nm PS-NH<sub>2</sub> on brine shrimp *A. franciscana* nauplii at 48 h: (a) control in  
386 NSW; (b, c) accumulation of green fluorescent PS-COOH (25 µg/ml) and (e, f) blue fluorescent PS-NH<sub>2</sub> (25 µg/ml)  
387 inside the digestive tract; (d) detail of fecal pellet containing PS-COOH aggregates. Images are representative of three  
388 independent experiments. Scale bar: 100 µm.


389



390

391 **Figure 2.** Effects of 50 nm PS-NH<sub>2</sub> on brine shrimp *A. franciscana* nauplii at 48 h. Upper images showing control in  
 392 NSW: detail of clear (a) sensorial antennules, (b) antennae and (c) abdomen. Lower images showing PS-NH<sub>2</sub> (50  
 393 µg/ml) exposed nauplii (48 h): detail of nanoplastics attached to (d) sensorial antennules, (e) sensory hairs of the  
 394 antennae and (f) abdomen region. Aggregates of PS-NH<sub>2</sub> are indicated by black arrows. Images are representative of  
 395 three independent experiments. Scale bar: 100 µm.

396



397

398 **Figure 3.** Two examples of molts found in wells containing brine shrimp *A. franciscana* nauplii after 48 h of exposure  
 399 to 50 nm PS-NH<sub>2</sub>. Anostracans such as *A. franciscana* are characterized by absence of carapace, therefore molts  
 400 appeared quite transparent and easily broken by other brine shrimps swimming through. 10X images are referred to 50  
 401 µg/ml exposed organisms and representative of three independent experiments. Scale bar: 200 µm.

402

### 403 **Recovery Experiment**

404 As reported by previous studies (Ates et al., 2013b; Cole et al., 2013; Lee et al., 2013), the ingestion  
 405 of synthetic micro-beads by zooplankton species can heavily hinder the digestive tract, thus limiting  
 406 feeding, growth and survival. In order to understand the extent of this phenomenon, a recovery  
 407 experiment was performed. After 48 h of exposure, brine shrimp larvae were left in clean NSW (no  
 408 PS NPs) for 24 h without feeding and then observed by light and fluorescent microscopy.

409 All brine shrimp nauplii (10 organisms in each experiments) earlier exposed to PS-COOH (0, 5, 25,  
 410 50 µg/ml) still presented aggregates in their gut, even at the lowest NPs concentrations, in

411 agreement with the retention of microplastics up to 7 days observed by (Cole et al., 2013) in marine  
412 copepods. Moreover, the presence of food during the recovery experiments has been shown to  
413 improve the elimination efficiency of NPs from the digestive tract of brine shrimp *A. franciscana*,  
414 even if a significant proportion was retained in the gut (Ates 2013a, Ates 2013b). On the  
415 opposite, larvae exposed to PS-NH<sub>2</sub> (0, 5, 25, 50 µg/ml) and transferred to clean NSW did not show  
416 neither aggregates in the gut nor NPs attached to the external surface and appendages. However,  
417 further studies are required to exclude any potential negative impact on brine shrimp larvae due to  
418 nano-PS exposure, since long-term exposures could provide in-depth information upon the effects  
419 of nanoplastics on brine shrimp *A. franciscana*.

420

421

#### 422 **4. Conclusion**

423 Our study suggests that PS NPs might pose a risk to marine zooplankton as a result of exposure to  
424 nanoplastics at the concentrations tested here. Nano-sized PS might be able to impair food uptake  
425 (feeding), behavior (motility) and physiology (multiple molting) of brine shrimp larvae *A.*  
426 *franciscana* with consequences not only at organism and population level but on the overall  
427 ecosystem based on the key role of zooplankton on marine food webs. In addition, our study again  
428 underline that careful assessment of NP properties and stability in NSW is needed in order to  
429 properly address their behavior towards marine organisms. Aggregation but more important surface  
430 charges (cationic vs anionic) may lead to different uptake and biodistribution and perhaps disrupt  
431 important physiological function linked to feeding and growth as observed in our study. The  
432 European Marine Strategy Framework Directive has a key goal of ensuring that marine litter is kept  
433 to a level that causes “no significant harm to the marine environment” and hence there are strong  
434 links with the development of sound environmental policy to meet this need at global level  
435 including to assess the impact of nano-debris as nanoplastics in the marine environment.

436

#### 437 **Acknowledgements**

438 This study was performed in the framework of the PhD project entitled "Polystyrene nanoparticles  
439 and their impact on marine ecosystems: accumulation, disposal and toxicity in marine species from  
440 Antarctic and Mediterranean Seas". PhD student Elisa Bergami, PhD School in Geological,  
441 Environmental and Polar sciences and technologies, Department of Physical, Earth and  
442 Environmental Sciences, University of Siena (Italy). Funded by the University of Siena and the  
443 Italian National Antarctic Museum "Felice Ippolito".

444 PS NPs characterization was performed at the Centre for BioNano Interactions (University College  
445 of Dublin) funded by the Erasmus Long Life Learning Program (year 2012-2013) and based  
446 onworks supported by the EU FP7 Capacities project QualityNano (grant no. INFRA-2010-  
447 262163). We thank Claudia Faleri (Dept Life Sciences, University of Siena) for her assistance in  
448 TEM images.  
449

450 **References**

451

- 452 Andrady, A. L., 2003. *Plastics and the Environment*. Wiley.
- 453 Arulvasu, C., Jennifer, S. M., Prabhu, D., Chandhirasekar, D., 2014. Toxicity effect of silver nanoparticles in  
454 brine shrimp *Artemia*. *ScientificWorldJournal*. 2014, 256919.
- 455 Ates, M., Daniels, J., Arslan, Z., Farah, I. O., 2013a. Effects of aqueous suspensions of titanium dioxide  
456 nanoparticles on *Artemia salina*: assessment of nanoparticle aggregation, accumulation, and toxicity.  
457 *Environ Monit Assess*. 185, 3339-48.
- 458 Ates, M., Daniels, J., Arslan, Z., Farah, I. O., Rivera, H. F., 2013b. Comparative evaluation of impact of Zn  
459 and ZnO nanoparticles on brine shrimp (*Artemia salina*) larvae: effects of particle size and solubility  
460 on toxicity. *Environ Sci Process Impacts*. 15, 225-33.
- 461 Auffan, M., Bertin, D., Chaurand, P., Pailles, C., Dominici, C., Rose, J., Bottero, J. Y., Thiery, A., 2013.  
462 Role of molting on the biodistribution of CeO<sub>2</sub> nanoparticles within *Daphnia pulex*. *Water Res*. 47,  
463 3921-30.
- 464 Bandyopadhyay, A., Basak, G. C., 2007. Studies on photocatalytic degradation of polystyrene. *Material*  
465 *Science and Technology*. 23, 307-317.
- 466 Barahona, M. V., Sánchez-Fortún, S., 1996. Comparative Sensitivity of Three Age Classes of *Artemia salina*  
467 Larvae to Several Phenolic Compounds. *Bull. Environ. Contam. Toxicol*. 56, 271-278.
- 468 Besseling, E., Wang, B., Lurling, M., Koelmans, A. A., 2014. Nanoplastic affects growth of *S. obliquus* and  
469 reproduction of *D. magna*. *Environ Sci Technol*. 48, 12336-43.
- 470 Bexiga, M. G., Kelly, C., Dawson, K. A., Simpson, J. C., 2014. RNAi-mediated inhibition of apoptosis fails  
471 to prevent cationic nanoparticle-induced cell death in cultured cells. *Nanomedicine (Lond)*. 9, 1651-  
472 64.
- 473 Bexiga, M. G., Varela, J. A., Wang, F., Fenaroli, F., Salvati, A., Lynch, I., Simpson, J. C., Dawson, K. A.,  
474 2011. Cationic nanoparticles induce caspase 3-, 7- and 9-mediated cytotoxicity in a human  
475 astrocytoma cell line. *Nanotoxicology*. 5, 557-67.
- 476 Bramini, M., Ye, D., Hallerbach, A., Raghnaill, M. N., Salvati, A., A<sup>o</sup>berg, C., Dawson, K. A., 2014.  
477 Imaging Approach to Mechanistic Study of Nanoparticle Interactions with the Blood-Brain Barrier.  
478 *ACS Nano*. 8, 4304-4312.
- 479 Browne, M. A., Crump, P., Niven, S. J., Teuten, E., Tonkin, A., Galloway, T., Thompson, R., 2011.  
480 Accumulation of microplastic on shorelines worldwide: sources and sinks. *Environ Sci Technol*. 45,  
481 9175-9.
- 482 Canesi, L., Frenzilli, G., Balbi, T., Bernardeschi, M., Ciacci, C., Corsolini, S., Della Torre, C., Fabbri, R.,  
483 Faleri, C., Focardi, S., Guidi, P., Kocan, A., Marcomini, A., Mariottini, M., Nigro, M., Pozo-  
484 Gallardo, K., Rocco, L., Scarcelli, V., Smerilli, A., Corsi, I., 2014. Interactive effects of n-TiO<sub>2</sub> and  
485 2,3,7,8-TCDD on the marine bivalve *Mytilus galloprovincialis*. *Aquat Toxicol*. 153, 53-65.
- 486 Casado, M. P., Macken, A., Byrne, H. J., 2013. Ecotoxicological assessment of silica and polystyrene  
487 nanoparticles assessed by a multitrophic test battery. *Environ Int*. 51, 97-105.
- 488 Chua, E. M., Shimeta, J., Nugegoda, D., Morrison, P. D., Clarke, B. O., 2014. Assimilation of  
489 polybrominated diphenyl ethers from microplastics by the marine amphipod, *Allorchestes*  
490 *compressa*. *Environ Sci Technol*. 48, 8127-34.
- 491 CNR, A. I., Metodo 8060 di valutazione della tossicità acuta con *Artemia* sp., Manuali e Linee Guida -  
492 Metodi analitici per le acque, Vol. 3. APAT IRSA CNR, 2003, pp. 1043-1049.
- 493 Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., Galloway, T. S., 2013.  
494 Microplastic ingestion by zooplankton. *Environ Sci Technol*. 47, 6646-55.
- 495 Cole, M., Webb, H., Lindeque, P. K., Fileman, E. S., Halsband, C., Galloway, T. S., 2014. Isolation of  
496 microplastics in biota-rich seawater samples and marine organisms. *Sci Rep*. 4, 4528.
- 497 Corsi, I., Cherr, G. N., Lenihan, H. S., Labille, J., Hasselov, M., Canesi, L., Dondero, F., Frenzilli, G.,  
498 Hristozov, D., Punes, V., Della Torre, C., Pinsino, A., Libralato, G., Marcomini, A., Sabbioni, E.,  
499 Matranga, V., 2014. Common Strategies and Technologies for the Ecosafety Assessment and Design  
500 of Nanomaterials Entering the Marine Environment. *Nano Focus*. 8, 9694-9709.
- 501 Cozar, A., Echevarria, F., Gonzalez-Gordillo, J. I., Irigoien, X., Ubeda, B., Hernandez-Leon, S., Palma, A.  
502 T., Navarro, S., Garcia-de-Lomas, J., Ruiz, A., Fernandez-de-Puelles, M. L., Duarte, C. M., 2014.  
503 Plastic debris in the open ocean. *Proc Natl Acad Sci U S A*. 111, 10239-44.



- 504 Cozar, A., Sanz-Martin, M., Marti, E., Gonzalez-Gordillo, J. I., Ubeda, B., Galvez, J. A., Irigoien, X.,  
505 Duarte, C. M., 2015. Plastic accumulation in the mediterranean sea. *PLoS One*. 10, e0121762.
- 506 Della Torre, C., Balbi, T., Grassi, G., Frenzilli, G., Bernardeschi, M., Smerilli, A., Guidi, P., Canesi, L.,  
507 Nigro, M., Monaci, F., Scarcelli, V., Rocco, L., Focardi, S. E., Monopoli, M., Corsi, I., 2015.  
508 Titanium dioxide nanoparticles modulate the toxicological response to cadmium in the gills of  
509 *Mytilus galloprovincialis*. *J Hazard Mater*. 297, 92-100.
- 510 Della Torre, C., Bergami, E., Salvati, A., Faleri, C., Cirino, P., Dawson, K. A., Corsi, I., 2014a.  
511 Accumulation and embryotoxicity of polystyrene nanoparticles at early stage of development of sea  
512 urchin embryos *Paracentrotus lividus*. *Environ Sci Technol*. 48, 12302-11.
- 513 Della Torre, C., Bocci, E., Focardi, S. E., Corsi, I., 2014b. Differential ABCB and ABCC gene expression  
514 and efflux activities in gills and hemocytes of *Mytilus galloprovincialis* and their involvement in  
515 cadmium response. *Mar Environ Res*. 93, 56-63.
- 516 EPA, U., Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and  
517 marine organisms. 2002.
- 518 Eriksen, M., Lebreton, L. C., Carson, H. S., Thiel, M., Moore, C. J., Borerro, J. C., Galgani, F., Ryan, P. G.,  
519 Reisser, J., 2014. Plastic Pollution in the World's Oceans: More than 5 Trillion Plastic Pieces  
520 Weighing over 250,000 Tons Afloat at Sea. *PLoS One*. 9, e111913.
- 521 Farrell, P., Nelson, K., 2013. Trophic level transfer of microplastic: *Mytilus edulis* (L.) to *Carcinus maenas*  
522 (L.). *Environ Pollut*. 177, 1-3.
- 523 Galgani, F., Fleet, D., Van Franeker, J., Katsanevakis, S., Maes, T., Mouat, J., Oosterbaan, L., Poitou, I.,  
524 Hanke, G., Thompson, R., Amato, E., Birkun, A., Janssen, C., Marine Strategy Framework  
525 Directive Task Group 10 Report Marine Litter. In: N. Zampoukas, (Ed.), 2010, pp. 57.
- 526 Gambardella, C., Mesaric, T., Milivojevic, T., Sepcic, K., Gallus, L., Carbone, S., Ferrando, S., Faimali, M.,  
527 2014. Effects of selected metal oxide nanoparticles on *Artemia salina* larvae: evaluation of mortality  
528 and behavioural and biochemical responses. *Environ Monit Assess*. 186, 4249-59.
- 529 Garaventa, F., Gambardella, C., Di Fino, A., Pittore, M., Faimali, M., 2010. Swimming speed alteration of  
530 *Artemia* sp. and *Brachionus plicatilis* as a sub-lethal behavioural end-point for ecotoxicological  
531 surveys. *Ecotoxicology*. 19, 512-9.
- 532 Handy, R. D., Owen, R., Valsami-Jones, E., 2008. The ecotoxicology of nanoparticles and nanomaterials:  
533 current status, knowledge gaps, challenges, and future needs. *Ecotoxicology*. 17, 315-25.
- 534 Hofer, T. N., 2008. *Marine Pollution: New Research*. Nova Science Publishers.
- 535 Innocenti, F. D., Biodegradability and compostability. In: E. Chiellini, Solaro, R., (Ed.), *Biodegradable*  
536 *Polymers and Plastics*. Kluwer Academic/Plenum Publishers, New York, 2003, pp. 33-46.
- 537 Klaine, S. J., Alvarez, P. J., Batley, G. E., Fernandes, T. F., Handy, R. D., Lyon, D. Y., Mahendra, S.,  
538 McLaughlin, M. J., Lead, J. R., 2008. Nanomaterials in the environment: behavior, fate,  
539 bioavailability, and effects. *Environ Toxicol Chem*. 27, 1825-1851.
- 540 Lee, K. W., Shim, W. J., Kwon, O. Y., Kang, J. H., 2013. Size-dependent effects of micro polystyrene  
541 particles in the marine copepod *Tigriopus japonicus*. *Environ Sci Technol*. 47, 11278-83.
- 542 Lesniak, A., Campbell, A., Monopoli, M. P., Lynch, I., Salvati, A., Dawson, K. A., 2010. Serum heat  
543 inactivation affects protein corona composition and nanoparticle uptake. *Biomaterials*. 31, 9511-8.
- 544 Lithner, D., Larsson, A., Dave, G., 2011. Environmental and health hazard ranking and assessment of plastic  
545 polymers based on chemical composition. *Sci Total Environ*. 409, 3309-24.
- 546 Liu, Y., Li, W., Lao, F., Liu, Y., Wang, L., Bai, R., Zhao, Y., Chen, C., 2011. Intracellular dynamics of  
547 cationic and anionic polystyrene nanoparticles without direct interaction with mitotic spindle and  
548 chromosomes. *Biomaterials*. 32, 8291-303.
- 549 Loos, C., Syrovets, T., Musyanovych, A., Mailander, V., Landfester, K., Nienhaus, G. U., Simmet, T., 2014.  
550 Functionalized polystyrene nanoparticles as a platform for studying bio-nano interactions. *Beilstein J*  
551 *Nanotechnol*. 5, 2403-12.
- 552 Lunov, O., Syrovets, T., Loos, C., Beil, J., Delacher, M., Tron, K., Nienhaus, G.U., Musyanovych, A.,  
553 Mailander, V., Landfester, K., Simmet, T., 2011. Differential Uptake of Functionalized Polystyrene  
554 Nanoparticles by Human Macrophages and a Monocytic Cell Line. *ACS Nano*. 5, 1657-1669.
- 555 Makridis, P., Vadstein, O., 1999. Food size selectivity of *Artemia franciscana* at three developmental stages.  
556 *Journal of Plankton Research*. 21, 2191-2201.
- 557 Manzo, S., Miglietta, M. L., Rametta, G., Buono, S., Di Francia, G., 2013. Embryotoxicity and  
558 spermotoxicity of nano-sized ZnO for Mediterranean sea urchin *Paracentrotus lividus*. *J Hazard*  
559 *Mater*. 254-255, 1-9.

560 Matranga, V., Corsi, I., 2012. Toxic effects of engineered nanoparticles in the marine environment: model  
561 organisms and molecular approaches. *Mar Environ Res.* 76, 32-40.

562 Moore, M. N., 2006. Do nanoparticles present ecotoxicological risks for the health of the aquatic  
563 environment? *Environ Int.* 32, 967-76.

564 Murray, F., Cowie, P. R., 2011. Plastic contamination in the decapod crustacean *Nephrops norvegicus*  
565 (Linnaeus, 1758). *Mar Pollut Bull.* 62, 1207-17.

566 Nowack, B., Bucheli, T. D., 2007. Occurrence, behavior and effects of nanoparticles in the environment.  
567 *Environ Pollut.* 150, 5-22.

568 Nunes, B. S., Carvalho, F. D., Guilhermino, L. M., Van Stappen, G., 2006. Use of the genus *Artemia* in  
569 ecotoxicity testing. *Environ Pollut.* 144, 453-62.

570 Persoone, G., Wells, P. G., *Artemia* in aquatic toxicology: a review. In: P. Sorgeloos, Bengtson, D. A.,  
571 Declair, W., Jaspers, E., (Ed.), *Artemia* research and its applications. Universa Press, Wetteren,  
572 Belgium, 1987, pp. 380.

573 Pretti, C., Oliva, M., Pietro, R. D., Monni, G., Cevasco, G., Chiellini, F., Pomelli, C., Chiappe, C., 2014.  
574 Ecotoxicity of pristine graphene to marine organisms. *Ecotoxicol Environ Saf.* 101, 138-45.

575 Rochman, C. M., Manzano, C., Hentschel, B. T., Simonich, S. L., Hoh, E., 2013. Polystyrene plastic: a  
576 source and sink for polycyclic aromatic hydrocarbons in the marine environment. *Environ Sci*  
577 *Technol.* 47, 13976-84.

578 Rodd, A. L., Creighton, M. A., Vaslet, C. A., Rangel-Mendez, J. R., Hurt, R. H., Kane, A. B., 2014. Effects  
579 of surface-engineered nanoparticle-based dispersants for marine oil spills on the model organism  
580 *Artemia franciscana*. *Environ Sci Technol.* 48, 6419-27.

581 Rossi, G., Barnoud, J., Monticelli, L., 2014. Polystyrene Nanoparticles Perturb Lipid Membranes. *The*  
582 *Journal of Physical Chemistry Letters.* 5, 241-246.

583 Ruppert, E. E., Fox, R. S., Barnes, R. D., 2004. *Invertebrate Zoology: A Functional Evolutionary Approach.*  
584 Thomson-Brooks/Cole.

585 Salvati, A., Åberg, C., dos Santos, T., Varela, J., Pinto, P., Lynch, I., Dawson, K. A., 2011. Experimental and  
586 theoretical comparison of intracellular import of polymeric nanoparticles and small molecules:  
587 toward models of uptake kinetics. *Nanomedicine.* 7, 818-26.

588 Setälä, O., Fleming-Lehtinen, V., Lehtiniemi, M., 2014. Ingestion and transfer of microplastics in the  
589 planktonic food web. *Environ Pollut.* 185, 77-83.

590 Silvestre, C., Duraccio, D., Cimmino, S., 2011. Food packaging based on polymer nanomaterials. *Progress in*  
591 *Polymer Science.* 36, 1766-1782.

592 SIRA RSS. [www.sira.arp.at.toscana.it](http://www.sira.arp.at.toscana.it)

593 Snell, T. W., Hicks, D. G., 2011. Assessing toxicity of nanoparticles using *Brachionus manjavacas*  
594 (Rotifera). *Environ Toxicol.* 26, 146-52.

595 Stone, V., Nowack, B., Baun, A., van den Brink, N., Kammer, F., Dusinska, M., Handy, R., Hankin, S.,  
596 Hasselov, M., Joner, E., Fernandes, T. F., 2010. Nanomaterials for environmental studies:  
597 classification, reference material issues, and strategies for physico-chemical characterisation. *Sci*  
598 *Total Environ.* 408, 1745-54.

599 UNEP, Marine Debris as a Global Environmental Problem. UNEP Scientific and Technical Advisory Panel,  
600 2011, pp. 40.

601 Vanhaecke, P., Persoone, G., Report on an Intercalibration exercise on a Short-term Standard Toxicity Test  
602 with *Artemia nauplii* (ARC-TEST). Vol. 106. Artemia Reference Center, Laboratory for Biological  
603 Research in Aquatic Pollution and Laboratory for Marine Culture, Ghent, Belgium, 1981, pp. 359-376.

604 Velzeboer, I., Kwadijk, C. J., Koelmans, A. A., 2014. Strong sorption of PCBs to nanoplastics, microplastics,  
605 carbon nanotubes, and fullerenes. *Environ Sci Technol.* 48, 4869-76.

606 Wang, F., Bexiga, M. G., Anguissola, S., Boya, P., Simpson, J. C., Salvati, A., Dawson, K. A., 2013a. Time  
607 resolved study of cell death mechanisms induced by amine-modified polystyrene nanoparticles.  
608 *Nanoscale.* 5, 10868-76.

609 Wang, F., Yu, L., Monopoli, M. P., Sandin, P., Mahon, E., Salvati, A., Dawson, K. A., 2013b. The  
610 biomolecular corona is retained during nanoparticle uptake and protects the cells from the damage  
611 induced by cationic nanoparticles until degraded in the lysosomes. *Nanomedicine.* 9, 1159-68.

612 Ward, J. E., Kach, D. J., 2009. Marine aggregates facilitate ingestion of nanoparticles by suspension-feeding  
613 bivalves. *Mar Environ Res.* 68, 137-42.

- 614 Watts, A. J., Lewis, C., Goodhead, R. M., Beckett, S. J., Moger, J., Tyler, C. R., Galloway, T. S., 2014.  
615 Uptake and retention of microplastics by the shore crab *Carcinus maenas*. *Environ Sci Technol.* 48,  
616 8823-30.
- 617 Wegner, A., Besseling, E., Foekema, E. M., Kamermans, P., Koelmans, A. A., 2012. Effects of  
618 nanopolystyrene on the feeding behavior of the blue mussel (*Mytilus edulis* L.). *Environ Toxicol*  
619 *Chem.* 31, 2490-7.
- 620 Wright, S. L., Rowe, D., Thompson, R. C., Galloway, T. S., 2013a. Microplastic ingestion decreases energy  
621 reserves in marine worms. *Curr Biol.* 23, R1031-3.
- 622 Wright, S. L., Thompson, R. C., Galloway, T. S., 2013b. The physical impacts of microplastics on marine  
623 organisms: a review. *Environ Pollut.* 178, 483-92.
- 624

**Nano-sized polystyrene affects feeding, behaviour and physiology of brine shrimp *Artemia franciscana* larvae**

Elisa Bergami<sup>1</sup>, Elena Bocci<sup>1</sup>, Maria Luisa Vannuccini<sup>1</sup>, Marco Monopoli<sup>2</sup>, Anna Salvati<sup>3</sup>, Kenneth A. Dawson<sup>2</sup>, Ilaria Corsi I.<sup>1\*</sup>

<sup>1</sup>Department of Physical, Earth and Environmental Sciences, University of Siena, Italy;

<sup>2</sup>Centre for BioNano Interactions, School of Chemistry and Chemical Biology, University College Dublin, Ireland;

<sup>3</sup>Department of Pharmacy, Division Pharmacokinetics, Toxicology and Targeting, University of Groningen, The Netherlands

**Supplementary material**

This document is an electronic supplement which provides additional information about this work.


It is organized as follows:

**Figure S1**

**Figure S2**

\*Corresponding Author

\* Phone (+39) 0577 232830; fax (+39) 0577 232806; e-mail: [ilaria.corsi@unisi.it](mailto:ilaria.corsi@unisi.it); mail: Department of Physical, Earth and Environmental Sciences, University of Siena, via Mattioli, 4, 53100 Siena, Italy



**Figure S1.** Primary characterization of PS NPs: TEM images showing an average size value of  $42 \pm 1.9$  nm PS-COOH (A) and  $53 \pm 2$  nm PS-NH<sub>2</sub> (B) suspended in distilled water. Scale bar: 200 nm.



**Figure S2.** Still of video data showing brine shrimp *A. salina* larvae after 48 h of exposure to PS-COOH (25  $\mu\text{g}/\text{ml}$ ), excreting a pellet containing PS-COOH aggregates. Scale bar: 100  $\mu\text{m}$ .

## Video

[Click here to download Video: VIDEO\\_ARTEMIA PSCOOH.mov](#)

Video Still  
[Click here to download high resolution image](#)

