

Multidisciplinary study of the Lower Palaeolithic site of Cimitero di Atella (Basilicata), Italy

This is a pre print version of the following article:
Original:
Rocca, R., Giannandrea, P., Pereira, A., Bahain, J, Boschin, F., Da Costa, A., et al. (2023). Multidisciplinary study of the Lower Palaeolithic site of Cimitero di Atella (Basilicata), Italy. QUATERNARY INTERNATIONAL, 676, 1-26 [10.1016/j.quaint.2023.09.004].
Availability:
This version is availablehttp://hdl.handle.net/11365/1245474 since 2023-09-22T13:59:16Z
Published:
DOI:10.1016/j.quaint.2023.09.004
Terms of use:
Open Access
The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license. For all terms of use and more information see the publisher's website.

(Article begins on next page)

Quaternary International

Multidisciplinary study of the Lower Palaeolithic site of Cimitero di Atella (Basilicata), Italy. --Manuscript Draft--

Manuscript Number:	
Article Type:	Regular Article
Corresponding Author:	Roxane Rocca Université Paris I Panthéon-Sorbonne Paris, Île-de-France FRANCE
First Author:	Roxane Rocca
Order of Authors:	Roxane Rocca
	Paolo Giannandrea
	Alison Pereira
	Jean-Jacques Bahain
	Francesco Boschin
	Amélie Da Costa
	Federico Di Rita
	François Fouriaux
	Alessio lannucci
	Lucie Germond
	Dario Gioia
	Donatella Magri
	Beniamino Mecozzi
	Sebastien Nomade
	Raffaele Sardella
	Marcello Schiattarella
	Pierre Voinchet
	Daniele Aureli
Abstract:	The Lower Palaeolithic site of Cimitero di Atella is located in the Basilicata region (Southern Italy), about 10 km south of the extinct Monte Vulture volcano. The site was discovered in the early 1990s and was continuously excavated for nearly twenty years under the supervision of Professor E. Borzatti von Löwenstern (University of Florence). This open-air site contained a five-metre-thick fluvio-lacustrine sequence characterized by the occurrence of two main archaeological units with lithic industries and faunal remains. Based on the composition of the lithic assemblages, and in particular the presence of handaxes in the Lower unit, Borzatti von Löwenstern (et al., 1997) attributed the site to the Early Acheulean. Cimitero di Atella was interpreted as the result of various lake shore occupations linked to the exploitation of large mammals (Palaeoloxodon antiquus and Bison sp.) and the opportunistic use of raw materials to produce simple small and large lithic tools (Borzatti von Löwenstern et al., 1997). The purpose of this paper is to present the last five years of research at the site. Geochronological, palaeontological and palynological investigations have been conducted to shed light on the environmental and chronological context of the site and the associated human occupations levels. This first essential step enables us to evaluate the nature and potential of archaeological data in order to place the results in a broader framework.
Suggested Reviewers:	Marta Arzarello

rzrmrt@unife.it
Biagio Giaccio biagio.giaccio@cnr.it
Jean-Luc Guadelli jean-luc.guadelli@u-bordeaux.fr
Małgorzata Kot m.kot@uw.edu.pl
Laurent Deschodt laurent.deschodt@inrap.fr

UNIVERSITÉ PARIS 1 PANTHÉON SORBONNE UFR HISTOIRE DE L'ART ET ARCHÉOLOGIE



Roxane ROCCA

Maîtresse de conférence Université Paris 1 (Panthéon-Sorbonne) Institut d'Histoire de l'Art et d'Archéologie 3, rue Michelet F-75005 Paris UMR 8068 – TEMPS (Technologie et Ethnologie des Mondes PréhistoriqueS) MSH Mondes, Bâtiment René Ginouvès, 21 Allée de l'Université, F-92023 Nanterre. <u>Roxane.rocca@univ-paris1.fr</u>

Cover letter to Quaternary International

Paris, 13/02/2023

Dear Editor,

I have the honour to submit, on behalf of all the collaborators, this article entitled "Multidisciplinary study of the Lower Palaeolithic site of Cimitero di Atella (Basilicata), Italy". This is a first synthetic work on the Lower Palaeolithic site of Cimitero di Atella in Italy. We resumed excavation on the site in 2016 and carried out new studies on the archaeological material, but also on the environmental context and the chronostratigraphic framework. Roxane ROCCA and Daniele AURELI have the excavation permit from the Italian Heritage and manage the field organisation and documentation together with Lucie GERMOND, Amélie DA COSTA and François FOURIAUX. Roxane ROCCA, Daniele AURELI, Lucie GERMOND and Amélie DA COSTA participate to the lithic study. Paolo GIANNANDREA, Dario GIOIA and Marcello SCHIATTARELLA, conducted the geological study, Alison PEREIRA and Sebastien NOMADE the Ar/Ar dating, Jean-Jacques BAHAIN and Pierre VOINCHET the ESR dating. The archaeozoological and palaeontological study was led by Francesco BOSCHIN, Alessio IANNUCCI, Beniamino MECOZZI and Raffaele SARDELLA. Donatella MAGRI and Federico DI RITA where in charge of the palynological analysis. All the co-author to the article produced the figures and text corresponding to their skill and have approved the final version of the manuscript.

Hoping that this article will be of interest to your review.

Best regards,

Roxane ROCCA

Université Paris 1 Panthéon-Sorbonne 12, place du Panthéon 75231 Paris Cedex 05 – Tél. : +33 (0)1 44 07 80 00 – www.univ-paris1.fr

UMR 8068, Technologie et Ethnologie des Mondes PréhistoriqueS MSH Mondes, Bâtiment René Ginouvès, 21 Allée de l'Université, F-92023 Nanterre Cedex

1 2	Multidisciplinary study of the Lower Palaeolithic site of <i>Cimitero di Atella</i> (Basilicata), Italy.
3	
4	
5	Roxane ROCCA ^a , Paolo GIANNANDREA ^b , Alison PEREIRA ^c , Jean-Jacques BAHAIN ^d ,
6	Francesco BOSCHIN ^e , Amélie DA COSTA ^f , Federico DI RITA ^g , François FOURIAUX ^h , Alessio
7	IANNUCCI ⁱ , Lucie GERMOND ⁱ , Dario GIOIA ^k , Donatella MAGRI ^I , Beniamino MECOZZI ⁱ ,
8	Sebastien NOMADE ^m , Raffaele SARDELLA ⁱ , Marcello SCHIATTARELLA ^b , Pierre VOINCHET ^d ,
9	Daniele AURELI ⁿ .
10	a. Université Paris 1 Panthéon-Sorbonne, école d'histoire de l'archéologie de la Sorbonne, UMR
11	8068, Technologie et Ethnologie des Mondes PréhistoriqueS (TEMPS), MSH Mondes, Bâtiment
12	René Ginouvès, 21 Allée de l'Université, 92023 Nanterre Cedex, France. roxane.rocca@univ-
13	paris1.fr
14	b. Dipartimento delle Culture Europee e del Mediterraneo, Università della Basilicata, via Lanera,
15	20 – 75100 Matera, Italy. <u>paolo.giannandrea@unibas.it</u> marcello.schiattarella@unibas.it
16	c. Université Paris-Saclay, Umr 8148 GEOPS, Campus de l'Université Paris Sud Bâtiment 504,
17	Rue du Belvédère, 91405 Orsay Cedex, France. alison.pereira@universite-paris-saclay.fr
18	d. Muséum national d'Histoire naturelle de Paris, Département « Homme et Environnement »,
19	UMR7194, Histoire Naturelle de l'Homme Préhistorique (HNHP), 1 rue René Panhard, 75013, Paris,
20	France. jean-jacques.bahain@mnhn.fr pierre.voinchet@mnhn.fr
21	e. Università Degli Studi di Siena, Dipartimento di Scienze Fisiche, Della Terra e Dell'Ambiente,
22	UR Preistoria e Antropologia, Via Laterina 8, IT 53100, Siena, Italy. <u>boschin@unisi.it</u>
23	f. Service départemental d'archéologie du Val d'Oise, UMR 7041, ArScAn, Equipe AnTET, MSH
24	Mondes, Bâtiment Max Weber, 21 Allée de l'Université, 92023 Nanterre Cedex, France.
25	amel.dacosta@gmail.com
26	g. Dipartimento di Biologia Ambientale, Sapienza Università di Roma, Piazzale Aldo Moro 5,

1

27	00185 Roma, Italy. federico.dirita@uniroma1.it donatella.magri@uniroma1.it
28	h. Centre Jean Bérard, École française de Rome, UMR 8546 AOROC, Campus Condorcet,
29	Bâtiment Recherche Nord, 14 Cours des Humanités, 93322 Aubervilliers, France.
30	francois.fouriaux@cnrs.fr
31	i. Department of Earth Sciences (PaleoFactory lab.), Sapienza University of Rome, Piazzale Aldo
32	Moro 5, 00185, Rome, Italy. alessio.iannucci@uniroma1.it beniamino.mecozzi@uniroma1.it
33	raffaele.sardella@uniroma1.it
34	j. <u>lucie.germond@hotmail.fr</u>
35	k. CNR-ISPC, Tito Scalo, 85050, Potenza, Italy. dario.gioia@cnr.it
36	m. CEA-CNRS, UMR 8212, Laboratoire des Sciences du Climat et de l'Environnement, CEA-
37	UVSQ et Université Paris Saclay, Bat 714, Orme des Merisiers, 91191 Gif sur Yvette, France. LSCE
38	sebastien.nomade@lsce.ipsl.fr
39	n. UMR 7041, ArScAn, Equipe AnTET, MSH Mondes, Bâtiment Max Weber, 21 Allée de
40	l'Université, 92023 Nanterre Cedex, France. danieleaureli1@gmail.com

41 Abstract:

42 The Lower Palaeolithic site of Cimitero di Atella is located in the Basilicata region (Southern Italy), about 10 km south of the extinct Monte Vulture volcano. The site was discovered in the early 43 44 1990s and was continuously excavated for nearly twenty years under the supervision of Professor E. Borzatti von Löwenstern (University of Florence). This open-air site contained a five-metre-thick 45 fluvio-lacustrine sequence characterized by the occurrence of two main archaeological units with 46 47 lithic industries and faunal remains. Based on the composition of the lithic assemblages, and in 48 particular the presence of handaxes in the Lower unit, Borzatti von Löwenstern (et al., 1997) 49 attributed the site to the Early Acheulean. Cimitero di Atella was interpreted as the result of various 50 lake shore occupations linked to the exploitation of large mammals (Palaeoloxodon antiquus and 51 Bison sp.) and the opportunistic use of raw materials to produce simple small and large lithic tools 52 (Borzatti von Löwenstern et al., 1997).

The purpose of this paper is to present the last five years of research at the site. Geochronological, palaeontological and palynological investigations have been conducted to shed light on the environmental and chronological context of the site and the associated human occupations levels. This first essential step enables us to evaluate the nature and potential of archaeological data in order to place the results in a broader framework.

58 Keywords:

Lower Palaeolithic, Quaternary, Monte Vulture Volcano, Geology, Geochronology, 40Ar/39Ar
geochronology, Lithic industry, Early–Middle Pleistocene Transition (EMPT)

61 1. Introduction

62 The Lower Palaeolithic site of Cimitero di Atella is located in the Basilicata region (Southern Italy), about 10 km south of the extinct Monte Vulture volcano (Figure 1). The site was discovered in 63 the early 1990s and was continuously excavated for nearly twenty years under the supervision of 64 65 Professor E. Borzatti von Löwenstern (University of Florence). This open-air site contained a fivemetre-thick fluvio-lacustrine sequence characterized by the occurrence of two main archaeological 66 67 units with lithic industries and faunal remains. Based on the lithology of the sedimentary layers and biochronological information from the faunal assemblages, the site was confidently attributed to the 68 Middle Pleistocene timescale (Borzatti von Löwenstern et al., 1990; Ciolli, 1997; Di Muro, 1999; 69 70 Zucchelli, 2002). The presence of volcanic materials (tephra, ignimbrites and reworked volcanic 71 minerals found in the fluvial deposits) typical of the early explosive activity of the Monte Vulture volcano indicates that human occupation took place between about 740 ka and 480 ka (Di Muro, 72 73 1999). Based on the composition of the lithic assemblages, and in particular the presence of handaxes in the Lower unit, Borzatti von Löwenstern (et al., 1997) attributed the site to the Early Acheulean. 74 Cimitero di Atella was interpreted as the result of various lake shore occupations linked to the 75 76 exploitation of large mammals (Palaeoloxodon antiquus and Bison sp.) and the opportunistic use of 77 raw materials to produce simple small and large lithic tools (Borzatti von Löwenstern et al., 1997). 78 In the light of increased early Acheulean evidence over the past decade (García-Medrano et al.,

79 2014; Moncel et al., 2016, 2019; Mosquera et al., 2016), as well as the still-open debate on the modes 80 and temporality of the emergence of handaxes in Europe (Moncel and Schreve, 2016; Moncel et al., 81 2018; Nicoud, 2013; Rocca et al., 2016a), Cimitero di Atella is of particular interest. The presence of 82 handaxes and large cutting-tools before 0.5 Ma is now well attested, mainly in South-western Europe. 83 This is generally explained by a second wave of settlement originating from Africa driving classical 84 Acheulean development. In many respects, this hypothesis is still tenuous as available data are still scarce for this pivotal period in Europe, between early settlement and the classical Acheulean. 85 86 Moreover, the comprehensive interpretation of settlement dynamics during this period is difficult, given the diversity of archaeological contexts (cave, palimpsest, snapshot, etc...), as well as the 87 88 plurality of lithic methodological approaches. In this background, the site of Cimitero di Atella will 89 undoubtedly help to test the various hypotheses concerning the diffusion of the Acheulean in Europe. 90 At the end of Borzatti von Löwenstern excavation at the site, many questions were still pending. 91 First of all, the chronological attribution based on volcano-stratigraphic correlation required 92 confirmation by sedimentological analysis and radiometric dating. Secondly, the question of site formation dynamics was crucial to interpret the archaeological record and assess the different 93 94 taphonomic aspects. Finally, it was indispensable to study the archaeological material with a 95 methodology adapted to topical issues in order to compare it to other contemporaneous sites.

A new multidisciplinary research project was thus launched at Atella in 2015, aiming to understand the site infilling and to clarify the archaeological context of the sequence. This work was rendered possible by the advancement of stratigraphic and palaeoenvironmental research on Middle Pleistocene volcanic and epiclastic successions outcropping at the Monte Vulture volcano and in the adjacent Atella and Venosa fluvio-lacustrine basins (Schiattarella et al., 2005; Giannandrea et al., 2006; Giannandrea, 2009; Schiattarella et al., 2016). New excavation campaigns were carried out and the material (lithic and fauna) preserved in the Melfi Museum collection was re-studied.

103 The purpose of this paper is to present the last five years of research at the site. Geochronological, 104 palaeontological and palynological investigations have been conducted to shed light on the 105 environmental and chronological context of the site and the associated human occupations levels.

106 This first essential step enables us to evaluate the nature and potential of archaeological data in order107 to place the results in a broader framework.

108 2. Regional setting

109 2.1 History of research at Atella

Since the end of the nineteenth century, the Monte Vulture region, and in particular the Venosaand Atella basins (see SM1).

Since the 2000s, research focused on the upper part of the stratigraphy, and revealed an alluvial environment with a high composition of volcanoclastic sediments (Borzatti von Löwenstern, 2011). In parallel, Borzatti von Löwenstern's team installed explanatory panels around the site and published a series of books geared towards the general public (Borzatti von Löwenstern, 1985; Borzatti von Löwenstern and Sozzi, 2001). The systematic excavation by the Florentine research group ended in 2012.

Renewed interest in the non-bifacial component of Lower Palaeolithic lithic industries led to a technological and techno-functional study of all the lithic industries from level F of the Atella site (Abruzzese, 2014). From 2017 to 2021, research at Atella was mainly funded by the École française de Rome as part of a larger programme on settlement dynamics during the Lower Palaeolithic in Italy (Rocca et al., 2018, 2020; Rocca and Aureli, 2019). Currently, the site of Atella is part of a valorization project funded by local and European institutions.

124 2.2 Geological context

The study area is in a volcanic district of Southern Italy, characterized by a complex stratigraphy of interbedded pyroclastic and sedimentary deposits spread over a 750 km² area along the external (i.e., north-eastern) sector of the Apennines chain (Figure 2, a). More precisely, the *Cimitero di Atella* site is located at the top of the fluvial-lacustrine succession of the Atella Basin (Giannandrea et al., 2006), about 10 km south of the Monte Vulture volcano (Figure 2, b). 130 The volcanic sequence has been subdivided into unconformity-bounded stratigraphic units (UBSU), in turn grouped into two Middle Pleistocene Supersynthems, labelled Monte Vulture and 131 Monticchio (Tab. 1). The Monte Vulture Supersystem includes the products of the central volcano 132 133 (subdivided into the Foggianello, Barile, and Melfi synthems), the underlying alluvial sediments, and 134 the fluvial-lacustrine infill of the Atella and Venosa basins (Figure 2), whereas the Monticchio Supersynthem comprises products correlated to some small eruptive vents. The age of the Monte 135 Vulture Supersynthem ranges from 698 ± 8 ka to 573 ± 4 ka, whereas the time-span of the Monticchio 136 137 Supersynthem is between 494 ± 5 ka and 141 ± 11 ka (Giannandrea et al., 2006; Villa and Buettner, 2009). 138

The lacustrine silty-sandy clay and the heteropic alluvial conglomerate of the Atella Basin i) cover the ignimbrite deposits of the Fara d'Olivo Subsynthem (Table 1), ii) are correlated with the pyroclastic and fall deposits of the Rionero Subsynthem, and iii) are covered by lava and pyroclastic flows of the Vulture - San Michele Subsynthem (Giannandrea et al., 2014; Schiattarella et al., 2016; Table 1). The latter are overlain by epiclastic deposits of the Piana del Gaudo and by travertines of Atella village. (Figure 3).

145 3. Material and methods

The new research on the Palaeolithic site of Cimitero di Atella begin in 2014 after excavations by the Borzatti von Löwenstern team had stopped. In this section we will present the material studied in this paper and the methods employed. These include excavation methods in the field, geological, geochronological and palaeoenvironment data, and the sampling and study methods applied to faunal and lithic remains.

151 3.1. Excavation method

The excavation strategy had to take into account constraints linked to the state of preservation of the site. During the initial excavation years in the early 1990s, the Borzatti von Löwenstern team decide to build a first protective cover. It consisted of a roof covering the excavation area, supported

155 by a first module made of metal tubes. Then, over the following years, a series of interventions were carried out. The covered surface was extended towards the cemetery wall, in order to increase the 156 157 excavation surface. When the new research project began in 2015, the site was therefore covered by 158 the metallic protective structure (SM2a) and included more than 2 metres section limiting the 159 excavation possibilities. Moreover, as some archaeological finds were still exposed on the excavation surface, in 2015 we first of all focused on the restoration and when possible the recording of finds. 160 161 During these two preliminary years of the project, we concentrated in particular on the revaluation of 162 the chronostratigraphic context, by drilling two cores, one from each side of the site, and collecting samples for radiometric dating. In 2016, 2017 and 2018, we began the excavation of several square 163 164 metres in the F levels inside the Borzatti von Löwenstern area, and in 2019 we opened a new 25 m² 165 area to the east of the former excavation (Figure 4). In 2020, we were able to enlarge the excavation area owing to a project promoting archaeology led by the Atella municipality. The Borzatti von 166 167 Löwenstern protective structure was dismantled and a new area was opened with mechanical diggers. The excavation area is now 250 m² and includes the Borzatti von Löwenstern area and the 2019 area 168 (Figure 4), which is now covered by a new adapted covering roof. 169

In each geological unit, the archaeological excavation proceeded by artificial levels of 5 cm, per quarter of a square metre, in order to locate the small elements found in the sediments during sieving. The three coordinates (x, y, z), the direction, dip and dip angle were recorded for all remains larger than 1 cm. The exploitation of these data is still in progress, and together with the micromorphological analysis, will provide information on the formation process of the levels.

175 3. 2. Geology

The stratigraphy of the exposed sediments of the Cimitero di Atella site was studied and the sediments from the drilling cores were analysed (Figure 4a). The detailed sedimentological analysis, which took into account grain size, sorting, and primary sedimentary structures, brought to light a series of lithofacies grouped into two associations, briefly described in Table 2. The facies were determined and interpreted on the basis of papers by Miall (1978), Walker and James (1992), Fisher

181 and Schmincke (1994), Sanders et al. (2009), Nehyba and Nývlt (2014), and Palladino et al., (2018). 182 In addition, some layers were sampled for palynological and geochronological analyses. In order to 183 establish lateral stratigraphic correlations between the deposits from the Cimitero di Atella site and 184 the Monte Vulture sequence, we performed a detailed field survey in the Atella village area, studying a topographic profile across the Atella plateau and drilling two cores (Figure 3, Figure 4a). From a 185 geomorphological viewpoint, the investigated site is located on a flat-topped hill, overlain by 186 187 travertine deposits. A straight scarp crosses this sub-horizontal land surface, due to the presence of a 188 high-angled normal fault cutting into the entire succession (Figure 5).

189 3. 3. Dating

Four sedimentary layers (Figure 6), labelled Atella INF (ash flow deposit Tng, 5 m below the archaeological excavation), F, I and L2, were dated by ⁴⁰Ar/³⁹Ar on single grains at Cimitero di Atella. In addition, two levels, L1 and L3 (Figure 6), were sampled for ESR dating of optically-bleached quartz grains.

194 $3.3.1.^{40}Ar/^{39}Ar$

195 The samples were crushed and sieved, and crystals were extracted from the 500-250 µm fraction size. They were then cleaned in distilled water using an ultrasonic bath. Unaltered and pristine 196 197 potassic feldspars (mainly sanidines) were then handpicked under a binocular microscope and leached with a 7 % HF solution for about 5 minutes to remove potential particles and alteration phases from 198 199 the surface of minerals. Thirty to fifty crystals were selected to be irradiated. They were loaded into 200 aluminium discs. Prior to mass spectrometric measurements, samples were activated in two distinct irradiations. All the samples were irradiated in the β 1 tube of the Osiris reactor (French Atomic Energy 201 202 Commission, Saclay France). Sample Atella INF was irradiated for 60 min (IRR 99) while samples F, I and L2 were irradiated for 90 min (IRR 108). Interference corrections were based on the 203 204 nucleogenic production ratios given in Guillou et al. (2018) for Osiris. After irradiation, crystals were transferred into a copper sample holder and individually loaded into a differential vacuum Cleartan© 205 206 window. The analytical procedure is described in detail in Nomade et al. (2010). Minerals were fused 207 one by one using a 25 Watts Synrad CO2 laser at about 10 to 15 % of nominal power. The extracted 208 gases were then purified for 10 min by two hot GP 10 and two GP 50 getters (ZrAl). Argon isotopes (40Ar, 39Ar, 38Ar, 37Ar and 36Ar) were successively measured using a VG 5400 mass spectrometer 209 210 equipped with an electron multiplier (Balzer SEV 217 SEN). Each argon isotope measurement 211 consisted of 20 cycles of peak-hopping. Neutron fluence J for each sample was calculated using co-212 irradiated Alder Creek sanidine standard (ACs at 1.1891 Ma, optimization calibrated age of Niespolo et al., 2017) and the 40K total decay constant of Renne et al., 2011). For the two irradiations (IRR 99 213 214 and 108), J-values were computed from a monitor co-irradiated with each dated sample (Atella INF: 215 $J = 0.000396998 \pm 0.00000199$; F: $J = 0.00039550 \pm 0.00000159$; I: $J = 0.00038743 \pm 0.00000155$. 216 L2: $J = 0.0004186 \pm 0.00000029$). Mass discriminations were monitored by the analysis of air pipettes throughout the analytical period, and relative to a ⁴⁰Ar/³⁶Ar ratio of 298.56 (Lee et al., 2006). 217 218 Procedural blank measurements were taken after every two or three unknown samples. For a typical 219 10-minute duration of isolation, typical backgrounds are about 2.0-3.0 x 10-17 and 5.0 to 6.0 x 10-19 220 moles for 40Ar and 36Ar, respectively.

3.3.2. ESR on bleached quartz

222 ESR analyses of quartz grains were performed using the multi-centre approach, based on the 223 measurement of both aluminium (Al) and titanium-lithium (Ti-Li) centres in a given quartz sample 224 (Toyoda et al., 2000; Tissoux et al., 2007). Indeed, these two ESR centres display different behaviours with regard to light on the one hand and irradiation on the other hand. Ti-Li centres are quickly and 225 226 totally bleached by solar light (in a few days), and are also much more radiosensitive than Al centres, 227 and saturate under irradiation much faster than the latter (Duval and Guilarte, 2015). In contrast, Al 228 centres are very stable under irradiation but are not totally reset even after long light exposure (several months). Therefore, a more complex protocol is required to determine the unbleachable part of the 229 230 dated quartz grains (Voinchet et al., 2004).

Firstly, the quartz grains were extracted from the samples using the chemical and physical protocol described by Voinchet et al. (2020). After extraction, each purified quartz sample was split into eleven aliquots. Nine of them were irradiated with a γ 60Co source (Commissariat à l'Énergie Atomique, 234 Saclay, France) at doses ranging from 264 to 12,500 Gy. One aliquot was conserved as a natural reference and the eleventh aliquot was exposed during 1,600 h to light in a Dr Honhle© SOL2 solar 235 simulator (light intensity between 3.2 and 3.4 105 Lux) in order to determine the unbleachable portion 236 237 of the ESR-Al signal. Each sample set of eleven aliquots was measured at least three times by ESR 238 at 107K using a Bruker[©] EMX spectrometer and every measure of each aliquot was undertaken three times after an approximatively 120° rotation of the tube in the ESR cavity, in order to consider angular 239 dependence of the signal due to sample heterogeneity. The ESR acquisition parameters used were 5 240 241 mW microwave power, 1024 point resolution, 20 mT sweep width, 100 kHz modulation frequency, 0.1 mT modulation amplitude, 40 ms conversion time, 20 ms time constant and 1 scan. 242

Signal intensity was then measured between the top of the first peak at g=2.018 and the bottom of the 16th peak at g=2.002 of the Aluminium hyperfine structure and by measuring the difference between the peak top (g= 1.913) of the Ti-Li signal and the baseline (Toyoda and Falguères, 2003). Equivalent doses (DE) were then determined from the obtained ESR intensities versus the dose growth curves using a coupled exponential and linear function with Microcal OriginPro 8 software with $1/I^2$ weighting (Voinchet et al., 2020).

External alpha and beta contributions to the dose rate were obtained using the dose-rate conversion 249 factors of Guérin et al. (2011) from the sediment radioelement contents (U, Th and K) measured in 250 251 the laboratory by high-resolution and low background gamma-spectrometry (Table 3). Gamma dose 252 rate was determined by in situ measurements with an Inspector 1000 (Canberra©) gamma spectrometer (Canberra©) using the threshold method (Mercier and Falguères, 2007). A k-value of 253 254 0.15 ± 0.1 (Laurent et al., 1998), alpha and beta attenuations from Brennan (2003) and Brennan et al. (1991), sediment water content estimated from the difference in mass between the natural sample and 255 the same sample dried in an oven at 40°C for a week and water attenuation formulae from Grün 256 257 (1994) were used in the age calculation. The cosmic dose rate was estimated from the equations of Prescott and Hutton (1994). The internal dose rate was considered to be negligible because of the low 258 259 radionuclide content usually found in quartz grains.

260 3. 4. Fauna

A preliminary identification was carried out on 621 faunal remains stored at the Heritage Office in 261 262 the Castle of Melfi (PZ). These remains are from Borzatti von Löwenstern's fieldwork and the new excavations carried out in 2015 and 2016. Most of the identifications are limited to the taxonomic 263 family or genus. Further work will attempt to refine the taxonomy of faunal remains from Atella. A 264 taphonomic approach was applied to a subsample, excluding specimens with modern fractures, or 265 incrusted by concretions. Specimen size was considered for 585 specimens, and the state of 266 267 preservation was observed on 600 remains. In particular, considering the geological setting of the site, 268 we classified specimens according to their level of rounding: i) specimens with no rounding; ii) specimens with slightly rounded edges; iii) specimens with rounded edges or completely reshaped by 269 270 agents of transport (probably water).

271 3. 5. Palynology

Fourteen samples were taken from core 2 of Cimitero di Atella for pollen analysis, from the following depths: 5.60, 6.80, 9.20, 9.60, 10.80, 11.20, 12.03, 12.40, 12.80, 13.20, 13.60, 14.02, 14.40, and 14.80 m. They were chemically treated with HCl (37%), HF (40%) and NaOH (10%) according to standard procedures for pollen extraction summarized in Magri and Di Rita (2015). Pollen concentrations were determined by adding tablets of spores of exotic *Lycopodium* to known weights of sediment. Observations were carried out by means of a light microscope at ×400 magnifications. For each sample, at least two slides were prepared and analysed.

Unfortunately, all the samples, excluding the sample at 14.02 m, were barren. In the sample at 14.02 m, after a series of three analysed slides, only one pollen grain of Poaceae was counted, which is clearly insufficient to define the past vegetation of the site. The lack of pollen grains can be ascribed to problems of pollen preservation, a very high sedimentation rate which would have diluted the pollen content within the sediment, or barren vegetation due to harsh glacial conditions. However, it is also likely that all these factors contributed to the absence of pollen.

11

285 3. 6. Lithic artefacts

The studied corpus is from the previous excavation led by Borzatti von Löwenstern and from the 286 new 2015 to 2018 excavations (Table 4). The old collection contains two main assemblages, 287 288 excluding surface finds, one from Level L, the other from Level F. Given the disparate states of 289 alteration, and some doubts concerning the anthropic nature of several pieces, only the more reliable 290 items were selected (Abruzzese et al., 2016). The following criteria were retained to improve the 291 reliability of the assemblage: type of blank and raw material, the degree of alteration, the type of patina and the presence of a double patina, and the presence or absence of knapping scars. We 292 293 subdivided the material into three groups, definite non-anthropic items, probable poorly-preserved 294 anthropic elements, and anthropic items. Only the last group was analysed. In level L, only 107 pieces 295 were selected out of 2553 (Borzatti von Löwenstern, 2005), and 666 of the 7887 pieces from level F 296 (Abruzzese et al., 2016).

Raw materials can be divided into three types: flint, quartzite and a highly altered raw material,
described as "porous radiolarite" (Borzatti von Löwenstern et al., 1997), which is probably a fine
limestone.

300 A technological approach was applied to the lithic assemblage of Atella (Boëda et al., 1990; Inizan 301 et al., 1999; Boëda, 2013), in order to describe the reduction sequence (selection, production aims, 302 use, management, etc.). In this paper, we focus on the production phase and the identification of the main technological categories: cores, debitage flakes, small tools, large tools, retouch flakes, based 303 304 on the diacritical scheme of each piece (chronology and direction of removal, measurements, angles, 305 etc.). We also began to analyse some piece with a techno-functional analysis (Lepot, 1991; Boëda, 2013; Aureli et al., 2016; Rocca et al., 2016) in order to identify the transformative and prehensile 306 307 part of the tools.

308 4. Results

309 4. 1. Stratigraphic and sedimentological analysis

310 The horizontally-bedded Quaternary succession outcropping in the study area unconformably 311 overlies the sedimentary bedrock, represented here by Numidian Flysch (age: Burdigalian -312 Langhian?). Miocene sediments only outcrop to the east of Atella village (Figure 3) and consist of 313 tilted quartzarenite beds. The Quaternary sequence of the site (Figure 6) is formed, from the base, of 314 light-grey horizontally-laminated volcaniclastic siltstone of the Sh lithofacies (Figure 7 and Figure 315 7a), attributed to sediment decantation in a lacustrine environment. Toward the north, such sediments 316 are in heteropic relationships with volcaniclastic alluvial conglomerates (Figure 3a). Above them, a 317 lava flow and pyroclastic and epiclastic deposits, labelled 3a, 3b and 4 in Figure 3, and swamp and 318 alluvial deposits, are recorded in the surveyed area. Finally, the travertine of Atella village (Figure 3) 319 unconformably overlies the previously described deposits, through an erosive surface.

320 In the Cimitero di Atella site, at the top of facies **Sh**, the base of the stratigraphic succession (Figure 321 6) is composed of horizontal beds of the **TLm** lithofacies (Figure 7b), interpreted as pyroclastic ash 322 flow deposits, with intercalations of fallout ash deposits, attributed to lithofacies Tng (Figure 7c) and 323 **Tm** (Figure 7d). Laterally (i.e., towards the north and to the west of Atella village), the pyroclastic 324 beds overlie a lava flow (Figure 3) attributed to the Vulture - San Michele Subsynthem by 325 Giannandrea et al. (2006). Moving up, about 2-m-thick lacustrine deposits of lithofacies Sh, Dh, and 326 Su (Figure 8a) are overlain by two epiclastic breccia (Figure 7e) with volcanic clasts (lithofacies 327 Bmli), connected to cohesive debris flow. Each breccia layer is covered by reworked sediments of 328 the Suli lithofacies (Figure 7e). Both lithofacies Bmli and Suli contain lithic industry and two tusks of Palaeoloxodon antiquus (Borzatti von Löwenstern et al., 1997). A westward-dipping erosive 329 330 surface is present at the top of these sediments (Figure 7).

The upper portion of the succession is characterized by alternating beds of primary volcanic deposits – made up of ash and lapilli fallout from lithofacies **Tm** and **Lm** (Figure 8d) and pyroclastic flow from lithofacies **LTm** (Figure 8b), **TLm**, and **TLBm** (Figure 8c) – and trough cross-bed epiclastic volcanic sediments corresponding to lithofacies SBt and Sr (Figure 8b and Figure 8d). It is noteworthy that Lm and SBt facies are frequently associated in the same horizon. Often, the beds at the base of lithofacies LTm, TLm, TLBm, and Bmli show water-escape pipes. These are softsediment deformation structures due to the unstable density of the water-saturated alluvial (Figure 8c and Figure 8d) and lacustrine (Figure 8) sediments. Here, such structures can be interpreted as loading structures resulting from the rapid sedimentation of mass deposits (Anketell et al., 1969).

340 This suggests that the deposition of facies **Bmli** occurred immediately after the lake was drained, 341 when sediments were still saturated. Is also likely that the cohesive debris flow deposits (facies Bmli) 342 were caused by the activation of water flows associated with lake drainage. The presence of such a 343 sedimentary structure shows continuity in sedimentation at the transition from lacustrine to alluvial deposits. Therefore, no stratigraphic unconformity is present at the base of that facies. Soft-sediment 344 deformation structures are also present in the upper part of the Cimitero di Atella succession (Figure 345 9). They consist of boudinage-like breccia structures, made of ash elements in alluvial epiclastic SBt-346 347 facies sandstone (Figure 8d and Figure 9) and represent a typical case of seismite. Therefore, there 348 must have been synsedimentary seismic activity at that time in the study area (e.g., Pope et al., 1997; Gibert et al., 2011). The above-described facies association can be referred to as a braided alluvial 349 plain with abrupt contributions of primary volcanic products by fallout process and connected to mass 350 351 deposition by pyroclastic flows.

Two fallout ash deposits (lithofacies **Tm**), separated by an eluvial-colluvial horizon (Figure 6), are present at the top of the pyroclastic/epiclastic sequence of the Atella site. Here, a southeast-dipping erosive surface intersects the whole sequence, cutting out a space subsequently filled by historical sediments (Figure 9).

356 4. 2. Dating

Atella INF - Ten crystals were individually dated. Nine of them provided a statistically equivalent age within uncertainty, confirming the primary volcanic nature of the deposit. The weighted mean age, calculated using a juvenile crystal population, is 655.2 ± 12 ka (MSWD = 0.2 and P = 1.0, full external errors at 2σ uncertainties). The ⁴⁰Ar/³⁶Ar initial ratio given by the inverse isochron of 290.8 \pm 56 (Table 3) is very unprecise but equivalent within uncertainty to the atmospheric ratio of 298.56 (Lee et al., 2006).

Atella F - Twelve sanidines were individually dated for this layer. The probability diagram obtained for this sample is multimodal and shows a high percentage of reworked volcanic minerals. Only six crystals constitute the homogeneous and youngest population. At least four older eruptive successions are evidenced. The weighted mean age calculated for the youngest population is 577.5 ± 6.4 ka (MSWD = 1.1 et P = 0.3, full external errors at 2σ uncertainties. The 40 Ar/ 36 Ar initial ratio given by the inverse isochron for this population is 299.4 ± 4.8 (Table 3), equivalent within uncertainty to the atmospheric ratio of 298.56.

Atella I - For this level, unfortunately only six crystals of potassic feldspars were found and successfully dated. Despite the limited number of dated crystals for this layer, these analyses provided significant information. The related probability diagram shows crystals dated from 715 to 575 ka. This crystal age dispersion is very similar to that of level F.

Atella L1 – ESR analyses of this sample provide similar equivalent doses and ages for both Al and Ti-Li centres, indicating good initial bleaching of both signals. A weighted quadratic mean age of 509 \pm 56 ka (2 σ , full external error, MSWD = 0.051, probability = 0.82) was calculated for this sample using IsoPlot 3.0 software (Ludwig, 2003).

Atella L2 - Thirteen sanidines were individually dated. Again, the related probability diagram is multimodal. Two major crystal populations are evidenced, as well as three older crystals ranging between about 650 and 720 ka (see Figure 10). The younger population included six crystals and gave a weighted mean age of 583 ± 4 ka (full external errors at 2σ uncertainties) (MSWD = 0.9 et P = 0.5 (11 ka). The older crystal population is dated to 610 ± 3 ka. The 40 Ar/ 36 Ar initial ratio of 297.6 \pm 3.0 for the younger population (see Table 3, 2σ analytical uncertainties) is equivalent to the atmospheric one (Table 3).

385 Atella L3 - Once again, equivalent doses and ages determined from both Al and Ti-Li centres

correspond, confirming good initial bleaching of the quartz grains and allowing for the calculation of a weighted quadratic mean age of 442 ± 39 ka (2σ , full external error, MSWD = 0.35, probability = 0.55).

389 4.3. Archaeological sequence

390 Our first hypothesis for the archaeological sequence is based on the results of the sedimentological 391 analysis and the first taphonomic observations on faunal and lithic remains. The Borzatti von 392 Löwenstern team initially proposed (Borzatti von Löwenstern et al., 1990) dividing the Cimitero di Atella sequence into 15 geological layers, and two large archaeological units: F and L (Borzatti von 393 394 Löwenstern et al., 1997). According to Borzatti von Löwenstern, the F level was the result of a 395 lakeshore occupation and level L a redeposition by the river of earlier occupations without handaxes 396 (Borzatti von Löwenstern and Sozzi, 1994; Borzatti von Löwenstern, 2005). Our aim was to re-397 evaluate site formation and to try to interpret the archaeological records of each level (Figure 11). The combination of an applied excavation method, the study of archaeological finds, and preliminary 398 399 sedimentological results yields a revised interpretation of the sequence. In the current state of 400 knowledge, we are able to identify four main archaeological units (F-Base, F-Deb2, F-Deb1, L), 401 resulting from different formation processes.

At the base of the sequence, we identified F-Base, a first archaeological layer partially excavated 402 by the previous team over about four square metres. This layer is still exposed but was not investigated 403 404 by us and our interpretation is thus only based on field observations. It contains a high concentration 405 of remains, a *Palaeoloxodon* tusk and other well-preserved macro-faunal remains, and lithic finds consisting of a few handaxes and many small tools and flakes. This horizon lies on top of a palustrine 406 407 diatomic laver (Figure 11). Human occupation occurred in level F-Base after the retreat of the lake 408 and was covered by a reworked colluvium of fine sediment composed of epiclastic volcanic 409 sandstone, with volcanic angular to sub-angular pebbles (ranging in size from 1 to 10 cm). 410 Preliminary observations on well-preserved remains (fauna and lithic industries), the combined 411 presence of small and large elements, along with the high concentration of finds in fine sediment,

412 indicate that this level is the result of a sub-primary occupation. This hypothesis needs to be confirmed413 by further analyses after the resumption of excavations in the field.

414 The second archaeological unit (F-Deb2) was extensively excavated by Borzatti von Löwenstern over a surface of 25 m² and also by our team over 4 m². The material from this level corresponds to 415 416 the base of level F of the Borzatti von Löwenstern excavation, with a lithic assemblage studied by 417 Abruzzese (et al., 2016). The lithic and faunal remains recorded in this level are in a poor state of 418 preservation. The lithic remains present smoothed edges, often with double patina. The fauna is 419 fragmented and rounded. The concentration of remains is very high in a poorly-sorted matrix supported epiclastic volcanic breccia, interpreted as a cohesive debris flow deposit. Thus, we 420 421 postulate that the deposit containing the archaeological record is the result of the reworking of one or 422 several occupations in a secondary position. At this stage, it is difficult to evaluate the degree of reworking of the deposit as well as distance from the primary occupations. Refitting should help to 423 424 determine whether this corresponds to a major or a light mass movement of the initial occupation.

At the top of this reworked deposit, we identified a third archaeological layer covered by the I 425 tephra (F-Deb1). This level was initially called H by Borzatti von Löwenstern and then mixed 426 together with the material from layer F. It was partially excavated by us over a surface of 7 m^2 . The 427 428 thickness of this level varied between a few centimetres to 15 cm, as the top was affected by erosion 429 before being covered by the overlying tephra layer (Figure 8). The lithic and faunal material recorded 430 in this layer is well preserved. The lithic finds present rather fresh edges and no double patina. The 431 assemblage contains a large tool in limestone, small tools and small flakes of different sizes. The 432 faunal remains are less fragmented and rounded and some pieces present fresh surfaces, as for 433 example two Cervidae remains (a coxal and a rib). The archaeological remains are in a fine sediment 434 of epiclastic volcanic sandstone, interpreted as a reworked low-intensity colluvium. The distribution 435 of the archaeological finds, the concentration of material, the presence of small and large pieces and the state of preservation of the remains indicate that the material has not undergone major reworking. 436 We propose that this layer is in a sub-primary position, partially intersected by erosion and quickly 437

438 covered by the I tephra layer. Further investigations (use-wear analysis, extension of the excavation, 439 refitting, taphonomic analysis on the fauna, etc) should clarify the degree of coherence of this level. 440 The last Pleistocene occupations recorded in the site of Atella are in the upper part of the sequence 441 (L). The archaeological finds were found in a coarse epiclastic volcanic sandstone and angular to sub-442 rounded epiclastic volcanic clast-supported breccia, interpreted as a channel infill by water stream 443 flow of reworked volcanic fallout and pyroclastic flow deposits. This part of the sequence was almost 444 exclusively excavated by the previous team, apart from a small pit in the new area with only a few 445 very small debris found during sieving. The studied lithic material of the Borzatti von Löwenstern 446 assemblage is differentially preserved. Some pieces look very fresh whereas others are smooth and 447 polished. The faunal remains are also differentially preserved. At this stage of the study, this large unit appears to contain material from partially preserved sporadic occupations on a river shore, with 448 449 differential sorting and erosion depending on the flow of the river.

450 Thus, based on the results obtained so far, we can identify four moments of occupation at Atella. These layers may be very different in terms of duration, as the F (Deb2) and the L accumulations 451 probably result from several occupations, and it is thus difficult to evaluate their duration. The 452 presence of a palimpsest in secondary position was already known, and we identified at least one 453 well-preserved layer between the F accumulation and the I tephra layer. Given these new and 454 455 promising data, further investigations at Atella will focus on the excavation of this horizon over a 456 larger surface. Another point to clarify will be the estimation of the time range between occupations, as for the time being, the resolution of absolute dating provides no information concerning the 457 458 timespan of each unit, nor the period of time between them. Further analyses, especially on the tephra 459 I, should reduce the imprecision of the dating resolution of the sequence.

460 4. 4. Fauna

The faunal remains are highly fragmented: the most represented size classes are 2 and 3 cm (22.4%
and 19.8% of remains respectively) and 81.4% of remains fall within the 1-5-cm interval (Table 6).
Most specimens from unit L are unaltered by transport (74%). On the other hand, 38.5% of specimens

464 are rounded (i.e., bone fragments with a spherical or subspherical polished shape) in phase F and 43.4% in phase H (Table 7). As most of the faunal remains from Atella are represented by incomplete 465 466 or fragmented specimens of poor diagnostic value, taxonomic identifications are mainly limited to family or genus, namely ungulates (Bovinae indet., Cervus elaphus, Dama sp.) or Palaeoloxodon 467 468 antiquus/Palaeoloxodon-size categories (Table 8). Apart from P. antiquus, Borzatti von Löwenstern et al. (1997) reported the presence of a few bovid and cervid remains, and a hyaena tooth. Zucchelli 469 470 (1999) attributed the Bovinae remains to Bos primigenius and Zucchelli (2002) identified Cervus 471 elaphus, Dama dama, and Capreolus sp. among the cervids. The revision of the Borzatti von 472 Löwenstern material, in addition to the study of new finds, reveals that some bovid postcranial 473 elements have Bison-like features, such as, for example, metapodials (e.g., interarticular incisure Ushaped and less proximally extended in relation to the margin of both condyles; parallel or slightly 474 475 convergent medial and lateral intercondylar crests). Nonetheless, pending the discovery of new 476 taxonomically significative fossils, the attribution of the bovid material remains opens (Bovinae indet.). No specimens identified so far within the studied sample indicate the presence of hyaenas or 477 Capreolus at Atella. 478

479 4. 5. Palynology

Unfortunately, all the analysed levels, excluding those at 6.30, 6.60 and 14.02 m, were barren. After a series of three analysed slides per sample, one *Zelkova* pollen grain was identified in the sample at 6.30 m, one *Pinus* grain at 6.60 m, and one Poaceae grain at 14.02 m, but they were insufficient to define the past vegetation of the site. The lack of pollen grains can be ascribed to factors preventing pollen preservation, the dilution of the pollen content in the sediment due to a very high sedimentation rate, or the occurrence of barren vegetation due to harsh glacial conditions. However, it is also possible that a combination of factors may have contributed to the absence of pollen.

487 4. 6. Lithic industry

The analysis of the lithic assemblage confirms the presence of the three main reduction sequences
already identified in the Borzatti von Löwenstern assemblage (Abruzzese et al., 2016): small tools,

490 debitage flakes and large tools (Table 5).

491 The first is aimed at small tools confection. This reduction sequence is therefore represented by small tools (266 items, see Table 5) and confection small flakes, namely notch flakes and retouch 492 493 flakes (268, see Table 5, Figure 12a). The first step consists of blank selection for the future tool. The 494 selected blanks, when determinable, can be small blocks of raw material, flakes, or more rarely residual core fragments. The study of the morphological and volumetric characters of the blank before 495 496 tool confection brought to light important criteria, such as the small dimensions, the relative thickness, 497 and the presence of a flat surface. Another element can also be present on the initial blank, such as an 498 abrupt surface, for example a back of a *debordant* flake or the butt of the flake. Then, the tool 499 confection phase depends on how the degree of affordance of the blank matches the sought-after tool. 500 This phase consists of shaping and retouching the blank to obtain several cutting edges. The removals 501 generally come from the flat surface of the blank and concern the transformative part of the future 502 tool and in some cases the prehensile part. On the cutting edges, several modalities of confection are 503 performed from short, low and parallel removals to notched abrupt retouch. The management of the 504 prehensile part is rarer and consists mainly of the correction or accentuation of an abrupt part. 505 Consequently, the flakes from the confection of these small tools are varied and depend on the features 506 of the initial blank and of the sought-after tool. Notch flakes are short and wide and characterized by 507 a thick and convex butt (Figure 12a). Retouch flakes are thin and curved, with a linear or punctiform 508 butt, and the dorsal surface can present previous retouch scars.

The aim of this reduction sequence is to obtain several cutting-edges on small tools. The tool sample is too small for the moment to describe this techno-functional group in detail. It consists mainly of rostrums, trihedral, and rectilinear edges, as observed in the previous publication (Abruzzese et al., 2016). The illustrated example (Figure 13) presents three transformative parts (TFU1, TFU2, TFU3) on the same piece, each associated with a prehensile part. The transformative part consists of a trihedron composed of a flat surface, two facets obtained by lateral notches, and a central ridge (Figure 14). Only part of the whole volume is modified to fashion the tool. The prehensile part, in this case a prepared back, can be adapted to several transformative parts. From a production point of view, several gestures and probably several hammerstones were used to make this small tool. For the management of convexities and the production of the flake, blows are internal, probably made with a hard hammer. The back is obtained by an orthogonal gesture (fracture blow). The notches blow, important to highlight the trihedrons are internal and were probably made with a smaller hammerstone.

522 The second reduction sequence consists of flake production. Due to the rarity of whole cores in 523 the assemblage (25 items, mainly fragments, Table 5), it is difficult at this stage of the study to define 524 flaking modes and the objectives of flake production based on core analysis. The few cores belong to 525 an additional production system (Boëda, 2013; De Weyer et al., 2022). The exploited volume of the 526 cores only represents a small part of the initial blank. No cores show global preparation of volume, or partial management of convexities. The flaking sequence is very short, with no more than five 527 528 removals per core. Flake scars indicate the production of elongated and quadrangular flakes, with 529 large butts, two cutting edges or one cutting edge and a back. The entire core and fragments are in 530 flint and quartzite. This scant information from the cores is consistent with data from flakes. The latter (401 items, Table 5) are elongated or quadrangular, with one or two scars on the dorsal surface 531 532 and often present a natural back (Figure 12b). These large flakes in flint and quartzite are never retouched, but present one or two cutting edges. The characteristics of these flakes, dimensions and 533 534 technical features are different from the small flakes issued from making small tools. This category of large flakes is compatible with core attributes, indicating that they are flakes from the debitage 535 536 reduction sequence. Some of these flakes may have been used as blanks for small tools, but as the 537 latter are highly reduced, it is difficult to identify the initial blank. The debitage reduction sequence 538 can also be interpreted as a goal in itself, to obtain a longer cutting edge. This component is almost 539 always present in Lower Palaeolithic assemblages. Further investigation into this debitage reduction 540 sequence and the relationship between small tools and flakes is required in the future, on a larger 541 sample, particularly of cores.

542 The large shaped tools constitute the third reduction sequence. This category comprises what we 543 can call, from a typological point of view, Large Cutting Tools, handaxes, unifacial pieces, and pebble 544 tools (28 items, Table 5). These tools are mainly made on poorly-preserved limestone pebbles and analysis is therefore difficult. From a production point of view, the chosen blank is plano-convex, and 545 546 can be a pebble or a large cortical flake. Shaping is mainly limited to the upper apical part of the 547 blank, and the basal part is unretouched. The shaping phase takes into account the volumetric characteristics of the chosen blank and only affects the future transformative part, as the prehensile 548 549 part is rarely modified. This very short shaping phase does not affect the volumetric structure of the blank. We were not able to identify flakes from this reduction sequence. This is maybe due to the poor 550 551 state of preservation of the limestone, particularly affecting on thin and small shaping flakes. It is also 552 important to recall that macro-tools are very rare in the assemblage and bear a low number of scars. The technical features of large tools show that shaping flakes may present a cortical, semi-cortical 553 554 surface or positive scar on the dorsal surface. Due to shaping modes, the classical expected features for shaping flakes, such as a curved profile, an open angle between the ventral surface and the butt, 555 and soft hammer percussion, may not be determinant here. Indeed, flakes from large tool shaping do 556 not bear specific features, and it is thus difficult to distinguish them from other flakes. 557

The identification of this techno-functional group is problematic due to the low number of large tools. At this stage of the analysis, the main categories of large tools are represented by distal rostrums and in one case by a lateral cutting edge (Figure 15).

Thus, the assemblage of Cimitero di Atella studied so far is composed of three main reduction sequences. The main one, in term of the number of items and techno-functional variability, is represented by the small tools. Large shaped tools and backed flakes are less frequent and for the moment, we can go no further with our interpretation due to the limited quantity of elements. 566 5. 1. Site of Atella

567 *5.1.1 Geology*

568 As a result of field investigations, following the stratigraphic framework proposed by Giannandrea 569 et al. (2006) and Schiattarella et al. (2016), we correlated the basal 1-m-thick portion of lacustrine sediments (facies Sh in Figure 6) with the Rionero Subsystem and, consequently, we were able to 570 571 correlate the overlying volcanic (facies Tm, Tng, LTm, and TLm) and lacustrine layers, up to the 572 epiclastic layers of the facies **Bmli** (unit F of Borzatti von Löwenstern et al., 1997), with the Vulture 573 - San Michele Subsynthem. The presence of soft-sediment deformation structures in lacustrine sediments below the Bmli facies suggests that the deposition of unit F of Borzatti von Löwenstern (et 574 575 al, 1997) occurred immediately after the lake was drained, when the fine sediments were still 576 saturated. It is also likely that the cohesive debris flows (facies Bmli) were deposited by the activation of water flows associated with lake drainage. This sedimentary structure shows continuity in 577 578 sedimentation at the transition from lacustrine to alluvial deposits. Therefore, no stratigraphic 579 unconformity is present at the base of the archaeological fluvial facies. Also, sedimentological 580 analyses performed in unit L of Borzatti von Löwenstern (et al., 1997) reveal tephra layer 581 intercalations, interpreted as pyroclastic flow, in the fluvial facies. Considering the presence of seismically-induced soft-sediment deformation structures, this can be ascribed to particularly intense 582 583 and voluminous eruptions of the Monte Vulture volcano.

584 *5.1.2 Geochronological interpretations*

Based on the geology, archaeology, and palaeontology of the site, Borzatti von Löwenstern et al. (1997) proposed a date for the Cimitero di Atella sequence between about 600 and 500 ka. Later, Di Muro (1999) suggested correlating stratigraphic levels A to E with the Rionero Subsynthem of Monte Vulture (i.e., 630 ± 20 ka to 714 ± 18 ka following Villa and Buettner, 2009). While our new individual data obtained by 40 Ar/ 39 Ar and ESR on bleached quartz appear insufficient for accurately determining the ages of the Cimitero di Atella archaeosurfaces, combining this new geochronological dataset with the available geological information helps to constrain the period of human occupations with greaterconfidence and accuracy.

Concerning the ⁴⁰Ar/³⁹Ar analyses, the lowest level dated by this technique, Atella INF (Tng 593 594 primary volcanic layer, see Table 3, Figure 10) is accurately dated to 655 ± 12 ka, the maximum age 595 for the entire sequence of the Cimitero di Atella archaeological site. The analytical data for levels F, I, and L2 are more difficult to interpret, as all the corresponding probability diagrams are complex 596 597 and multimodal (see Figure 10), suggesting the massive reworking of volcanic materials from several 598 eruptive events, the youngest of which is dated to around 570-580 ka. The various ages of the oldest 599 populations of crystals can be attributed to the first part of the Monte Vulture volcano eruptive activity 600 (between about 610 and 760 ka), corresponding to the San Michele and Rionero Subsynthem 601 activities (Villa and Buettner, 2009). For these three dated stratigraphic levels, the youngest sanidines are all dated to around 580-570 ka. This age can only be considered here as a maximum age for the 602 603 deposition of the Cimitero di Atella archaeological horizons. It is noteworthy that even if the Monte 604 Vulture volcano was very active from the beginning of the Middle Pleistocene until 490 ka, it was nonetheless characterized by a long phase of quiescence between about 570 and at least 530 ka, 605 606 followed by an eruptive sequence after local tectonic activity (i.e., Case Lopes subsynthem). 607 Therefore, the fact that no eruptions more recent than about 580-570 ka were recorded in levels F, I, 608 and L2, can be taken as a further chronological indicator. Deposits from the Case Lopes Subsynthem 609 are dated to 530 ± 22 ka and 494 ± 5 ka (Villa and Buettner, 2009, Monticchio Supersynthem) and 610 found a few kilometres NE of the site but not found in the reworked and dated volcanic material at 611 Cimitero di Atella. Even though the absence of such deposits cannot be considered as chronological 612 proof of the antiquity of the archaeological remains, it nevertheless implies that the Cimitero di Atella 613 sequence predates the Case Lopes eruptive phase. This hypothesis is reinforced by the ESR ages 614 calculated for L1 (509 ± 46 ka) whereas L3 (442 ± 39 ka) suggests a younger age for the deposition 615 of the upper part of the sequence. Even if these ages lack precision, they provide a needed chronological framework for the deposition age of the sedimentary sequence of Cimitero di Atella. 616

They also demonstrate the continuity of the stratigraphic sequence and reinforce interpretations based on the geology and 40 Ar/ 39 Ar data. They allow us to determine a deposition age between L1 and L3 (509 ± 46 ka - 442 ± 39 ka). The age of 509 ± 46 ka also provides a minimum age for the archaeological levels underlying L1 (F unit). The combined geochronological data therefore suggest that at least the lower Palaeolithic F levels can be securely dated between 509 ± 46 ka and 570-580 ka, covering the end of MIS 15 until MIS 13 (Figure 10).

623 *5.1.3 Palaeobotany*

Previous pollen analyses at Cimitero di Atella were carried out on 15 pollen samples by Borzatti 624 von Löwenstern von Löwenstern et al. (1998). However only four levels presented enough pollen 625 626 grains to describe the surrounding environment, namely levels A01, A02, A03 and I. According to the new chronology of the Borzatti von Löwenstern sequence presented here, these levels were deposited 627 between 655 ± 12 and 575 ± 7 ka BP. The pollen content of A02, A03, and I, showing Arboreal Pollen 628 (AP) percentages above 88%, Abies up to 15%, Quercus deciduous ranging from 7 to 35%, and many 629 630 other deciduous and evergreen broadleaved trees in low frequencies, suggests a cool and wet phase of the interglacial stage corresponding to MIS 15. Conversely, in the underlying level A01, the AP 631 632 percentage (76%) with 53% *Pinus*, and high proportions of bush and herbaceous xeric taxa, such as 633 Ephedra, Artemisia, and Poaceae, suggest an arid phase during the MIS 16 glacial stage.

634 *5.1.4 Occupation modes*

In the current state of knowledge, it is difficult to accurately reconstruct the human occupations of 635 Cimitero di Atella. However, some new data contribute to a better understanding of the type of 636 occupation. Overall, the stratigraphic succession registered several episodes of human presence 637 between 0.58 and 0.44 Ma. At that time, the landscape changed from a lacustrine to a fluvial 638 639 environment in a territory highly impacted by the activity of Monte Vulture. The recording of human 640 occupations is different in term of temporality. Some layers, such as F-Deb2, are probably a mix of several occupations, of unknown duration, whereas others, like F-Base and F-Deb1, probably result 641 642 from a single "moment" of occupation, including tool making and faunal exploitation. The future 643 excavation of unit L, corresponding to the fluviatile sequence, will provide crucial information. The

644 archaeological record shows that such contexts are propitious to simultaneous faunal and human occupation and to the quick burial of their tracks. Unit L, as already observed by Borzatti von 645 Löwenstern (2005, 2011), could contain several more or less well-preserved occupation "spots", 646 647 depending on site formation taphonomy (velocity of sedimentation, the riverbed position and flow 648 intensity). The presence of rich but mixed units alongside short well-preserved occupations in the 649 sequence is interesting and is in our opinion complementary. The secondary position (colluvium) of 650 the unit is not conducive to spatial or techno-economic analysis, but the material is useful for a more 651 global view of technical variability. Conversely, the level interpreted as an occupation surface may 652 contain important information on occupation modes, spatial organization, faunal exploitation and tool 653 manufacture and use. At Atella, given the type of sedimentation, we are probably not in presence of a genuine snapshot, as observed at Ficoncella (Aureli et al., 2015; Boschin et al., 2018), Soucy 3 654 655 (Lhomme, 2007), La Polledrara di Cecanibbio (Santucci et al., 2016), and Schöningen (Serangeli et 656 al., 2015), but rather a sub-primary position site resulting from one or more occupations in a relatively short time period, like in some levels of Notarchirico (Piperno et al., 1999) and Isernia La Pineta 657 658 (Peretto et al., 2004). It is interesting to note that in both levels (F-Base and F-Deb1), like in other many contemporaneous sites, human occupation is linked to an aquatic environment (lake and river) 659 with large herbivores, especially elephants (Kodinaris et al., 2016; Rocca et al., 2021). The 660 661 relationship between humans and elephants also needs to be reassessed, beyond the hunting/scavenging debate, in order to really focus on exploitation modes and purposes. For meat? 662 For toolmaking or other technical activities? What was the impact of the elephant on the landscape? 663 664 Was the carcass a landmark in the territory? etc. We hope that the extension of the excavation surface to about 250 m² will reveal new information on site formation and spatial organization, in order to 665 enhance our interpretations of occupations at Atella. 666

667 5.2. Atella in a broader context

- 668 5.2.1. Is it relevant to speak about an Early Acheulean?
- The south European Lower Palaeolithic panorama has been extensively reshaped in recent years.

670 Until the end of the 1990s, this cultural period was divided into two lengthy phases on the basis of 671 typological studies; an older one, called the pre-Acheulean without handaxes, and a younger one, including Acheulean, Clactonian and Tayacian industries (Palma di Cesnola, 2001; Grifoni and Tozzi, 672 673 2006). New data, mainly from the sites of Isernia La Pineta and Notarchirico, call into question this 674 model. At the end of the 1990s, new dates from Notarchirico significantly pushed back the age of early handaxes in Europe (Pilleyre et al., 1999; Piperno et al., 1999). Then, conversely, in the 2010s, 675 676 the new dating of Isernia La Pineta, a site without handaxes, attributed the site to a more recent period 677 (Peretto et al., 2015). Those new data led to a reconsideration of the Lower Palaeolithic periodisation, 678 alongside the development of lithic industry study methods. The Isernia La Pineta team applied new 679 methods, such as technology, experimentation, use-wear analysis, resulting in the criticism of 680 Tayacian typology (Crovetto, 1993; Crovetto et al., 1994; Peretto et al., 1994; Longo et al., 1997). The discovery of new early sites with handaxes in Europe in the 2010s led to the construction of a 681 682 new chrono-cultural model (Moncel et al., 2013, Garcia-Medrano et al., 2014, Mosquera et al., 2016). These early occupations (Arzarello et al., 2012; Ollé et al., 2013; Michel et al., 2017; Despriée et al., 683 684 2018) brought to light a new conception of tools in Europe from about 0.8/0.7 Ma, based on the handaxe or Large Cutting-Tool, called the Early Acheulean (Moncel et al., 2015, 2020; Moncel and 685 686 Schreve, 2016). Thus paradoxically, after a short parenthesis focusing on the reduction sequence and 687 technical diversity, we are now returning to a very typological definition based on the Acheulean 688 guiding fossil: the biface. Indeed, the origins of the Early Acheulean in Europe are often implicitly 689 and schematically explained using the following model:

Presence of a phase of core and flake industries between 1.4 Ma and 0.8 Ma (Mode 1) and the arrival from Africa of new industries with handaxes (Mode 2), from 0.8/0.7 Ma onwards, spreading rapidly to the north (La Noira) and south (Notarchirico, la Boella) of the European continent. Can we really define this period on the basis of a single tool? What can other ways of making tools, such as flake production or the so-called small tools tell us? Do we have solid proof of an external arrival of this new technical tradition? Is it relevant for researchers to coin the term Early Acheulean in studies of this key period of the Palaeolithic? How can the lithic industries of Cimitero di Atella contribute toan enhanced definition of this period?

The assemblage of Cimitero di Atella studied so far is consistent with the other contemporaneous 698 699 technical system, the so-called Early Acheulean. As in other sites, abundant small tools and/or flakes 700 play an important role in tool kit variability, while large shaped tools are rare (Nicoud et al., 2016; 701 Tourloukis et al., 2018; Grimaldi et al., 2020). Furthermore, large tools are significantly different to 702 the Acheulean handaxes that emerge in the second part of the Lower Palaeolithic, in that shaping does 703 not affect the global volume of the blank and is aimed at obtaining a specific cutting edge, mainly 704 rostrums. Edge-point bifacial tools are very rare, as well as lateral cutting edges. At Atella, as 705 elsewhere, these shaped tools display wide technical diversity: each specimen is almost a specific 706 tool type. This diversity is also present for example at Notarchirico (Moncel et al., 2019, 2020; Nicoud 707 2011), and at la Boella (Mosquera et al., 2016).

708 Further discussion is required to enhance our understanding of the specificity of this ambiguous 709 category of large tools in the Early Acheulean. Nicoud (2011) proposes the term "pebble with bifacial 710 removals", as does Moncel (2019, 2020), to distinguish "bifaces and bifacial tools" from "pebble 711 tools" at Notarchirico. Mosquera (et al., 2016) groups together all the shaped pieces under the term 712 "large cutting-tools". A more in-depth investigation of this category of tools is now required, focusing 713 on techno-functional aspects. Paradoxically, early large cutting-tools are overrepresented in 714 publications, despite their small numbers, but are rarely studied in relation with other 715 contemporaneous tools and compared with younger handaxes. We choose to call these pieces from 716 Atella large shaped tools, because they cannot all be considered to be bifacially shaped from a 717 technological point of view. Yet it would be illogical to distinguish bifacial from non-bifacial large 718 tools, as they can present the same morpho-technical features. However, we can note that these pieces 719 are different from classical handaxes in terms of volumetric structure and techno-functional features. 720 Even more surprisingly, some transformative parts, such as the rostrum, for example, seem to be comparable to ones on small tools in terms of technical-functional features and techno-productional 721

modes. In terms of weight and dimensions, use, but also probably prehension modes, these tools are clearly distinguishable from their smaller counterparts. But it is nonetheless interesting to approach the relationships between several categories without always considering these first handaxes as an archaic model of future Acheulean bifacial pieces.

726 The small tool category also necessitates further discussion. The presence of small tools was 727 recognized in Tayacian typology (Tayac and Quinson point), but the variability and importance of this category was subsequently re-evaluated in Central European and Middle Eastern assemblages 728 729 (Burdukiewic and Ronen, 2003). In line with this work, we attempted to better define this part of the 730 lithic assemblage from a technological and techno-functional point of view (Aureli et al., 2016; Rocca 731 2016; Rocca et al., 2016). The term small tool can be misleading, since it emphasizes size, rather than 732 the technical reality of this specific reduction sequence. Indeed, the presence of small and thick tools on specific selected blanks with a flat surface opposed to a convex surface, with cutting edges, is 733 734 widespread. Regardless of the term, this specific tool kit is a crucial element for characterizing the technical system of this phase of Lower Palaeolithic. 735

In sum, the results obtained on the lithic assemblage of Cimitero di Atella contribute to a better knowledge of the technical system in Southern Europe. The tool kit is mainly obtained by the confection of small tools and secondarily by shaping large tools and flaking backed flakes. Many questions still need to be addressed in the future, including the selection/production of blanks and the techno-functional diversity of small and large tools. But in our opinion, in order to better comprehend this phase of the Lower Palaeolithic, we need to examine the whole assemblage, not only LCTs, from a technological perspective.

743 5.2.2. *The fauna of Atella in a broader context*

The Early Acheulean spread throughout Europe alongside considerable faunal turnover, in a broader context of remarkable environmental and climatic changes. Consequently, the period between the latest Early and Early Middle Pleistocene (~1.2-0.5 Ma) is often referred to as the Mid-Pleistocene Revolution, or Early–Middle Pleistocene Transition (EMPT). The EMPT witnessed a climatic change 748 of paramount importance: the increase in the amplitude of glacial-interglacial oscillations, a shift that 749 in turn prompted sharper recurrent changes in associated phenomena, such as, for instance, ice volume 750 cycles (Clark et al., 2006; Head and Gibbard, 2015; Maslin and Brierley, 2015). The mutated (and 751 more instable, at a ~100 kyr temporal scale) climate conditions favoured the spread of adaptable 752 species, which eventually completely replaced previous Villafranchian mammals by the end of the EMPT (Gliozzi et al., 1997). These animals include several forms closely related to extant lineages 753 754 or species, for instance, red deer, wild boars, hyenas, and wolves (van der Made et al., 2017; Iannucci 755 et al., 2021a; Iurino et al., 2022). The general pattern is quite clear, but these bioevents did not all occur at once, as the different species reacted variously to environmental changes. As emphasised by 756 757 several authors (Martinez-Navarro et al., 2010), investigating the faunal response and the timing of 758 bioevents can provide relevant clues on potential coincidental diffusions of hominin populations or lithic tradition. 759

760 The faunal list of Atella should still be considered to be provisional, as the ongoing excavations 761 will yield new material that will contribute to refining current attributions and possibly identify new species. However, some considerations can be noted. The straight-tusked elephant, Palaeoloxodon 762 antiquus, is the best-represented species at Atella. An isolated molar recovered from the latest Early 763 764 Pleistocene of Slivia, in north-eastern Italy (~0.9–0.8 Ma) is the earliest record of this elephant in Europe (Bon et al., 1992), which becomes much more abundant during the Middle Pleistocene 765 766 (Palombo and Ferretti, 2005). In Italy, the latest well-dated occurrences of the species are not recorded after MIS 5 (Braun and Palombo, 2021; Mecozzi et al., 2021b; Pieruccini et al., 2022). The straight-767 768 tusked elephant is more frequently reported during interglacial stages, but the species was a mixed 769 feeder and inhabited different environments (Palombo et al., 2005; Rivals et al., 2012).

The red deer, *Cervus elaphus*, is also documented in some latest Early Pleistocene localities, including Atapuerca, Dorn-Dürkheim, and Slivia (Bon et al., 1992; Franzen et al., 2000; van der Made et al., 2017), while it becomes widespread in Middle Pleistocene assemblages (Di Stefano et al., 2015; van der Made et al., 2017). Again, *C. elaphus* is an adaptable species with a wide ecological tolerance and flexible feeding behaviour (Gebert and Verheyden-Tixier, 2001; Benvenuti et al., 2017).

775 Few remains attest to the presence of the fallow deer, Dama sp., at Atella. The taxonomy of Dama-776 like deer is far from unanimously agreed upon (e.g., Di Stefano and Petronio, 2002; Breda and Lister, 2013; van der Made et al., 2017; Croitor, 2018), and thus it is somewhat contentious to attempt to 777 778 place occurrences of the genus in a chronological context. Differences in Middle Pleistocene fallow deer are mainly documented on antlers, and when these are not available, specific attributions are 779 often based on chronological grounds. In any case, fallow deer are common during the Middle 780 781 Pleistocene, but are usually reported in localities related to interglacial stages (Mecozzi et al., 2021c). 782 Moreover, although the diet of extant fallow deer populations varies in many respects (Esattore et al., 783 2022), a browsing behaviour is generally inferred for Middle Pleistocene fallow deer (Rivals and 784 Ziegler, 2018; Strani et al., 2022).

The Atella bovids cannot be confidently identified at a species level (contra Zucchelli, 1999), even 785 786 though some bones show Bison-like features. Nonetheless, this group arouses much interest within the context of Acheulean diffusion in Europe, with Martínez-Navarro et al. (2010) emphasising a 787 788 coincidental spread of the Acheulean culture and Bos. This was supported in particular by the 789 attributions of the material of Notarchirico and Loreto to *B. primigenius* (Caloi and Palombo, 1979; 790 Cassoli et al., 1999). However, subsequent research has pushed back the chronology of Early 791 Acheulean diffusion into Europe (see section 5.2.1.), while the taxonomic identification of the scant 792 postcranial bovid material recovered from the aforementioned localities of the Venosa Basin requires 793 further investigation (Iannucci et al., 2021b). Incontestable B. primigenius cranial remains are only 794 documented since ~ 0.5 Ma. As in the case of the other species, the representativeness of B. 795 primigenius in the fossil record increased at a later stage, during the late Middle to Late Pleistocene 796 (Wright, 2013; Iannucci et al., 2021b).

To sum up, the species encountered at Atella represent a typical assemblage for the Middle Pleistocene of Italy. Most of them are characterized by high ecological tolerance, but some indications (abundance of *P. antiquus*, presence of *Dama* sp., absence of species mainly related to cold and open 800 habitats) suggest a warm, perhaps interglacial, environment.

801 *5.2.3. Environmental evolution*

In Central and Southern Italy, the 650-500 cal. BP time interval, encompassing the site of Cimitero di Atella, is characterized by major vegetation turnover, revealed by a number of pollen records spanning at least part of this interval, stretching from MIS 16 to the first part of MIS 13.

MIS 16 is recorded at Vallo di Diano, the largest tectonic basin of the Southern Apennines, located in the Campania region, ca. 50 km south of Cimitero di Atella (Russo Ermolli, 1994; Russo Ermolli and Cheddadi, 1997). The pollen record reveals a semi-open landscape with steppe vegetation dominated by Poaceae, *Artemisia* and other Asteraceae species. Conifers were mostly represented by *Pinus* and *Abies*, especially at the transition with MIS 15, when steppe taxa declined. Apart from oaks, broadleaved trees were represented by scattered occurrences. The contents of pollen sample A01 in the Borzatti von Löwenstern sequence match the final part of MIS 16 at Vallo di Diano.

812 MIS15 is recorded at Vallo di Diano and in the Sessano Basin (Russo Ermolli et al., 2010), a large 813 tectonic depression located in Molise. The pollen record of this interglacial period points to a 814 remarkable development of broadleaved woodlands dominated by deciduous oaks with Ulmus, 815 Zelkova, Carpinus, and Acer, accompanied by sparse evergreen formations composed of evergreen 816 oaks and Ericaceae. The permanence of conifers in the landscape is revealed by significant 817 frequencies of Abies, suggesting cool and wet conditions, and Pinus. Some steppe vegetation was 818 also present in this environment, despite increased humidity. Overall, the vegetation features 819 documented by samples A02, A03 and I of the Borzatti von Löwenstern sequence are similar to some 820 MIS15 pollen spectra from Vallo di Diano. A remarkable ecological and biostratigraphic aspect of the 821 Borzatti von Löwenstern sequence is the occurrence of *Tsuga* (<3%) in samples A01 and A02. In Central Italy, the last significant abundance (>5%) of this conifer occurs in the Torre Mucchia record, 822 823 correlated with MIS 17 (Pieruccini et al., 2016; Magri et al., 2017). It probably persisted until MIS 824 13, when it occurred at Sessano (Russo Ermolli et al., 2010). Thus, the presence of *Tsuga* at Atella is consistent with the new chronology of the site and shows that sparse populations of this tree were still 825
826 present in the Italian Peninsula during the Middle Pleistocene.

827 MIS 14 is well represented in the pollen records of Vallo di Diano, Sessano, and Acerno. The latter 828 site is a narrow tectonic depression located in the central sector of the Picentini Mts, ca 50 km west 829 of Cimitero di Atella (Munno et al., 2001). The vegetation history of this period points to a new 830 decline of forests, in particular broadleaved trees and shrubs, alongside a marked expansion of steppe vegetation, but with clear floristic differences in the three sites. At Vallo di Diano, a progressive 831 832 decline in deciduous oaks is coeval with an abrupt conifer turnover, with a clear reduction of Abies 833 and a new expansion of Pinus. At Sessano, deciduous oaks declined more markedly, accompanied by 834 the disappearance of almost all the broadleaved trees and Abies. At Acerno, broadleaved trees were 835 still abundant, at some point with *Betula*. In general, climate conditions during glacial phases MIS 16 and MIS 14 do not seem sufficiently harsh to wipe out temperate forests, and plants and animals may 836 have persisted for some time in glacial refuges in Southern Italy (Orain et al., 2013). 837

838 The vegetation history of the MIS 13 interglacial phase has been reconstructed from several sites in Central and Southern Italy, including (from north to south) Rignano Flaminio (Di Rita and Sottili, 839 840 2019) and Ceprano from Lazio (Margari et al., 2018), Sessano (Russo Ermolli et al., 2010) and Boiano from Molise (Orain et al., 2015), Acerno (Munno et al., 2001) and Vallo di Diano (Russo Ermolli et 841 842 al., 1994, Petrosino et al., 2014a) from Campania, and the Mercure Basin at the Calabria-Basilicata boundary (Petrosino et al., 2014b). These sequences suggest that local environmental factors 843 844 (including edaphic, topographic and climatic conditions) led to the development of specific forest landscape characters in each site, which are difficult to place in regional vegetation patterns (Russo 845 846 Ermolli et al., 2015). In the pollen records closest to Cimitero di Atella, namely Vallo di Diano and Acerno, forests were dominated by deciduous *Quercus* and *Abies*, with a still conspicuous presence 847 of Pinus. It is likely that these trees were also the main floristic elements of the woodlands 848 849 surrounding Atella during MIS 13.

850 6. Conclusion

851 The new stratigraphic, sedimentological and geochronological data identified tephra layers in the

33

852 sediments of the Cimitero di Atella archaeological site, interpreted as fallout deposits and pyroclastic 853 flows related to the eruptive activity of the upper portion of the Monte Vulture Supersynthem (starting at the boundary between the Rionero and Vulture-San Michele subsynthems). ⁴⁰Ar/³⁹Ar dating on the 854 855 lowest tephra layer (Atella INF) of the measured stratigraphic section dated the base of the Vulture – 856 San Michele Subsynthem to 655 ± 12 ka. Sedimentological analysis also correlates the **Bmli** and **Suli** facies (archaeological unit F of Borzatti von Löwenstern et al., 1997) with the draining of the lake. 857 According to our ⁴⁰Ar/³⁹Ar dating of the F horizon of Atella, this dramatic event should have taken 858 859 place at about 570-580 ka. The erosional surface at the top of these sediments indicates a 860 morphological development of an alluvial channel, subsequently filled by 2.5-m-thick volcaniclastic 861 braided-type alluvial facies with intercalations of primary volcanic layers. These sedimentological 862 data suggest a volcanic reactivation of Monte Vulture Volcano, leading to an increase in the sedimentary load in an alluvial environment. The presence of seismically-induced soft-sediment 863 864 deformation structures can be related to eruptions with sufficient energy to quake the Atella territory. Geochronological analyses carried out in the alluvial part of the stratigraphic section suggest 865 sedimentation between 509 ± 46 ka and 570-580 ka. These data do not allow us to accurately identify 866 the upper limit of the Vulture - San Michele Subsynthem nor to link the deposits from the upper 867 portion of the Cimitero di Atella section to the younger eruptive stages of the Monte Vulture 868 869 Supersynthem. Therefore, in order to date the archaeological site more accurately, and also in relation 870 to the stratigraphic scheme of Monte Vulture Volcano by Giannandrea et al. (2006), in the future we 871 aim to collect further sedimentological, geochemical, and geochronological data from the portion 872 revealed by ongoing excavations of the site and to perform stratigraphic checks in the Monte Vulture 873 volcanic sequence.

Archaeological investigations identify at least four occupation episodes at the site during the second phase of the Lower Palaeolithic. They are different in terms of preservation and probably accumulation duration, but these occupations belong to the same techno-cultural context, and are consistent with other more or less contemporaneous sites in Italy and Western Europe. The tool kit is 878 obtained in three main ways: small tool confection, flake debitage and LCT shaping. The specific 879 productional and functional attributes of these categories need to be clearly defined in the future, along with discussions of the definition of these typical Early Acheulean productions. The still 880 881 preliminary results from the techno-functional analysis of small tools reveal the technical complexity 882 of this phase of the Lower Palaeolithic. The nature of interactions between human groups and fauna will also be developed, in particular after the extensive excavation of level F-Deb1. the most 883 884 promising in terms of preservation and deposition duration. The faunal elements identified so far 885 constitute a typical assemblage for the Middle Pleistocene of Italy, with abundant straight-tusked 886 elephants Palaeoloxodon antiquus, bovids and cervids. Palaeoloxodon antiquus and fallow deer 887 Dama sp. are more frequently recorded during interglacial stages, and Middle Pleistocene fallow deer 888 are usually associated with a browsing feeding behaviour. The identification of an elephant footprint 889 at the top of tephra I by Borzatti von Löwenstern will need to be confirmed by fieldwork. Last year, 890 new work focused on the ichnological sources, of animals but also humans (Altamura et al., 2020; 891 Duveau et al., 2019; Mayoral et al., 2021). The development of this approach, based on more samples 892 and new technical methods, will also reconsider the Atella trace found in the 1990s and explicitly 893 look for other possible footprints. The study of the fauna will focus on faunal procurement but also 894 on carcass exploitation for alimentary or technical purposes, as in other contemporaneous contexts (Boschian and Sacca, 2015; van Kolfschoten et al., 2016; Tourkoulis et al., 2018; Aranguren et al., 895 896 2019; Villa et al., 2021). Future investigations at Atella, in the field and on the material will assess 897 the degree of preservation of each occupation and better define the technical system. Then these 898 results need to be compared with results from other sites in Italy and in Europe also under study, and 899 placed in a broader context.

900 Acknowledgment:

We would like to thank the institutions that support the Atella project. The Soprintendenza
Archeologia Belle Arti e Paesaggio della Basilicata, the French school of Rome, the Basilicata region
and the municipal administration of Atella.

904

- 905
- 906 Table:
- 907Table 1: Description of Monte Vulture Volcano stratigraphic units (after Giannandrea et al., 2004, 2006; Stoppa et al., 2008). Unit
codes correspond to those of the official 1:50,000-scale geological sheet (Foglio 451 Melfi, Carta Geologica d'Italia, 2010).
- 909 Table 2: List of pyroclastic and epiclastic lithofacies with brief descriptions and interpretations
- 910 Table 3: ESR data and ages obtained for Atella samples.
- 911 Table 4: Number of archaeological finds

912 Table 5: Distribution of lithic artefacts by technological category. Retouch flakes: small retouch flakes, notch flakes, small flakes 913 of from the confection of small tools; Small tools: entire and fragmented small tools; Cores: entire cores and core fragments; Flakes: 914 production flakes from a flaking reduction sequence; Large tools: Handaxes and large shaped tools; Indet: debris, indeterminate small 915 fragments, chunks

- 916 Table 6: Size of faunal specimens in the whole site
- 917 Table 7: Alteration of fragments
- 918 Table 8: Taxonomy of faunal remains according to stratigraphy
- 919 Figures:

920 Figure 1: Map of the region. A. Atella and Venosa Basin in Southern Italy; B. Picture of the Atella Basin (archivio scavo Borzatti 921 von Löwenstern).

- 922 Figure 2: a. Geological sketch map of Southern Italy; b. DEM-based sketch map of Monte Vulture volcano and Atella and Venosa 923 mid-Pleistocene basins, with location of the Cimitero di Atella site.
- 924 Figure 3: a. Detailed geological map of the Atella area; b. geological cross-section through the Cimitero di Atella site.
- 925 Figure 4: (a), position of the cores (b) Cimitero di Atella excavation plan with the location of the different excavation areas from 1990 to 2021.
- 927 Figure 5: Panoramic view of the study area with Monte Vulture volcano in the background showing the Atella sub-horizontal landsurface.

929 Figure 6: Sedimentary log with indications of lithofacies and facies associations, correlation with previously published units (Borzatti von Löwenstern et al., 1997; Giannandrea et al., 2006), and 40Ar/39Ar and ESR dating samples.

Figure 7: Photos showing some Cimitero di Atella site lithofacies: a) thinly laminated epiclastic volcanic siltstone, Sh; b) fine- to coarse-grained massive tuff, TLm (black triangles indicate co-current falls); c) fine- to medium-grained normal graded tuff, Tng; d) three beds made of fine-grained structureless (the lowermost bed) and poorly laminated (the upper two beds) tuff, Tm; e) epiclastic volcanic breccia (lithofacies Bmli) sandwiched abruptly between two reworked colluvium deposits, Suli (black triangles indicate the basal erosive surface; white triangles indicate the upper boundary of the Bmli bed consisting of an articulated erosive surface); at the top, a tusk of Palaeoloxodon antiquus unconformably covers the Bmli and Suli lithofacies. Details of lithofacies codes in Table 2.

Figure 8: Photos showing some lithofacies from the upper part of the Cimitero di Atella site: a) fine-grained lacustrine beds of lithofacies Sh, Dh, and Su, under the lithofacies Bmli, coarse-grained deposits (triangles indicate soft-sediment deformation structures: black triangle indicates dome-like flame structure, and yellow triangle indicates simple load casts); b) alluvial fine (Sr)- to coarse (SBt)-grained epiclastic sandstone abruptly overlain by a massive pyroclastic deposit of lithofacies LTm; c) pyroclastic bed of lithofacies TLBm, showing at the base dome-like flame structure (white triangles); d) from the bottom: bed of pyroclastic deposit of lithofacies TLm, sandwiched between alluvial sediments made of a mixture of pyroclastic (Lm) and epiclastic (SBt) lithofacies (yellow triangles indicate simple load cast structures; white triangles indicate the erosive surface at the top of the lithofacies Lm; black triangles indicate the depositional surface of the lithofacies Lm above the alluvial sediment SBt); in the upper portion, irregular balland-pillow morphology (red triangles indicate boudinage-like breccia structures). Details of lithofacies codes in Table 2.

- Figure 9: Depositional architecture of alluvial facies association outcropping in the upper portion of the Cimitero di Atella
 succession (yellow triangles indicate load structures; red triangles indicate irregular ball-and-pillow structures); lithofacies codes
 reported in Table 2.
- 949 Figure 10: ⁴⁰Ar/³⁹Ar and ESR results obtained on Atella samples, presented as probability diagrams. Ages are calculated at 95% of confidence.
- 951 *Figure 11: Correlation between the log of Borzatti von Löwenstern (et al., 1997), the new log (Giannandrea this work) and the archaeological levels.*

Figure 12: Flakes from Cimitero di Atella unit F (a): flakes from the small tool reduction sequence. (1), (2): flint confection flakes;
(3): flint resharpening flakes; (4), (5): flint notch flakes. (b) Flakes from the debitage reduction sequence. (6): coarse flint flake; (7) limestone flake proximal fragment; (8), (9) small flint flakes.

956 957 958 959 960 961 962 963 964 Figure 13: Diacritical analysis of a small tool, piece n° 892, F-Deb 1, dimension 47x38x12 mm. (a): Order and direction of removals; (b): Schematic description of the phases. The piece is a coarse flint flake, with a fresh overall physical state and a slight polish. The presence of partial cortex and the series of first removals (1 to 6) indicate that the flake (7) comes from the initial phases of core reduction with partial management of convexities. Phase I: 1 to 6, series of removals showing opposite and orthogonal direction with respect to the future flake (7) decorticating the block and managing the distal and lateral convexities of the debitage surface. Some of these first removals, all smaller and thinner than the future flake (7), such as, for example removal (6), can have functional lateral cutting edges. 7, detachment of thick flake, blank of future tool. Phase II: 8, 9: Orthogonal blow fracture (8) starting from lower face and adjustment (9) starting from upper face. These removals create the back of the future tool. 10 to 16: creation of TFU1 composed of trihedron and adjacent cutting edge. Two notch blows (10, 11) starting from lower face created trihedral functional part and series 965 of small alternate retouch (12 to 16) to create a saw cutting edge. Phase III: 17 to 20: Series of scaly blows starting from upper face 966 967 to create the flat surface of future TFU2. 21 to 26: Creation of TFU2 composed of trihedron. Series of small blows (21, 24, 26) push back the edge of flake, while three notch blows (22, 23, 26) create the trihedral functional part. All these blows start from the upper 968 face. Phase IV: 27 to 31: Creation of TFU3 composed of a trihedron.

Figure 14: Techno-functional analysis of a small tool, n° 892, F-Deb 1. (a) Prehensile part (TFU p in green) composed of a flat/convex thick volume and lateral back prepared by a facture blow; transformative part (TFU t in red) made up of a trihedron and cutting-edge. (b) Prehensile part (TFU p in green) composed of a flat/convex thick volume and lateral back prepared by a facture blow; transformative part (TFU p in green) composed of a flat/convex thick volume and lateral back prepared by a facture blow;
transformative part (TFU t in red) formed by a trihedron. (c) Prehensile part (TFU p in green) composed of a flat/convex thick volume and lateral back prepared by a facture blow; transformative part (TFU t in red) formed by a trihedron.

Figure 15: Large shaped tools from unit F. Due to the very poor state of preservation, it was not possible to record the angles and
features of the section. (a) Shaped tool on a limestone nodule. Diacritical analysis: first phase (in grey, 1 to 10): removals linked to
volumetric construction; second phase (in green, 11 to 19): series of removals to complete the prehensile part; third phase (in red, 20
to 25): series of removals to implement the transformative part. Techno-functional analysis: prehensile part (TFUp in green) consisting
of a natural thick volume bordered by a natural back and a prepared back; two transformative parts (TFUt in red): TFUt 1 tip-edge
and TFUt 2 a serrated edge. (b) Shaped tool on a limestone pebble. Diacritical analysis: first phase (in grey, 1 to 16) removals linked
to the volumetric construction; second phase (in green, 17 to 21) series of removals to complete the prehensile part; third phase (in
gent and TFUt 2 to 29), series of removals to implement the transformative part. Techno-functional analysis: prehensile part; third phase (in
gent and TFUt 2 to 29), series of removals to implement the transformative part. Techno-functional analysis: prehensile part; third phase (in
gent and TFUt 2 to 29), series of removals to implement the transformative part. Techno-functional analysis: prehensile part (TFUp in green)
consisting of a thick natural volume and a prepared lateral back; two transformative parts (TFUt in red): TFUt 1 rostrum and TFUt 2
gent and gent analysis: prehensile part (TFUp in green)
gent and transformative part. Techno-functional analysis: prehensile part (TFUp in green)
gent and the gent and gent and gent and gent and gent analysis: prehensile part (TFUp in green)

984 References

- Abruzzese, C., 2014. Il sito paleolitico di Atella (PZ) nel quadro del primo popolamento europeo:
- 986 studio tecnologico dell'industria litica (Tesi di Laurea Magistrale). Università di Napoli
- 987 "L'Orientale," Naples.
- 988 Abruzzese, C., Aureli, D., Rocca, R., 2016. Assessment of the Acheulean in Southern Italy: New
- study on the Atella site (Basilicata, Italy). Quaternary International, The first peopling of Europe
- and technological change during the Lower-Middle Pleistocene transition 393, 158–168.
- 991 <u>https://doi.org/10.1016/j.quaint.2015.06.005</u>
- 992 Altamura, F., Bennett, M.R., Marchetti, L., Melis, R.T., Reynolds, S.C., Mussi, M., 2020.
- 993 Ichnological and archaeological evidence from Gombore II OAM, Melka Kunture, Ethiopia: An
- integrated approach to reconstruct local environments and biological presences between 1.2 and
- 995 0.85 Ma. Quaternary Science Reviews 244, 106506.
- 996 https://doi.org/10.1016/j.quascirev.2020.106506

- Anketell, J.M., Cegla, J., Dzulynski, S., 1969. Unconformable surfaces formed in the absence of
 current erosion. Geologica Romana 8, 41–46.
- 999 Aranguren, B., Grimaldi, S., Benvenuti, M., Capalbo, C., Cavanna, F., Cavulli, F., Ciani, F.,
- 1000 Comencini, G., Giuliani, C., Grandinetti, G., Mariotti Lippi, M., Masini, F., Mazza, P.P.A.,
- 1001 Pallecchi, P., Santaniello, F., Savorelli, A., Revedin, A., 2019. Poggetti Vecchi (Tuscany, Italy): A
- 1002 late Middle Pleistocene case of human–elephant interaction. Journal of Human Evolution 133, 32–
- 1003 60. https://doi.org/10.1016/j.jhevol.2019.05.013
- Arzarello, M., Pavia, G., Peretto, C., Petronio, C., Sardella, R., 2012. Evidence of an Early Pleistocene
 hominin presence at Pirro Nord (Apricena, Foggia, southern Italy): P13 site. Quaternary
 International, The genus Homo from Africa to Europe: evolution of terrestrial ecosystems and
 dispersal routes 267, 56–61. https://doi.org/10.1016/j.quaint.2011.01.042
- 1008 Aureli, D., Contardi, A., Giaccio, B., Jicha, B., Lemorini, C., Madonna, S., Magri, D., Marano, F.,
- 1009 Milli, S., Modesti, V., Palombo, M.R., Rocca, R., 2015. Palaeoloxodon and Human Interaction:
- 1010 Depositional Setting, Chronology and Archaeology at the Middle Pleistocene Ficoncella Site
- 1011 (Tarquinia, Italy). PLOS ONE 10, e0124498. https://doi.org/10.1371/journal.pone.0124498
- 1012 Aureli, D., Rocca, R., Lemorini, C., Modesti, V., Scaramucci, S., Milli, S., Giaccio, B., Marano, F.,
- 1013 Palombo, M.R., Contardi, A., 2016. Mode 1 or mode 2? "Small tools" in the technical variability
- 1014 of the European Lower Palaeolithic: The site of Ficoncella (Tarquinia, Lazio, central Italy).
- 1015 Quaternary International, The first peopling of Europe and technological change during the Lower-
- 1016 Middle Pleistocene transition 393, 169–184. https://doi.org/10.1016/j.quaint.2015.07.055Barral,
- L., Simone, S., 1983. Le bassin fluvio-lacustre de Venosa (Basilicate, Italie). Bull. Mus. anthropol.
 préhist. Monaco 5–19.
- 1019 Benvenuti, M., Bahain, J.-J., Capalbo, C., Capretti, C., Ciani, F., D'Amico, C., Esu, D., Giachi, Gi.,
- 1020 Giuliani, C., Gliozzi, E., Lazzeri, S., Macchioni, N., Lippi, M.M., Masini, F., Mazza, P.P.A.,
- 1021 Pallecchi, P., Revedin, A., Savorelli, A., Spadi, M., Sozzi, L., Vietti, A., Voltaggio, M., Aranguren,
- 1022 B., 2017. Paleoenvironmental context of the early Neanderthals of Poggetti Vecchi for the late

- 1023 middle Pleistocene of Central Italy. Quaternary Research 88, 327-344.
- 1024 https://doi.org/10.1017/qua.2017.51
- Blanc, A.C., 1953. Venosa, gisement à industrie tayacienne et micoquienne de Loreto. IV Congr.
 Intern. INQUA 63–68.
- 1027 Boëda, É., 2013. Techno-logique & technologie, @rchéo-éditions. ed. Prigonrieux.
- Boëda, É., Geneste, J.-M., Meignen, L., 1990. Identification de chaînes opératoires lithiques du
 Paléolithique ancien et moyen. Paléo 2, 43–80.
- 1030 Bon, M., Piccoli, G., Sala, B., 1992. La fauna pleistocenica della breccia di Slivia (Carso triestino)
- 1031 nella collezione del Museo Civico di Storia Naturale di Trieste. Atti Mus. Civico Storia Nat.
 1032 Trieste 44, 33–51.
- 1033 Borzatti von Löwenstern, E., 1985. Sulle rive di un lago scomparso. L'Universo, Firenze.

1034

- Borzatti von Löwenstern, E., 1998. Il Bacino di Atella nella Preistoria. Studi per l'ecologia del
 Quaternario 7–39.
- Borzatti von Löwenstern, E., 2005. Il sito Acheuleano antico del Cimitero di Atella: gli strati alti del
 sedimento preistorico. Studi per l'ecologia del Quaternario 7–20.
- Borzatti von Löwenstern, E., 2011. L'antico lago di Atella: verso la fine di un ambiante geografico.
 Studi per l'ecologia del Quaternario 57.
- Borzatti von Löwenstern, E., Sozzi, M., 1994. Prime ipotesi paleogeografiche sul sito acheuleano del
 Cimitero di Atella. Studi per l'ecologia del Quaternario 16.
- Borzatti von Löwenstern, E., Sozzi, M., 2001. Il Bacino di Atella, 10.000 strumenti di pietra. Libria,
 Melfi.
- Borzatti von Löwenstern, E., Vianello, F., 1990. Nuove ricerche a Masseria Palladino nel Bacino di
 Atella (Filiano-PZ). Studi per l'ecologia del Quaternario 12, 31.
- Borzatti von Löwenstern, E., Vianello, F., 1989. L'Acheuleano antico di Masseria Palladino nel
 Bacino di Atella (Potenza). Studi per l'ecologia del Quaternario 11, 9.
- 1049 Borzatti von Löwenstern, E., Sozzi, M., Vannucci, S., Vianello, F., 1990. L'acheuleano del cimitero

- di Atella. Prime indagini sulla stratigrafia del sedimento e sulle industrie litiche. Studi per
 l'ecologia del Quaternario 12, 9–30.
- 1052 Borzatti von Löwenstern, E., Palchetti, A., Sozzi, M., Maestrini, M., 1997. Témoignages de
- 1053 l'Acheuléen inférieur en Italie méridionale : Le gisement du Cimitero di Atella (Basilicata).
 1054 L'Anthropologie 101, 617–638.
- Borzatti von Löwenstern, E., Fabiano, M., Secci, M.M., Sozzi, M., 1998. Paleoenvironmental
 investigations on the Acheulean site at the Cimitero di Atella (Potenza, Southern Italy). Atti del
 XIII Congr. UISPP, Forlì.
- 1058 Boschian, G., Saccà, D., 2015. In the elephant, everything is good: Carcass use and re-use at Castel
- 1059 di Guido (Italy). Quaternary International, The Origins of Recycling: A Paleolithic Perspective
- 1060 361, 288–296. https://doi.org/10.1016/j.quaint.2014.04.030
- 1061 Boschin, F., Rocca, R., Aureli, D., 2018. New archaeozoological and taphonomic analysis on macro-
- and megafauna remains from the Lower Palaeolithic site of Ficoncella (Tarquinia, Central Italy).
- 1063 Quaternaire. Revue de l'Association française pour l'étude du Quaternaire 13–20.
- Breda, M., Lister, A.M., 2013. Dama roberti, a new species of deer from the early Middle Pleistocene
- 1065 of Europe, and the origins of modern fallow deer. Quaternary Science Reviews 69, 155–167.
- 1066 <u>https://doi.org/10.1016/j.quascirev.2013.01.029</u>
- Brennan, B, Lyons, R, Phillips, S. 1991. Attenuation of alpha particle track dose for spherical grains.
 Nuclear Tracks Radiation Measurements 18, 249-253.
- 1069 Brennan, B. 2003. Beta doses to spherical grains. Radiation Measurements 37, 299-303.
- Burdukiewicz, J.M., Ronen, A., 2003. Lower Palaeolithic Small Tools in Europe and the Levant.
 British Archaeological Reports, Oxford.
- 1072 Caloi L., Palombo M.R., 1979. La fauna quaternaria di Venosa: bovidi. Boll. Serv. Geol. Ital. 100,
 1073 101–140.
- 1074 Cassoli, F., Di Stefano, G., Tagliacozzo, A., 1999. I Vertebrati dei livelli superiori (Alfa ed A) della
- 1075 serie stratigrafica di Notarchirico, in: Piperno, M. (Ed.), Notarchirico. Un Sito Del Pleistocene

- 1076 Medio-Antico Nel Bacino Di Venosa (Basilicata). Osana edizione, Venosa, pp. 361–438.
- 1077 Ciolli, N., 1997. *Elephas antiquus* Falconer & Cautley del Cimitero di Atella (Pz). Studi per
 1078 l'ecologia del Quaternario 25–34.
- 1079 Croitor, R., 2018. Plio-Pleistocene deer of western palearctic: taxonomy, systematics, phylogeny.
 1080 Institute of Zoology of the Academy of Sciences of Moldova, Chisinău.
- 1081 Clark, P.U., Archer, D., Pollard, D., Blum, J.D., Rial, J.A., Brovkin, V., Mix, A.C., Pisias, N.G., Roy,
- 1082 M., 2006. The middle Pleistocene transition: characteristics, mechanisms, and implications for
- 1083 long-term changes in atmospheric pCO2. Quaternary Science Reviews, Critical Quaternary
 1084 Stratigraphy 25, 3150–3184. https://doi.org/10.1016/j.quascirev.2006.07.008
- 1085 Crovetto, C., 1993. Le Paléolithique inférieur de Loreto (Venosa, Basilicate, Italie). Bulletin du
- 1086 Musée d'anthropologie préhistorique de Monaco 31–57.
- 1087 Crovetto, C., Ferrari, M., Peretto, C., Longo, L., Vianello, F., 1994. The carinated denticulates from 1088 the Paleolithic site of Isernia La Pineta (Molise, Central Italy): Tools or flaking waste? The results
- 1089 of the 1993 lithic experiments. Human Evolution 9, 175–207.
- 1090 Weyer, L.D., Pérez, A., Hoguin, R., Forestier, H., Boëda, E., 2022. Time, memory and alterity in
- 1091 prehistoric lithic technology: Synthesis and perspectives of the French technogenetic approach.
- 1092 Journal of Lithic Studies 9, 46 p.-46 p. https://doi.org/10.2218/jls.7020
- 1093 Despriée, J., Moncel, M.-H., Arzarello, M., Courcimault, G., Voinchet, P., Bahain, J.-J., Falguères,
- 1094 C., 2018. The 1-million-year-old quartz assemblage from Pont-de-Lavaud (Centre, France) in the
- European context. Journal of Quaternary Science 33, 639–661. https://doi.org/10.1002/jqs.3042
- 1096 Di Muro, A., 1999. Inquadramento tefrostratigrafico del sito acheuleano del Cimitero di Atella
- 1097 (Basilicata-Italia). Studi per l'ecologia del Quaternario 7–14.
- 1098 Di Stefano, G., Pandolfi, L., Petronio, C., Salari, L., 2015. The morphometry and the occurrence of
- 1099 *Cervus elaphus* (Mammalia, Cervidae) from the Late Pleistocene of the Italian peninsula. Riv. Ital.
- 1100 Paleontol. Stratigr. 121(1), 103–120.
- 1101 Di Stefano, G., Petronio, C., 2002. Systematics and evolution of the Eurasian Plio-Pleistocene tribe

- 1102 Cervini (Artiodactyla, Mammalia). Geol. Romana 36, 311–334.
- 1103 Duval, M., Guilarte, V., 2015. ESR dosimetry of optically bleached quartz grains extracted from Plio-
- quaternary sediment: evaluating some key aspects of the ESR signal associated to the Ti-center.
 Radiation Measurement 78, 28–41.
- 1106 Duveau, J., Berillon, G., Verna, C., Laisné, G., Cliquet, D., 2019. The composition of a Neandertal
- social group revealed by the hominin footprints at Le Rozel (Normandy, France). PNAS 116,
- 1108 19409–19414. https://doi.org/10.1073/pnas.1901789116
- 1109 Esattore, B., Saggiomo, L., Sensi, M., Francia, V., Cherin, M., 2022. Tell me what you eat and I'll tell
- 1110 you...where you live: an updated review of the worldwide distribution and foraging ecology of 1111 the fallow deer (Dama dama). Mamm Biol. https://doi.org/10.1007/s42991-022-00250-6
- Franzen, J.L., Gliozzi, E., Jellinek, T., Scholger, R., Weidenfeller, M., 2000. Die spätaltpleistozäne
 Fossillagerstätte Dorn-Dürkheim 3 und ihre Bedeutung für die Rekonstruktion der Entwicklung
 des rheinischen Flußsystems. Senckenbergiana lethaea 80, 305–353.
 https://doi.org/10.1007/BF03043674
- 1116 Fisher, R.V., Schmincke, H.-U., 2012. Pyroclastic Rocks. Springer.
- 1117 García-Medrano, P., Ollé, A., Mosquera, M., Cáceres, I., Díez, C., Carbonell, E., 2014. The earliest
- 1118 Acheulean technology at Atapuerca (Burgos, Spain): Oldest levels of the Galería site (GII Unit).
- 1119 Quaternary International, Environmental History of European High Mountains 353, 170–194.
- 1120 <u>https://doi.org/10.1016/j.quaint.2014.03.053</u>
- 1121 Giannandrea, P., 2009. Evoluzione sedimentaria della successione alluvionale e lacustre quaternaria
- del Bacino di Venosa (Italia meridionale). Alpine and Mediterranean Quaternary 22, 269–290.
- 1123 Giannandrea, P., La Volpe, L., Principe, C., Schiattarella, M., 2006. Unità stratigrafiche a limiti
- 1124 inconformi e storia evolutiva del vulcano medio-pleistocenico di Monte Vulture (Appennino
- 1125 meridionale, Italia). Bollettino-Società Geologica Italiana 125, 67.
- 1126 Giannandrea, P., Marino, M., Romeo, M., Schiattarella, M., 2014. Pliocene to Quaternary evolution
- 1127 of the Ofanto Basin in southern Italy: an approach based on the unconformity-bounded

- 1128 stratigraphic units. IJG 133, 27–44. <u>https://doi.org/10.3301/IJG.2013.11</u>
- 1129 Gebert, C., Verheyden-Tixier, H., 2001. Variations of diet composition of Red Deer (Cervus elaphus
- 1130 L.) in Europe. Mammal Review 31, 189–201. https://doi.org/10.1111/j.1365-2907.2001.00090.x
- 1131 Gibert, L., Alfaro, P., García-Tortosa, F.J., Scott, G., 2011. Superposed deformed beds produced by
- 1132 single earthquakes (Tecopa Basin, California): Insights into paleoseismology. Sedimentary
- 1133 Geology, Recognising triggers for soft-sediment deformation: Current understanding and future
- 1134 directions 235, 148–159. <u>https://doi.org/10.1016/j.sedgeo.2010.08.003</u>
- 1135 Gliozzi, E., Abbazzi, L., Argenti, P., Azzaroli, A., Caloi, L., Barbato, L.C., Stefano, G.D., Esu, D.,
- 1136 Ficcarelli, G., Girotti, O., Kotsakis, T., Masini, F., Mazza, P., Mezzabotta, C., Palombo, M.R.,
- 1137 Petronio, C., Rook, L., Sala, B., Sardella, R., Zanalda, E., Torre, D., 1997. Biochronology of
- selected mammals, molluscs and ostracods from the middle Pliocene to the late Pleistocene in
- 1139 Italy. The state of the art. Rivista Italiana di Paleontologia e Stratigrafia 103.
 1140 https://doi.org/10.13130/2039-4942/5299
- Grifoni, R., Tozzi, C., 2006. L'émergence des identités culturelles au Paléolithique inférieur : le cas
 de l'Italie. Comptes Rendus Palevol, 5, 137–148.
- 1143 Grimaldi, S., Santaniello, F., Angelucci, D.E., Bruni, L., Parenti, F., 2020. A Techno-Functional
- 1144 Interpretation of the Lithic Assemblage from Fontana Ranuccio (Anagni, Central Italy): an Insight
- 1145 into a MIS 11 Human Behaviour. J Paleo Arch. https://doi.org/10.1007/s41982-020-00064-3
- Grün, R., 1994. A cautionary note: use of the "water content" and "depth for cosmic ray dose rate" in
 AGE and DATA. Ancient TL 12, 50- 51.
- Guérin, G., Mercier, N., Adamiec, G., 2011. Dose-rate conversion factors: update. Ancient TL, 29, 58.
- 1150 Guillou H, Scao V., Nomade S., 2018. 40Ar/39Ar age of cryptochron C2r.2r-1 as recorded in a lava
- sequence within the Ko'olau volcano (Hawaii, USA). Quaternary Geochronology 43,91–101. doi
- 1152 : 10.1016/j.quageo.2017.10.005
- 1153 Head, M.J., Gibbard, P.L., 2015. Early–Middle Pleistocene transitions: Linking terrestrial and marine

- realms. Quaternary International, The Jaramillo Subchron and the Early-Middle Pleistocene
 transition in continental records from a multidisciplinary perspective 389, 7–46.
 <u>https://doi.org/10.1016/j.quaint.2015.09.042</u>
- Iannucci, A., Mecozzi, B., Sardella, R., Iurino, D.A., 2021a. The extinction of the giant hyena
 Pachycrocuta brevirostris and a reappraisal of the Epivillafranchian and Galerian Hyaenidae in
- 1159 Europe: Faunal turnover during the Early–Middle Pleistocene Transition. Quaternary Science
- 1160 Reviews 272, 107240. <u>https://doi.org/10.1016/j.quascirev.2021.107240</u>
- 1161 Iannucci, A., Mecozzi, B., Sardella, R., 2021b. Large Mammals from the Middle Pleistocene (Mis
- 1162 11) site of Fontignano 2 (Rome, Central Italy), with an overview of "San Cosimato" assemblages.
- Alpine and Mediterranean Quaternary 34, 155–164. https://doi.org/10.26382/AMQ.2021.07
- Inizan, M.-L., Reduron-Ballinger, M., Roche, H., Tixier, J., 1999. Technology and Terminology of
 Knapped Stone. Préhistoire de la Pierre Taillée. Éditions du CREP. Nanterre.
- 1166 Iurino, D.A., Mecozzi, B., Iannucci, A., Moscarella, A., Strani, F., Bona, F., Gaeta, M., Sardella, R.,

1167 2022. A Middle Pleistocene wolf from central Italy provides insights on the first occurrence of

- 1168 Canis lupus in Europe. Sci Rep 12, 2882. <u>https://doi.org/10.1038/s41598-022-06812-5</u>
- 1169 Konidaris, G.E., Athanassiou, A., Tourloukis, V., Thompson, N., Giusti, D., Panagopoulou, E.,
- 1170 Harvati, K., 2018. The skeleton of a straight-tusked elephant (Palaeoloxodon antiquus) and other
- 1171 large mammals from the Middle Pleistocene butchering locality Marathousa 1 (Megalopolis Basin,
- 1172 Greece): preliminary results. Quaternary International, The Gates of Europe 497, 65–84.
- 1173 https://doi.org/10.1016/j.quaint.2017.12.001
- 1174 Laurent, M., Falguères, C., Bahain, J-J., Rousseau, L., Van Vliet-Lanoë, B. 1998. ESR dating of
- 1175 quartz extracted from Quaternary and Neogene sediments: method, potential and actual limits.
- 1176 Quaternary Science Reviews 17, 1057-1061.
- 1177 Lepot, M., 1993. Approche techno-fonctionnelle de l'outillage lithique Moustérien : essai de
- 1178 classification des parties actives en terme d'efficacité technique (Mémoire de Maîtrise). Université
- 1179 Paris X Nanterre, Nanterre.

- 1180 Lhomme, V., 2007. Tools, space and behaviour in the Lower Palaeolithic: discoveries at Soucy in the
- 1181 Paris basin. Antiquity 81, 536–554.
- 1182 Lee, J.Y., Marti, K., Severinghaus, J.P., Kawamura, K., Hee-Soo, Y., Lee, J.B., Kim, J.S., 2006. A
- 1183 redetermination of the isotopic abundances of atmospheric Ar. Geochimica et Cosmochimica Acta
- 1184 70, 4507-4512. doi: 10.1016/j.gca.2006.06.1563
- 1185 Longo, L., Peretto, C., Sozzi, M., Vannucci, S., Leroy-Prost, C., 1997. Artefacts, outils ou supports
- 1186 épuisés ? Une nouvelle approche pour l'étude des industries du Paléolithique ancien : Le cas
- d'Isernia La Pineta (Molise, Italie centrale). L'Anthropologie 101, 579–596.
- Ludwig K. R., 2003. Isoplot 3.0, a geochronological toolkit for Microsoft Excel. Berkeley
 Geochronology Center Special Publication, 4, 71p.
- Maslin, M.A., Brierley, C.M., 2015. The role of orbital forcing in the Early Middle Pleistocene
 Transition. Ouaternary International. The Jaramillo Subchron and the Early-Middle Pleistocene
- transition in continental records from a multidisciplinary perspective 389, 47–55.
 https://doi.org/10.1016/j.quaint.2015.01.047
- Magri, D., Rita, F., 2015. Archaeopalynological Preparation Techniques.<u>https://doi.org/10.1007/978-</u>
 <u>3-319-19944-3_27</u>
- Margari, V., Roucoux, K., Magri, D., Manzi, G., Tzedakis, P.C., 2018. The MIS 13 interglacial at
 Ceprano, Italy, in the context of Middle Pleistocene vegetation changes in southern Europe.
 Quaternary Science Reviews 199, 144–158. https://doi.org/10.1016/j.quascirev.2018.09.016
- 1199 Martínez-Navarro, B., Rook, L., Papini, M., Libsekal, Y., 2010. A new species of bull from the Early
- 1200 Pleistocene paleoanthropological site of Buia (Eritrea): Parallelism on the dispersal of the genus
- 1201 Bos and the Acheulian culture. Quaternary International, Quaternary Changes of Mammalian
- 1202 Communities Across and Between Continents 212, 169–175. 1203 https://doi.org/10.1016/j.quaint.2009.09.003
- 1204 Mayoral, E., Díaz-Martínez, I., Duveau, J., Santos, A., Ramírez, A.R., Morales, J.A., Morales, L.A.,
- 1205 Díaz-Delgado, R., 2021. Tracking late Pleistocene Neandertals on the Iberian coast. Sci Rep 11,

- 1206 4103. https://doi.org/10.1038/s41598-021-83413-8
- Mecozzi, B., Iannucci, A., Sardella, R., Curci, A., Daujeard, C., Moncel, M.-H., 2021a. *Macaca* ulna
 from new excavations at the Notarchirico Acheulean site (Middle Pleistocene, Venosa, southern)
- 1209 Italy). Journal of Human Evolution 153, 102946. https://doi.org/10.1016/j.jhevol.2020.102946
- 1210 Mecozzi, B., Bellucci, L., Giustini, F., Iannucci, A., Iurino, D.A., Mazzini, I., Strani, F., Sardella, R.,
- 1211 2021b. A reappraisal of the Pleistocene mammals from the karst infilling deposits of the Maglie
- 1212 area (Lecce, Apulia, Southern Italy). Rivista Italiana di Paleontologia e Stratigrafia 127.
- 1213 https://doi.org/10.13130/2039-4942/15776
- 1214 Mecozzi, B., Iannucci, A., Mancini, M., Sardella, R., 2021c. Redefining Ponte Molle (Rome, Central
- 1215 Italy): an important locality for Middle Pleistocene mammal assemblages of Europe. Alpine and
- 1216 Mediterranean Quaternary 34, 131–154. <u>https://doi.org/10.26382/AMQ.2021.09</u>
- Mercier, N., Falguères, C., 2007. Field gamma dose-rate measurement with a NaI(Tl) detector:
 reevaluation of the "threshold" technique. Ancient TL, 25,1.
- 1219 Miall, A.D., 1978. Tectonic setting and syndepositional deformation of molasse and other nonmarine-
- 1220 paralic sedimentary basins. Can. J. Earth Sci. 15, 1613–1632. https://doi.org/10.1139/e78-166
- 1221 Michel, V., Shen, C.-C., Woodhead, J., Hu, H.-M., Wu, C.-C., Moullé, P.-É., Khatib, S., Cauche, D.,
- 1222 Moncel, M.-H., Valensi, P., Chou, Y.-M., Gallet, S., Echassoux, A., Orange, F., Lumley, H., 2017.
- 1223 New dating evidence of the early presence of hominins in Southern Europe. Scientific Reports 7,
- 1224 10074. https://doi.org/10.1038/s41598-017-10178-4
- 1225 Moncel, M., Schreve, D., 2016. The Acheulean in Europe: Origins, evolution and dispersal.
- 1226 Quaternary International, The Acheulean in Europe: origins, evolution and dispersal 411, Part B,
- 1227 1–8. https://doi.org/10.1016/j.quaint.2016.08.039
- 1228 Moncel, M.-H., Despriée, J., Voinchet, P., Tissoux, H., Moreno, D., Bahain, J.-J., Courcimault, G.,
- 1229 Falguères, C., 2013. Early Evidence of Acheulean Settlement in Northwestern Europe La Noira
- 1230 Site, a 700 000 Year-Old Occupation in the Center of France. PLOS ONE 8, e75529.
- 1231 https://doi.org/10.1371/journal.pone.0075529

- Moncel, M.-H., Ashton, N., Lamotte, A., Tuffreau, A., Cliquet, D., Despriée, J., 2015. The Early
 Acheulian of north-western Europe. Journal of Anthropological Archaeology 40, 302–331.
- 1234 Moncel, M.-H., Despriée, J., Voinchet, P., Courcimault, G., Hardy, B., Bahain, J.-J., Puaud, S., Gallet,
- 1235 X., Falguères, C., 2016. The Acheulean workshop of la Noira (France, 700 ka) in the European
- 1236technological context. Quaternary International, The first peopling of Europe and technological1237changeduringtheLower-MiddlePleistocenetransition393,112–136.
- 1238 https://doi.org/10.1016/j.quaint.2015.04.051
- 1239 Moncel, M.-H., Arzarello, M., Boëda, É., Bonilauri, S., Chevrier, B., Gaillard, C., Forestier, H.,
- 1240 Yinghua, L., Sémah, F., Zeitoun, V., 2018. The assemblages with bifacial tools in Eurasia (first
- 1241 part). What is going on in the West? Data on western and southern Europe and the Levant. Comptes
- 1242 Rendus Palevol, Hominins and tools. Expansion from Africa towards Eurasia / Homininés et outils.
- 1243 Expansions depuis l'Afrique vers l'Eurasie 17, 45–60. https://doi.org/10.1016/j.crpv.2015.09.009
- 1244 Moncel, M.-H., Santagata, C., Pereira, A., Nomade, S., Bahain, J.-J., Voinchet, P., Piperno, M., 2019.
- 1245 A biface production older than 600 ka ago at Notarchirico (Southern Italy) contribution to
- understanding early Acheulean cognition and skills in Europe. PLOS ONE 14, e0218591.
- 1247 https://doi.org/10.1371/journal.pone.0218591
- 1248 Moncel, M.-H., Santagata, C., Pereira, A., Nomade, S., Voinchet, P., Bahain, J.-J., Daujeard, C., Curci,
- 1249 A., Lemorini, C., Hardy, B., Eramo, G., Berto, C., Raynal, J.-P., Arzarello, M., Mecozzi, B.,
- 1250 Iannucci, A., Sardella, R., Allegretta, I., Delluniversità, E., Terzano, R., Dugas, P., Jouanic, G.,
- 1251 Queffelec, A., d'Andrea, A., Valentini, R., Minucci, E., Carpentiero, L., Piperno, M., 2020. The
- 1252 origin of early Acheulean expansion in Europe 700 ka ago: new findings at Notarchirico (Italy).
- 1253 Sci Rep 10, 13802. https://doi.org/10.1038/s41598-020-68617-8.
- 1254 Mosquera, M., Ollé, A., Saladié, P., Cáceres, I., Huguet, R., Rosas, A., Villalaín, J., Carrancho, A.,
- Bourlès, D., Braucher, R., Pineda, A., Vallverdú, J., 2016. The Early Acheulean technology of
- 1256 Barranc de la Boella (Catalonia, Spain). Quaternary International, The first peopling of Europe
- 1257 and technological change during the Lower-Middle Pleistocene transition 393, 95–111.

- 1258 https://doi.org/10.1016/j.quaint.2015.05.005
- Nehyba, S., Nývlt, D., 2014. Deposits of pyroclastic mass flows at Bibby Hill (Pliocene, James Ross
 Island, Antarctica). Czech Polar Reports 4, 103–122. https://doi.org/10.5817/CPR2014-2-11
- 1261 Nicoud, E., 2011. Le phénomène acheuléen en Europe occidentale : Approche chronologique,
- 1262 technologie lithique et implications culturelles (Thèse de doctorat). Université Aix Marseille 1,
- 1263 Aix-en-Provence.
- Nicoud, E., 2013. What does the Acheulean consist of? The example of Western Europe (MIS 16-9).
 Mitteilungen der Gesellschaft für Urgeschichte 22, 41–60.
- 1266 Nicoud, E., Aureli, D., Pagli, M., Villa, V., Chaussé, C., Agostini, S., Bahain, J.-J., Boschian, G.,
- 1267 Degeai, J.-P., Fusco, F., Giaccio, B., Hernandez, M., Kuzucuoglu, C., Lahaye, C., Lemorini, C.,
- 1268 Limondin-Lozouet, N., Mazza, P., Mercier, N., Nomade, S., Pereira, A., Robert, V., Rossi, M.A.,
- 1269 Virmoux, C., Zupancich, A., 2016. Preliminary data from Valle Giumentina Pleistocene site
- 1270 (Abruzzo, Central Italy): A new approach to a Clactonian and Acheulian sequence. Quaternary
- 1271 International, Special Issue: The Hoslteinian period in Europe (MIS 11-9) 409, Part B, 182–194.
- 1272 https://doi.org/10.1016/j.quaint.2015.08.080
- Niespolo, E.M., Rutte, D., Deino, A. et Renne, P.R., 2017. Intercalibration and age of the Alder Creek
 sanidine 40Ar/39Ar standard. Quaternary Geochronology in press. http://dx.doi.
 org/10.1016/j.quageo.2016.09.004
- - Nomade, S., Gauthier, A., Guillou, H., Pastre, J.F., 2010. 40Ar/39Ar temporal framework for the
 Alleret maar lacustrine sequence (French Massif-Central): Volcanological and paleoclimatic
 implications. Quaternary Geochronology 5, 20-27.
 - 1279 Ollé, A., Mosquera, M., Rodríguez, X.P., de Lombera-Hermida, A., García-Antón, M.D., García-
 - 1280 Medrano, P., Peña, L., Menéndez, L., Navazo, M., Terradillos, M., Bargalló, A., Márquez, B., Sala,
 - 1281 R., Carbonell, E., 2013. The Early and Middle Pleistocene technological record from Sierra de
 - 1282 Atapuerca (Burgos, Spain). Quaternary International, East meets West: First settlements and
 - 1283 human evolution in Eurasia 295, 138–167. https://doi.org/10.1016/j.quaint.2011.11.009

- Orain, R., Lebreton, V., Ermolli, E.R., Combourieu-Nebout, N., Sémah, A.-M., 2013. *Carya* as
 marker for tree refuges in southern Italy (Boiano basin) at the Middle Pleistocene.
 Palaeogeography, Palaeoclimatology, Palaeoecology 369, 295–302.
 https://doi.org/10.1016/j.palaeo.2012.10.037
- 1288 Orain, R., Russo Ermolli, E., Lebreton, V., Di Donato, V., Bahain, J.-J., Sémah, A.-M., 2015.
- 1289 Vegetation sensitivity to local environmental factors and global climate changes during the Middle
- Pleistocene in southern Italy—A case study from the Molise Apennines. Review of Palaeobotany
 and Palynology 220, 69–77. https://doi.org/10.1016/j.revpalbo.2015.05.002
- 1292 Palladino, G., Giannandrea, P., Siniscalchi, A., Magri, C., Loiacono, F., 2018. Quaternary
- 1293 tectonostratigraphic evolution of the Vlora Basin, south-western Albania. Geological Journal 53,
- 1294 1698–1715. https://doi.org/10.1002/gj.2992
- Palma Di Cesnola, A., 2001. Il paleolitico inferiore e medio in Italia, Millenni. Museo Fiorentino di
 Preistoria "Paolo Graziosi," Firenze.
- 1297 Palombo, M.R., Ferretti, M.P., 2005. Elephant fossil record from Italy: knowledge, problems, and
- 1298 perspectives. Quaternary International, Studying Proboscideans: knowledge, Problems and
- 1299 Perspectives. Selected papers from "The world of Elephants" Congress, Rome 126–128, 107–136.
- 1300 <u>https://doi.org/10.1016/j.quaint.2004.04.018</u>
- Palombo, M.R., Filippi, M.L., Iacumin, P., Longinelli, A., Barbieri, M., Maras, A., 2005. Coupling
 tooth microwear and stable isotope analyses for palaeodiet reconstruction: the case study of Late
 Middle Pleistocene Elephas (Palaeoloxodon) antiquus teeth from Central Italy (Rome area).
- 1304 Quaternary International, Studying Proboscideans: knowledge, Problems and Perspectives.
- 1305 Selected papers from "The world of Elephants" Congress, Rome 126–128, 153–170.
- 1306 https://doi.org/10.1016/j.quaint.2004.04.020
- 1307 Pereira, A., Nomade, S., Voinchet, P., Bahain, J.J., Falguères, C., Garon, H., Lefèvre, D., Raynal, J.P.,
- 1308 Scao, V., Piperno, M., 2015. The earliest securely dated hominin fossil in Italy and evidence of
- 1309 Acheulian occupation during glacial MIS 16 at Notarchirico (Venosa, Basilicata, Italy). Journal of

- 1310 Quaternary Science 30, 639–650.
- Peretto, C., Terzani, C., Vannucci, S., Vaselli, O., Vianello, F., 1994. Le industrie litiche del
 giacimento paleolitico di Isernia La Pineta: la tipologia, le tracce di utilizzazione, la
 sperimentazione, Cosmo Iannone edizione. ed. Campobasso.
- 1314 Peretto, C., Arzarello, M., Gallotti, R., Lembo, G., Minelli, A., Thun Hohenstein, U., 2004. Middle
- 1315 Pleistocene behavioural strategies: the contribution of Isernia La Pineta site, in: Baquedano Pérez,
- 1316 E., Rubio-Jara, S., Aguirre, E. (Eds.), Miscelánea En Homenaje a Emiliano Aguirre. Comunidad
- 1317 de Madrid, Museo Arqueológico Regional, Madrid, pp. 369–381.
- 1318 Peretto, C., Arnaud, J., Moggi-Cecchi, J., Manzi, G., Nomade, S., Pereira, A., Falguères, C., Bahain,
- 1319 J.-J., Grimaud-Hervé, D., Berto, C., Sala, B., Lembo, G., Muttillo, B., Gallotti, R., Hohenstein,
- U.T., Vaccaro, C., Coltorti, M., Arzarello, M., 2015. A Human Deciduous Tooth and New
 40Ar/39Ar Dating Results from the Middle Pleistocene Archaeological Site of Isernia La Pineta,
 Southern Italy. PLOS ONE 10, e0140091. https://doi.org/10.1371/journal.pone.0140091
- 1323 Petrosino, P., Jicha, B.R., Mazzeo, F.C., Russo Ermolli, E., 2014a. A high resolution
- tephrochronological record of MIS 14–12 in the Southern Apennines (Acerno Basin, Italy).
- Journal of Volcanology and Geothermal Research 274, 34–50.
 https://doi.org/10.1016/j.jvolgeores.2014.01.014
- Petrosino, P., Ermolli, E.R., Donato, P., Jicha, B., Robustelli, G., Sardella, R., 2014b. Using
 Tephrochronology and palynology to date the MIS 13 lacustrine sediments of the Mercure basin
 (Southern Apennines–Italy). Italian Journal of Geosciences 133, 169–186.
- 1330 Pieruccini, P., Forti, L., Mecozzi, B., Iannucci, A., Yu, T.-L., Shen, C.-C., Bona, F., Lembo, G.,
- 1331 Muttillo, B., Sardella, R., Mazzini, I., 2022. Stratigraphic reassessment of Grotta Romanelli sheds
- 1332 light on Middle-Late Pleistocene palaeoenvironments and human settling in the Mediterranean.
- 1333 Sci Rep 12, 13530. https://doi.org/10.1038/s41598-022-16906-9
- Pilleyre, T., Sanzelle, S., Fain, J., Miallier, D., Montret, M., 1999. Essai de datation par
 thermoluminescence des depots du site acheuleen de Notarchirico, in: Piperno, M. (Ed.),

- Notarchirico. Un Sito Del Pleistocene Medio-Antico Nel Bacino Di Venosa (Basilicata),. Osana
 edizione, p. 235.
- Piperno, M. (Ed.), 1999. Notarchirico: un sito del Pleistocene medio antico nel bacino di Venosa.
 Osana edizione, Venosa.
- 1340 Pope, M.C., Fred Read, J., Bambach, R., Hofmann, H.J., 1997. Late Middle to Late Ordovician
- seismites of Kentucky, southwest Ohio and Virginia: Sedimentary recorders of earthquakes in the
- 1342 Appalachian basin. Geological Society of America Bulletin 109, 489–503.
- Prescott, J.R., Hutton, J.T., 1994. Cos3ic ray contributions to dose rates for luminescence and ESR
 dating: large depths and long-term time variations. Radiation Measurements, 23, 497–500.
- 1345 Rivals, F., Semprebon, G., Lister, A., 2012. An examination of dietary diversity patterns in Pleistocene
- 1346 proboscideans (Mammuthus, Palaeoloxodon, and Mammut) from Europe and North America as
- 1347 revealed by dental microwear. Quaternary International, Mammoths and Their Relatives 1:
- 1348 Biotopes, Evolution and Human Impact V International Conference, Le Puy-en-Velay, 2010 255,
- 1349 188–195. <u>https://doi.org/10.1016/j.quaint.2011.05.036</u>
- 1350 Rivals, F., Ziegler, R., 2018. High-resolution paleoenvironmental context for human occupations
- during the Middle Pleistocene in Europe (MIS 11, Germany). Quaternary Science Reviews 188,
- 1352 136–142. https://doi.org/10.1016/j.quascirev.2018.03.026
- 1353 Rocca, R., 2016. First settlements in Central Europe: Between originality and banality. Quaternary
- 1354 International, Special Issue: The Hoslteinian period in Europe (MIS 11-9) 409, Part B, 213–221.
- 1355 https://doi.org/10.1016/j.quaint.2015.08.066
- 1356 Rocca, R., Abruzzese, C., Aureli, D., 2016. European Acheuleans: Critical perspectives from the East.
- 1357 Quaternary International, The Acheulean in Europe: origins, evolution and dispersal 411, Part B,
- 1358 402–411. https://doi.org/10.1016/j.quaint.2016.01.025
- 1359 Rocca, R., Aureli, D., 2019. Cimitero di Atella. Résultats de la mission 2018. Chronique des activités
- 1360 archéologiques de l'École française de Rome. https://doi.org/10.4000/cefr.3329
- 1361 Rocca, R., Aureli, D., Abruzzese, C., 2018. Cimitero di Atella. Nouvelles recherches et valorisation

- 1362 d'un gisement du Paléolithique ancien en Méditerranée. Chronique des activités archéologiques
- de l'École française de Rome. https://doi.org/10.4000/cefr.1910
- 1364 Rocca, R., Boschin, F., Aureli, D., 2021. Around an elephant carcass: Cimitero di Atella and
- 1365 Ficoncella in the behavioural variability during the Early Middle Pleistocene in italy, in: Konidaris,
- 1366 G., Barkai, R., Tourloukis, V., Harvati, K. (Eds.), Human-Elephant Interactions: From Past to
- 1367 Present, Tübingen Paleoanthropology Book Series Volume Contributions in Paleoanthropology.
- 1368 Tübingen University Press, Tübingen, pp. 287–302.
- 1369 Rocca, R., Da Costa, A., Germond, L., Aureli, D., 2020. Cimitero di Atella : résultats de la mission
- 1370 2019. Chronique des activités archéologiques de l'École française de Rome.
 1371 https://doi.org/10.4000/cefr.4436
- Sanders, D., Ostermann, M., Kramers, J., 2009. Quaternary carbonate-rocky talus slope successions
 (Eastern Alps, Austria): sedimentary facies and facies architecture. Facies 55, 345.
- 1374 https://doi.org/10.1007/s10347-008-0175-z
- 1375 Santagata, C., Moncel, M.-H., Piperno, M., 2020. Bifaces or not bifaces? Role of raw materials in the
- 1376 Middle Pleistocene. The example of levels E-E1, B and F (610–670 ka) at Notarchirico (Italy).
- 1377JournalofArchaeologicalScience:Reports33,102544.1378https://doi.org/10.1016/j.jasrep.2020.102544
- Santucci, E., Marano, F., Cerilli, E., Fiore, I., Lemorini, C., Palombo, M.R., Anzidei, A.P., Bulgarelli, 1379 1380 G.M., 2016. Palaeoloxodon exploitation at the Middle Pleistocene site of La Polledrara di 1381 Cecanibbio (Rome, Italy). Quaternary International, VIth International Conference on Mammoths 1382 and their Relatives, Part 2 406, Part Β, 169–182. 1383 https://doi.org/10.1016/j.quaint.2015.08.042Segre, A.G., 1962. Considerazioni preliminari sul 1384 paleolitico inferiore e sulla stratigrafia del bacino di Atella-Vitalba, Lucania. Quaternaria IV, 199-1385 203.
- Schiattarella, M., Beneduce, P., Di Leo, P., Giano, S.I., Giannandrea, P., Principe, C., 2005. Assetto
 strutturale ed evoluzione morfotettonica quaternaria del vulcano del Monte Vulture (Appennino

- 1388 Lucano). Bollettino-Società Geologica Italiana 124, 543.
- Schiattarella, M., Giannandrea, P., Principe, C., La Volpe, L., 2016. Note Illustrative della Carta
 Geologica d'Italia alla scala 1:50.000 Foglio 451 Melfi, in: Carta Geologica d'Italia Alla Scala
 1:50.000. Servizio Geologico d'Italia Litografia Artistica Cartografica, Firenze.
- 1392 Segre, A.G., 1957. Considerazioni preliminari sul Paleolitico inferiore e sulla stratigrafia del bacino
- di Atella-Vitalba (Lucania). Quaternaria IV, 199–203.
- 1394 Serangeli, J., Böhner, U., Van Kolfschoten, T., Conard, N.J., 2015. Overview and new results from
- 1395 large-scale excavations in Schöningen. Journal of Human Evolution, Special Issue: Excavations
- at Schöningen: New Insights into Middle Pleistocene Lifeways in Northern Europe 89, 27–45.

1397 <u>https://doi.org/10.1016/j.jhevol.2015.09.013</u>

- Strani, F., Bellucci, L., Iannucci, A., Iurino, D.A., Mecozzi, B., Sardella, R., 2022.
 Palaeoenvironments of the MIS 15 site of Cava di Breccia Casal Selce 2 (central Italian
 Peninsula) and niche occupation of fossil ungulates during Middle Pleistocene interglacials.
 Historical Biology 34, 555–565. https://doi.org/10.1080/08912963.2021.1935920
- 1402 Tourloukis, V., Thompson, N., Panagopoulou, E., Giusti, D., Konidaris, G.E., Karkanas, P., Harvati,
- 1403 K., 2018. Lithic artifacts and bone tools from the Lower Palaeolithic site Marathousa 1,
- 1404 Megalopolis, Greece: Preliminary results. Quaternary International, The Gates of Europe 497, 47–
- 1405 64. https://doi.org/10.1016/j.quaint.2018.05.043
- Tissoux, H., Falguères C., Voinchet P., Toyoda S., Bahain J.J., Despriée J. 2007. Potential use of Ti
 centre in ESR dating of Fluvial Sediment. Quaternary Geochronology, 2, 1-4, 367-372
- 1408 Tourloukis, V., Thompson, N., Panagopoulou, E., Giusti, D., Konidaris, G.E., Karkanas, P., Harvati,
- 1409 K., 2018. Lithic artifacts and bone tools from the Lower Palaeolithic site Marathousa 1,
- 1410 Megalopolis, Greece: Preliminary results. Quaternary International, The Gates of Europe 497, 47–
- 1411 64. <u>https://doi.org/10.1016/j.quaint.2018.05.043</u>
- 1412 Toyoda, S., Falguères, C., 2003. The method to represent the ESR signal intensity of the aluminium
- hole centre in quartz for the purpose of dating. Adv. ESR Appl. 20, 7–10.

- 1414 Toyoda, S., Voinchet, P., Falguères, C., Dolo, J.M., Laurent, M., 2000. Bleaching of ESR signal by
- the sunlight: a laboratory experiment for establishing the ESR dating of sediments. AppliedRadiation Isotopes, 52, 5, 1357–1362.
- 1417 van der Made, J., Rosell, J., Blasco, R., 2017. Faunas from Atapuerca at the Early–Middle Pleistocene
- 1418 limit: The ungulates from level TD8 in the context of climatic change. Quaternary International,
- 1419 What's happening now in Atapuerca? Latest research at the Sierra de Atapuerca sites 433, 296–
- 1420 346. https://doi.org/10.1016/j.quaint.2015.09.009
- 1421 van Kolfschoten, T., Parfitt, S.A., Serangeli, J., Bello, S.M., 2015. Lower Paleolithic bone tools from
- 1422 the 'Spear Horizon' at Schöningen (Germany). Journal of Human Evolution, Special Issue:
- 1423 Excavations at Schöningen: New Insights into Middle Pleistocene Lifeways in Northern Europe
- 1424 89, 226–263. https://doi.org/10.1016/j.jhevol.2015.09.012
- Villa, I.M., Buettner, A., 2009. Chronostratigraphy of Monte Vulture volcano (southern Italy):
 secondary mineral microtextures and 39Ar-40Ar systematics. Bull Volcanol 71, 1195.
 https://doi.org/10.1007/s00445-009-0294-6
- Villa, P., Boschian, G., Pollarolo, L., Saccà, D., Marra, F., Nomade, S., Pereira, A., 2021. Elephant
 bones for the Middle Pleistocene toolmaker. PLOS ONE 16, e0256090.
 https://doi.org/10.1371/journal.pone.0256090
- 1431 Voinchet, P., Bahain, J.-J., Falguères, C., Laurent, M., Dolo, J., Despriée, J., Gageonnet, R., Chaussé,
- 1432 C., 2004. ESR dating of quartz extracted from Quaternary sediments application to fluvial terraces
- 1433 system of northern France [Datation par résonance paramagnétique électronique (RPE) de quartz
- 1434 fluviatiles quaternaires: application aux systèmes de terrasses du nord de la France.]. Quaternaire
- 1435 15, 135–141. https://doi.org/10.3406/quate.2004.1761
- 1436 Voinchet, P., Pereira, A., Nomade, S., Falguères, C., Biddittu, I., Piperno, M., Moncel, M.-H., Bahain,
- 1437 J.-J., 2020. ESR dating applied to optically bleached guartz a comparison with 40Ar/39Ar
- 1438 chronologies on Italian Middle Pleistocene sequences, Quaternary International, 556, 113-123.
- 1439 Walker, R.G., James, N.P., 1992. Facies Models: Response to Sea Level Change. Geological

- 1440 Association of Canada, Toronto.
- 1441 Zucchelli, M., 1999. Bos primigenius Bojanus del Bacino di Atella (Potenza). Studi per l'ecologia
 1442 del Quaternario 19–30.
- 1443 Zucchelli, M., 2002. I cervidi dell'Acheuleano antico del Cimitero di Atella (Potenza). Studi per
- 1444 l'ecologia del Quaternario 3–17.
- 1445 Wright E., 2013. The history of the European aurochs (*Bos primigenius*) from the Middle Pleistocene
- 1446 to its extinction: an archaeological investigation of its evolution, morphological variability and
- 1447 response to human exploitation. PhD thesis. University of Sheffield.
































Stratigraphic units								
Supersynthem	Synthem	Subsynthem	Units code	Brief description of lithology and deposit interpretation	Thickness (m)	(Ma)		
MONTICCHIO	Laghi di Monticchio	Serra di Braida	LGM5	Alternating beds of massive (fall deposits) to cross-laminated (pyroclastic surge) tuff and lapilli stone.	4 - 5			
		Lago Piccolo	LGM4	Alternating beds (cm-thick) of massive to cross-laminated tuff and lapilli stone (pyroclastic surge).	5	0.141±11		
		Lago Grande	LGM3	Massive (fall deposits) to cross-laminated (pyroclastic surge) tuff with scattered blocks (cm to m-sized).	5			
		Piano Comune	LGM2	Fine to coarse-grained parallel laminated tuff locally with pisolites and cross- laminations (pyroclastic surge).	7			
		Casa Rossa	LGM1	Volcanic breccia deposit covered by fine-grained, cross-laminated tuff (pyroclastic surge).	1-5			
	Valle dei Grig – Fosso del	Masseria di Cuscito	VGC3	Fine-grained cross-laminated tuff (pyroclastic surge) cut by an erosive surface and covered by two massive breccia deposits (pyroclastic flow).	7			
	Corbo	Imbandina	VGC2	Massive beds of tuff and blocks (pyroclastic flow) with subordinate fine- grained cross-laminated tuff (pyroclastic surge); 4-m-thick clast-supported, coarse-grained, sedimentary and lava blocks bearing lapilli tuffs unconformably lie on the previous unit and underlie in turn 1-m-thick welded, agglutinated lapilli tuff (carbonatite).	8			
		Case Lopes	VGC1	Breccia deposits and thick beds of fine- to coarse-grained tuff, with cross- lamination (pyroclastic surge); at the base, a 4-m-thick black scoriae (basanite) is present.	15	0.530±22 0.494±5		
MONTE VULTURE	Melfi	Solagne - Arcidiaconata	SMF4	Travertine deposits.	4-5			
		Piano di Croce	SMF3	Lava flow (haüynitite), covered by 3-m-thick massive to cross-bedded tuff (pyroclastic surge).	6			
		Castello di Melfi	SMF2	Dark-grey lava flow ("Melfi haüynophyre" auctt.),	1 to 20	0.573±4		
		Gaudo	SMF1	Epiclastic volcanic alluvial conglomerates, with rare intercalations of sands, laterally heteropic to swamp sediments.	12			
	Barile	Ventaruolo	SBL4	Dark-grey-green purnice fall deposit, in dm-sized layers alternated with massive yellow tuff with blocks (pyroclastic flow), yellow tuff with accretionary lapilli (fall deposits) and yellow to red cross-bedded tuff (pyroclastic surge).	1.5-8	<0,610		
		Vulture – San Michele	SBL3	Massive deposits (10-15 m thick) of tuff with dm-sized heterolithologic blocks (pyroclastic flow), alternated with: i) levels of pumices (fallout deposits), ii) lava flow (foidites, tephro-foidites, tephrites, basanites); these units are laterally correlated with epiclastic volcanic conglomerates, sands, and massive pelites (alluvia) and locustring deposits).	500	0.609±9 0.611±6		
		Rionero	SBL2	Composed of three lithofacies. The lowermost lithofacies is made of massive dm-thick beds of pale-yellow fine-grained tuff (fall deposits) with horizons of accretionary lapilli and m-thick beds of fine-grained, cross-bedded tuff with lenses bearing cm-sized white punices (pyroclastic surge); the intermediate lithofacies is made of lapilli fallout (thephrite, foidites) alternated with dm-sized massive or trough cross-beds of yellowish tuff (pyroclastic flow); the uppermost lithofacies is constituted of massive proximal fall and surge deposits, with m-sized ballistic lava blocks. In the Atella and Venosa basins, the volcanic succession is heteropic to lacustrine and alluvial deposits).	44	0.630±20 0.672±6 0.714±18 <0.720±15		
		Toppo San Paolo	SBL1	Pale gray lava dome (haüyne phonolite).	140	0.673±19		
	Foggianello	Fara d'Olivo	FGG3	Composed of two ignimbrite units, both made of massive to planar cross- bedded tuff with sparse cm-sized pumices (trachyphonolite) and rare lithics (pyroclastic flow deposits).	10	Upper ignimbrite <0.755±21; <0.781±29 Lower ignimbrite 0.742±22; 0.737±16		
		Campanile	FGG2	Sequence of massive (fall deposits) and subordinate cross-bedded (pyroclastic surge) tuff.	3-4			
		Spinoritola	FGG1	Alluvial conglomerate with subordinate lava clast of the Spinoritola dikes (haüina-trachyte composition).	25-30	0.698±25; 0.698±48; 0.678±9; 0.698±8		

 Table 1: Description of the Monte Vulture Volcano stratigraphic units (after Giannandrea et al., 2004, 2006, and Stoppa et al., 2008).

 Unit codes correspond to those of the official 1:50,000-scale geological sheet (Foglio 451 Melfi, Carta Geologica d'Italia, 2010)

Table 2: List of pyroclastic and epiclastic lithofacies with a brief description and interpretation

Facies code	Facies	Sedimentary structures	Interpretation
Bmli	Poorly-sorted matrix-supported epiclastic volcanic breccia, with lithic industry. Angular to sub-angular pebbles ranging in size from 1 to 10 cm. Matrix consists of grey middle- to coarse-grained epiclastic volcanic sandstone.	Structureless/massive beds	Cohesive debris flows deposits
Suli	Epiclastic volcanic sandstone, with volcanic angular to sub-angular pebble (ranging in size from 1 to 10 cm) and lithic industry.	Poorly-defined undulated beds	Reworked colluvium
SBt	Coarse epiclastic volcanic sandstone and angular to sub-rounded epiclastic volcanic clast-supported breccia, with scattered cobbles (10-50 cm size).	Trough cross-beds	Channel infill by water stream flow of the reworked volcanic fallout and pyroclastic flow deposits
Sr	Gray fine-grained epiclastic volcanic sandstone.	Ripple cross-lamination	Ripples (lower flow regime)
Sh	Light grey epiclastic volcanic siltstone.	Horizontally thinly laminated	Decantation, lake deposits.
Dh	White to light-grey silty-clay diatomites.	1-15-cm-thick massive to laminated beds	Shallow-lake deposits
Su	Red siltstone.	Massive appearance or with poorly defined planar lamination in undulated beds	Hydrothermal alteration of volcanic material
TLBm	Medium- to coarse-grained tuff with fine- to coarse- grained angular lapilli stones and blocks (max 27 cm size).	Structureless/massive beds	Pyroclastic debris flows
LTm	Fine- to coarse-grained lapilli stone, with medium- grained tuff. Presence of some angular blocks max 10 cm size.	Structureless/massive beds	Pyroclastic debris flows
Lm	Coarse-grained clast-supported lapilli stone with grey pumice.	Structureless/massive beds	Lapilli fallout deposits
TLm	Grey fine- to coarse-grained tuff with fine-grained scattered lapilli, locally covered by 1-3-cm-thick light-grey fine-grained tuff.	Massive appearance or with poorly- defined planar lamination	Lower-concentration pyroclastic currents and co- current falls
Tng	Grey fine- to medium-grained tuff.	Normal grading	Decantation of a volcanic fall deposit in shallow fresh water
Tm	Fine- to coarse-grained tuff.	Structureless/massive or poorly- defined laminated beds	Fallout deposits

Table 3:ESR data and ages obtained for Atella samples.

Level	²³⁸ U (ppm)		23 (p	²³² Th K (ppm) (%)			Dα (μGy/a)	D _β (μGy/a)	D _{γ+cosmic} (μGy/a)	
L3	7.09± 0.15 20.		20.19	0 ± 0.28	1.35 ± 0.02		179 ± 3	2083 ± 32	2018 ± 102	
L1	4.44 ± 0.1	17	18.37	7 ± 0.32	1.19 ± 0.02		138 ± 4	1652 ± 38	2375 ± 110	
Level	Used center	Bl max. (%)		W (%)	De (Gy)		Da (µGy/a)	Ages (ka)	Mean Ages (ka)	
1.2	Al		51	10 . 5	1870 ± 100		4207 . 111	435 ± 23	442 ± 39	
L3	Ti -Li		100	10 ± 5	1990 ± 175		4297±111	463 ± 41		
L1	Al		53	10 ± 5	2134 ± 159		4147 + 101	515 ± 39	500 . 50	
	Ti -Li		100		2081 ± 175		$414/\pm 121$	502 ± 42	509 ± 50	

Table 4: Number of archaeological finds

Levels	Old excavation (Borzatti)	New excavation (EFR)	Total
L complex	107	0	107
		Deb 1: 261	_
F complex		Deb 2: 129	_
	667	273	940
Total	774	273	1047

Table 5: Distribution of lithic artefacts by technological category. Retouch flakes and flakes from shaping small tools: small retouch flakes, notch flakes, small flakes from shaping the edges of small tools; Small tools: entire and fragmented small tools; Core: entire core and fragment of core; Flakes: production flakes from a flaking reduction sequence; Large tools: Handaxes and large shaped tools; Indet: debris, indeterminate small fragment, chunks.

Technological categories	Retouch and small tool- shaping flakes	Small tools	Cores	Flakes	Large tools	Indet	Total
Level L	46	16	2	15	0	28	107
Level F	222	250	23	386	28	31	940
Total	268	266	25	401	28	59	1047

Size	ТОТ	TOT %
1 cm	54	9.2
2 cm	131	22.4
3 cm	116	19.8
4 cm	108	18.5
5 cm	67	11.5
6 cm	29	5.0
7 cm	20	3.4
8 cm	11	1.9
9 cm	7	1.2
10 cm	9	1.5
11 cm	4	0.7
12 cm	5	0.9
13 cm	3	0.5
14 cm	3	0.5
15 cm	5	0.9
16 cm	0	0.0
17 cm	3	0.5

Table 6: Size of faunal specimens in the whole site

18 cm	2	0.3
19 cm	2	0.3
20 cm	1	0.2
21 cm	1	0.2
25 cm	0	0.0
28 cm	1	0.2
30 cm	3	0.5
ТОТ	585	100

Table 7: Alteration of fragments

	F	Η	L
Rounded fragments	38.5	43.4	7.1
Slightly rounded edges	22.3	22.6	18.8
Not altered	39.2	34.0	74.1
ТОТ	462	53	85

Table 8: Taxonomy of faunal remains per level

	Levels							
Taxon	F	G	G/H	Н	L	М		
Bovinae indet.	9	0	0	0	3	0		
Bovid/Cervid	1	0	0	0	1	0		
Caprine	0	0	0	1	0	0		
Cervus elaphus	10	0	0	2	7	0		
Dama sp.	0	0	0	1	3	0		
Cervid	19	0	0	1	6	0		
Ungulate	9	0	0	3	3	0		
Palaeoloxodon	24	0	0	1	3	0		
antiquus								
Palaeoloxodon size	11	0	0	0	5	0		
Unidentified	385	2	9	45	54	1		
Small mammal	1	0	0	0	0	0		
Testudines	0	0	0	1	0	0		
ТОТ	469	2	9	55	85	1		

Declaration of interests

 \boxtimes The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

□The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

E-component/Supplementary Material

Click here to access/download **E-component/Supplementary Material** Atella_SM.pdf