



Searches for Continuous Gravitational Waves from Supernova Remnants in the First Part of the LIGO-Virgo-KAGRA Fourth Observing Run

A. G. Abac¹ , I. Abouelfettouh², F. Acernese^{3,4}, K. Ackley⁵ , C. Adamcewicz⁶ , S. Adhicary⁷ , D. Adhikari^{8,9}, N. Adhikari¹⁰ , R. X. Adhikari¹¹ , V. K. Adkins¹², S. Afroz¹³ , A. Agapito¹⁴, D. Agarwal¹⁵ , M. Agathos¹⁶ , N. Aggarwal¹⁷, S. Aggarwal¹⁸, O. D. Aguiar¹⁹ , I.-L. Ahrend²⁰, L. Aiello^{21,22} , A. Ain²³ , P. Ajith²⁴ , T. Akutsu^{25,26} , S. Albanesi^{27,28} , W. Ali^{29,30}, S. Al-Kershi^{8,9}, C. Alléne³¹, A. Allocca^{4,32} , S. Al-Shammari³³, P. A. Altin³⁴ , S. Alvarez-Lopez³⁵ , W. Amar³¹, O. Amarasinghe³³, A. Amato^{36,37} , F. Amicucci^{38,39} , C. Amra⁴⁰, A. Ananyeva¹¹, S. B. Anderson¹¹ , W. G. Anderson¹¹ , M. Andia⁴¹ , M. Ando⁴², M. Andrés-Carcasona⁴³ , T. Andric^{8,9,44,45} , J. Anglin⁴⁶, S. Ansoldi^{47,48} , J. M. Antelis⁴⁹ , S. Antier⁴¹ , M. Aoumi⁵⁰, E. Z. Appavuravther^{51,52} , S. Appert¹¹, S. K. Apple⁵³ , K. Arai¹¹ , A. Araya⁴² , M. C. Araya¹¹ , M. Arca Sedda^{44,45} , J. S. Areeda⁵⁴ , N. Aritomi², F. Armato^{29,30} , S. Armstrong⁵⁵ , N. Arnaud⁵⁶ , M. Arogeti⁵⁷ , S. M. Aronson¹², G. Ashton⁵⁸ , Y. Aso^{25,59} , L. Asprea²⁸, M. Assiduo^{60,61} , S. Assis de Souza Melo⁶², S. M. Aston⁶³, P. Astone³⁸ , F. Attadio^{38,39} , F. Aubin⁶⁴ , K. AultONeal⁶⁵ , G. Avallone⁶⁶ , E. A. Avila⁴⁹ , S. Babak²⁰ , C. Badger⁶⁷ , S. Bae⁶⁸ , S. Bagnasco²⁸ , L. Baiotti⁶⁹ , R. Bajpai⁷⁰ , T. Baka^{37,71} , A. M. Baker⁶, K. A. Baker⁷², T. Baker⁷³ , G. Baldi^{74,75} , N. Baldicchi^{51,76} , M. Ball⁷⁷ , G. Ballardín⁶², S. W. Ballmer⁷⁸ , S. Banagiri⁶ , B. Banerjee⁴⁴ , D. Bankar⁷⁹ , T. M. Baptiste¹², P. Baral¹⁰ , M. Baratti^{80,81} , J. C. Barayoga¹¹, B. C. Barish¹¹, D. Barker², N. Barman⁷⁹, P. Barneo^{82,83,84} , F. Barone^{4,85} , B. Barr⁸⁶ , L. Barsotti³⁵ , M. Barsuglia²⁰ , D. Barta⁸⁷ , A. M. Bartoletti⁸⁸, M. A. Barton⁸⁶ , I. Bartos⁴⁶, A. Basalae^{8,9} , R. Bassiri⁸⁹ , A. Basti^{80,81} , M. Bawaj^{51,76} , P. Baxi⁹⁰, J. C. Bayley⁸⁶ , A. C. Baylor¹⁰ , P. A. Baynard II⁵⁷, M. Bazzan^{91,92} , V. M. Bedakihale⁹³, F. Beirnaert⁹⁴ , M. Bejger⁹⁵ , D. Belardinelli²² , A. S. Bell⁸⁶ , D. S. Bellie⁹⁶ , L. Bellizzi^{80,81} , W. Benoit¹⁸ , I. Bentara⁵⁶ , J. D. Bentley⁹⁷ , M. Ben Yaala⁵⁵, S. Bera^{98,99} , F. Bergamin³³ , B. K. Berger⁸⁹ , S. Bernuzzi²⁷ , M. Beroiz¹¹ , D. Bersanetti²⁹ , T. Bertheas¹⁰⁰, A. Bertolini^{36,37} , J. Betzwieser⁶³ , D. Beveridge⁷² , G. Bevilacqua¹⁰¹ , N. Bevis¹⁰² , R. Bhandare¹⁰³, R. Bhatt¹¹, D. Bhattacharjee^{104,105} , S. Bhattacharyya¹⁰⁶ , S. Bhaumik⁴⁶ , V. Biancalana¹⁰¹ , A. Bianchi^{37,107} , I. A. Bilenko¹⁰⁸, G. Billingsley¹¹ , A. Binetti¹⁰⁹ , S. Bini^{11,74,75} , C. Binu¹¹⁰, S. Biot¹¹¹ , O. Birnholtz¹¹² , S. Biscoveanu⁹⁶ , A. Bisht⁹, M. Bitossi^{62,80} , M.-A. Bizouard¹¹³ , S. Blaber¹¹⁴ , J. K. Blackburn¹¹ , L. A. Blagg⁷⁷ , C. D. Blair^{63,72} , D. G. Blair⁷² , N. Bode^{8,9} , N. Boettner⁹⁷, G. Boileau¹¹³ , M. Boldrini³⁸ , G. N. Bolingbroke¹¹⁵ , A. Boliand^{40,116} , L. D. Bonavena⁴⁶ , R. Bondarescu⁸² , F. Bondu¹¹⁷ , E. Bonilla⁸⁹ , M. S. Bonilla⁵⁴ , A. Bonino¹¹⁸, R. Bonnand^{31,116} , A. Borchers^{8,9}, V. Boschi⁸⁰ , S. Bose¹¹⁹, V. Bossilkov⁶³ , Y. Bothra^{37,107} , A. Boudon⁵⁶, L. Bourg⁵⁷, M. Boyle¹²⁰, A. Bozzi⁶², C. Bradaschia⁸⁰ , P. R. Brady¹⁰ , A. Branch⁶³ , M. Branchesi^{44,45} , I. Braun¹⁰⁴, T. Briant¹²¹ , A. Brillet^{113,319} , M. Brinkmann^{8,9}, P. Brockill¹⁰, E. Brockmueller^{8,9} , A. F. Brooks¹¹ , B. C. Brown⁴⁶, D. D. Brown¹¹⁵, M. L. Brozzetti^{51,76} , S. Brunett¹¹, G. Bruno¹⁵, R. Bruntz¹²² , J. Bryant¹¹⁸ , Y. Bu¹²³ , F. Bucci⁶¹ , J. Buchanan¹²², O. Bulashenko^{82,83} , T. Bulik¹²⁴ , H. J. Bulten³⁷, A. Buonanno^{1,125} , K. Burtnyk² , R. Busicchio^{126,127} , D. Buskalic³¹, C. Buy¹⁰⁰ , R. L. Byer⁸⁹, G. S. Cabourn Davies⁷³ , R. Cabrita¹⁵ , V. Cáceres-Barbosa⁷ , L. Cadonati⁵⁷ , G. Cagnoli¹²⁸ , C. Cahillane⁷⁸ , A. Calafat⁹⁸ , T. A. Callister¹²⁹, E. Calloni^{4,32} , S. R. Callos⁷⁷ , G. Caneva Santoro⁴³ , K. C. Cannon⁴² , H. Cao³⁵, L. A. Capistran¹³⁰, E. Capocasa²⁰ , E. Capote^{2,11} , G. Capurri^{80,81} , G. Carapella^{66,131} , F. Carbognani⁶², M. Carlassara^{8,9}, J. B. Carlin¹²³ , T. K. Carlson¹³² , M. F. Carney¹⁰⁴ , M. Carpinelli^{62,126} , G. Carrillo⁷⁷, J. J. Carter^{8,9} , G. Carullo^{118,133} , A. Casallas-Lagos¹³⁴ , J. Casanueva Diaz⁶² , C. Casentini^{22,135} , S. Y. Castro-Lucas¹³⁶ , S. Caudill¹³² , M. Cavaglia¹⁰⁵ , R. Cavalieri⁶² , A. Ceja⁵⁴ , G. Cella⁸⁰ , P. Cerdá-Durán^{137,138} , E. Cesarini²² , N. Chabbra³⁴, W. Chaibi¹¹³, A. Chakraborty¹³ , P. Chakraborty^{8,9} , S. Chakraborty¹⁰³, S. Chalathadka Subrahmanya⁹⁷ , J. C. L. Chan¹³⁹ , M. Chan¹¹⁴, K. Chang¹⁴⁰, S. Chao^{140,141} , P. Charlton¹⁴² , E. Chassande-Mottin²⁰ , C. Chatterjee¹⁴³ , Debarati Chatterjee⁷⁹ , Deep Chatterjee³⁵ , M. Chaturvedi¹⁰³ , S. Chaty²⁰ , A. Chen¹⁴⁴ , A. H.-Y. Chen¹⁴⁵, D. Chen¹⁴⁶ , H. Chen¹⁴¹, H. Y. Chen¹⁴⁷ , S. Chen¹⁴³ , Yanbei Chen¹⁴⁸ , Yitian Chen¹²⁰ , H. P. Cheng¹⁴⁹, P. Chessa^{51,76} , H. T. Cheung⁹⁰ , S. Y. Cheung⁶, F. Chiadini^{131,150} , G. Chiarini^{8,9,92} , A. Chiba¹⁵¹ , A. Chincarini²⁹ , M. L. Chiofalo^{80,81} , A. Chiummo^{4,62} , C. Chou¹⁴⁵, S. Choudhary⁷² , N. Christensen^{113,152} , S. S. Y. Chua³⁴ , G. Ciani^{74,75} , P. Ciecielag⁹⁵ , M. Cieřlar¹²⁴ , M. Cifaldi²² , B. Cirok¹⁵³ , F. Clara², J. A. Clark^{11,57} , T. A. Clarke⁶ , P. Clearwater¹⁵⁴, S. Clesse¹¹¹, F. Cleva^{113,116} , E. Coccia^{43,44,45} , E. Codazzo^{155,156} , P.-F. Cohadon¹²¹ , S. Colace³⁰ , E. Colangeli⁷³, M. Colleoni⁹⁸ , C. G. Collette¹⁵⁷, J. Collins⁶³, S. Colloms⁸⁶ , A. Colombo^{127,158} , C. M. Compton², G. Connolly⁷⁷, L. Conti⁹² , T. R. Corbitt¹² , I. Cordero-Carrión¹⁵⁹ , S. Corezzi^{51,76} , N. J. Cornish¹⁶⁰ , I. Coronado¹⁶¹ , A. Corsi¹⁶² , R. Cottingham⁶³, M. W. Coughlin¹⁸ , A. Couineaux³⁸, P. Couvares^{11,57} , D. M. Coward⁷² , R. Coyne¹⁶³ , A. Cozzumbo⁴⁴, J. D. E. Creighton¹⁰ , T. D. Creighton¹⁶⁴, P. Cremonese⁹⁸ , S. Crook⁶³, R. Crouch² , J. Csizmazia², J. R. Cudell¹⁶⁵ , T. J. Cullen¹¹ , A. Cumming⁸⁶ , E. Cuoco^{166,167} , M. Cusinato¹³⁷ , L. V. Da Conceição¹⁶⁸ , T. Dal Canton⁴¹ , S. Dal Pra¹⁶⁹ , G. Dálya¹⁰⁰ , B. D'Angelo²⁹ , S. Danilishin^{36,37} , S. D'Antonio³⁸ , K. Danzmann^{8,9}, K. E. Darroch¹²², L. P. Darte⁶³ , R. Das¹⁰⁶, A. Dasgupta⁹³, V. Dattilo⁶² , A. Daumas²⁰ , N. Davari^{170,171} , I. Dave¹⁰³, A. Davenport¹³⁶, M. Davies⁴¹, T. F. Davies⁷² , D. Davis¹¹ , L. Davis⁷² , M. C. Davis¹⁸ , P. Davis^{172,173} , E. J. Daw¹⁷⁴ , M. Dax¹ , J. De Bolle⁹⁴ , M. Deenadayalan⁷⁹, J. Degallaix¹⁷⁵ , M. De Laurentis^{4,32}

F. De Lillo²³, S. Della Torre¹²⁷, W. Del Pozzo^{80,81}, A. Demagny³¹, F. De Marco^{38,39}, G. Demasi^{61,176},
 F. De Matteis^{21,22}, N. Demos³⁵, T. Dent¹⁷⁷, A. Depasse¹⁵, N. DePergola¹⁰², R. De Pietri^{178,179}, R. De Rosa^{4,32},
 C. De Rossi⁶², M. Desai³⁵, R. DeSalvo¹⁸⁰, A. DeSimone¹⁸¹, R. De Simone^{131,150}, A. Dhani¹, R. Diab⁴⁶, M. C. Díaz¹⁶⁴,
 M. Di Cesare^{4,32}, G. Dideron¹⁸², T. Dietrich¹, L. Di Fiore⁴, C. Di Fronzo⁷², M. Di Giovanni^{38,39}, T. Di Girolamo^{4,32},
 D. Diksha^{36,37}, J. Ding^{20,183}, S. Di Pace^{38,39}, I. Di Palma^{38,39}, D. Di Piero^{48,184}, F. Di Renzo⁵⁶, Divyajyoti³³,
 A. Dmitriev¹¹⁸, J. P. Docherty⁸⁶, Z. Doctor⁹⁶, N. Doerken¹⁶⁸, E. Dohmen², A. Doke¹³², A. Domiciano De Souza¹⁸⁵,
 L. D’Onofrio³⁸, F. Donovan³⁵, K. L. Dooley³³, T. Dooney⁷¹, S. Doravari⁷⁹, O. Dorosh¹⁸⁶, W. J. D. Doyle¹²²,
 M. Drago^{38,39}, J. C. Driggers², L. Dunn¹²³, U. Dupletsa⁴⁴, P.-A. Duverne²⁰, D. D’Urso^{155,170}, P. Dutta Roy⁴⁶,
 H. Duval¹⁸⁷, S. E. Dwyer², C. Eassa², M. Ebersold^{31,188}, T. Eckhardt⁹⁷, G. Eddolls⁷⁸, A. Effler⁶³, J. Eichholz³⁴,
 H. Einsle¹¹³, M. Eisenmann²⁵, M. Emma⁵⁸, K. Endo¹⁵¹, R. Enficiaud¹, L. Errico^{4,32}, R. Espinosa¹⁶⁴, M. Esposito^{4,32},
 R. C. Essick¹⁸⁹, H. Estellés¹, T. Etzel¹¹, M. Evans³⁵, T. Evstafyeva¹⁸², B. E. Ewing⁷, J. M. Ezquiaga¹³⁹,
 F. Fabrizi^{60,61}, V. Fafone^{21,22}, S. Fairhurst³³, A. M. Farah¹²⁹, B. Farr⁷⁷, W. M. Farr^{190,191}, G. Favaro⁹¹,
 M. Favata¹⁹², M. Fays¹⁶⁵, M. Fazio⁵⁵, J. Feicht¹¹, M. M. Fejer⁸⁹, R. Felicetti^{48,184}, E. Fenyvesi^{87,193}, J. Fernandes¹⁹⁴,
 T. Fernandes^{137,195}, D. Fernando¹¹⁰, S. Ferraiuolo^{38,39,196}, T. A. Ferreira¹², F. Fidecaro^{80,81}, P. Figura⁹⁵, A. Fiori^{80,81},
 I. Fiori⁶², M. Fishbach¹⁸⁹, R. P. Fisher¹²², R. Fittipaldi^{131,197}, V. Fiumara^{131,198}, R. Flaminio⁵¹, S. M. Fleischer¹⁹⁹,
 L. S. Fleming²⁰⁰, E. Floden¹⁸, H. Fong¹¹⁴, J. A. Font^{137,138}, F. Fontinele-Nunes¹⁸, C. Foo¹, B. Fornal²⁰¹, K. Franceschetti¹⁷⁸,
 F. Frappez³¹, S. Frasca^{38,39}, F. Frasconi⁸⁰, J. P. Freed⁶⁵, Z. Frei²⁰², A. Freise^{37,107}, O. Freitas^{137,195}, R. Frey⁷⁷,
 W. Frischhertz⁶³, P. Fritschel³⁵, V. V. Frolov⁶³, G. G. Fronze²⁸, M. Fuentes-Garcia¹¹, S. Fujii²⁰³, T. Fujimori²⁰⁴, P. Fulda⁴⁶,
 M. Fyffe⁶³, B. Gadre⁷¹, J. R. Gair¹, S. Galaudage¹⁸⁵, V. Galdi²⁰⁵, R. Gamba⁷, A. Gamboa¹, S. Gamoji¹⁸⁰,
 D. Ganapathy²⁰⁶, A. Ganguly⁷⁹, B. Garaventa²⁹, J. García-Bellido²⁰⁷, C. García-Quirós¹⁸⁸, J. W. Gardner³⁴,
 K. A. Gardner¹¹⁴, S. Garg⁴², J. Gargiulo⁶², X. Garrido⁴¹, A. Garron⁹⁸, F. Garufi^{4,32}, P. A. Garver⁸⁹, C. Gasbarra^{21,22},
 B. Gateley², F. Gautier²⁰⁸, V. Gayathri¹⁰, T. Gayer⁷⁸, G. Gemme²⁹, A. Gennai⁸⁰, V. Gennari¹⁰⁰, J. George¹⁰³,
 R. George¹⁴⁷, O. Gerberding⁹⁷, L. Gergely¹⁵³, Archisman Ghosh⁹⁴, Sayantan Ghosh¹⁹⁴, Shaon Ghosh¹⁹²,
 Shrobana Ghosh^{8,9}, Suprovo Ghosh²⁰⁹, Tathagata Ghosh⁷⁹, J. A. Giaime^{12,63}, K. D. Giardina⁶³, D. R. Gibson²⁰⁰,
 C. Gier⁵⁵, S. Gkaitatzis^{80,81}, J. Glanzer¹¹, F. Glotin⁴¹, J. Godfrey⁷⁷, R. V. Godley^{8,9}, P. Godwin¹¹, A. S. Goettel³³,
 E. Goetz¹¹⁴, J. Golomb¹¹, S. Gomez Lopez^{38,39}, B. Goncharov⁴⁴, G. González¹², P. Goodarzi²¹⁰, S. Goode⁶,
 A. W. Goodwin-Jones¹⁵, M. Gosselin⁶², R. Gouaty³¹, D. W. Gould³⁴, K. Govorkova³⁵, A. Grado^{51,76}, V. Graham⁸⁶,
 A. E. Granados¹⁸, M. Granata¹⁷⁵, V. Granata^{131,211}, S. Gras³⁵, P. Grassia¹¹, J. Graves⁵⁷, C. Gray², R. Gray⁸⁶, G. Greco⁵¹,
 A. C. Green^{37,107}, L. Green²¹², S. M. Green⁷³, S. R. Green²¹³, C. Greenberg¹³², A. M. Gretarsson⁶⁵, H. K. Griffin¹⁸,
 D. Griffith¹¹, H. L. Griggs⁵⁷, G. Grignani^{51,76}, C. Grimaud³¹, H. Grote³³, S. Grunewald¹, D. Guerra¹³⁷,
 D. Guetta²¹⁴, G. M. Guidi^{60,61}, A. R. Guimaraes¹², H. K. Gulati⁹³, F. Gulminelli^{172,173}, H. Guo¹⁴⁴, W. Guo⁷²,
 Y. Guo^{36,37}, Anuradha Gupta²¹⁵, I. Gupta⁷, N. C. Gupta⁹³, S. K. Gupta⁴⁶, V. Gupta¹⁸, N. Gupte¹, J. Gurs⁹⁷,
 N. Gutierrez¹⁷⁵, N. Guttman⁶, F. Guzman¹³⁰, D. Haba²¹⁶, M. Haberland¹, S. Haino²¹⁷, E. D. Hall³⁵, E. Z. Hamilton⁹⁸,
 G. Hammond⁸⁶, M. Haney³⁷, J. Hanks², C. Hanna⁷, M. D. Hannam³³, O. A. Hannuksela²¹⁸, A. G. Hanselman¹²⁹,
 H. Hansen², J. Hanson⁶³, S. Hanumasagar⁵⁷, R. Harada⁴², A. R. Hardison¹⁸¹, S. Harikumar¹⁸⁶, K. Haris^{37,71},
 I. Harley-Trochimczyk¹³⁰, T. Harmark¹³³, J. Harms^{44,45}, G. M. Harry²¹⁹, I. W. Harry⁷³, J. Hart¹⁰⁴, B. Haskell^{95,220,221},
 C.-J. Haster²¹², K. Haughian⁸⁶, H. Hayakawa⁵⁰, K. Hayama²²², M. C. Heintze⁶³, J. Heinze¹¹⁸, J. Heinzl³⁵,
 H. Heitmann¹¹³, F. Hellman²⁰⁶, A. F. Helmling-Cornell⁷⁷, G. Hemming⁶², O. Henderson-Sapir¹¹⁵, M. Hendry⁸⁶,
 I. S. Heng⁸⁶, M. H. Hennig⁸⁶, C. Henshaw⁵⁷, M. Heurs^{8,9}, A. L. Hewitt^{223,224}, J. Heynen¹⁵, J. Heyns³⁵,
 S. Higginbotham³³, S. Hild^{36,37}, S. Hill⁸⁶, Y. Himemoto²²⁵, N. Hirata²⁵, C. Hirose²²⁶, D. Hofman¹⁷⁵, B. E. Hogan⁶⁵,
 N. A. Holland^{37,107}, K. Holley-Bockelmann¹⁴³, I. J. Hollows¹⁷⁴, D. E. Holz¹²⁹, L. Honet¹¹¹, D. J. Horton-Bailey²⁰⁶,
 J. Hough⁸⁶, S. Hourihane¹¹, N. T. Howard¹⁴³, E. J. Howell⁷², C. G. Hoy⁷³, C. A. Hrishikesh²¹, P. Hsi³⁵,
 H.-F. Hsieh¹⁴¹, H.-Y. Hsieh¹⁴¹, C. Hsiung²²⁷, S.-H. Hsu¹⁴⁵, W.-F. Hsu¹⁰⁹, Q. Hu⁸⁶, H. Y. Huang¹⁴⁰, Y. Huang⁷,
 Y. T. Huang⁷⁸, A. D. Huddart²²⁸, B. Hughey⁶⁵, V. Hui³¹, S. Husa⁹⁸, R. Huxford⁷, L. Iampieri^{38,39}, G. A. Iandolo³⁶,
 M. Ianni^{21,22}, G. Iannone¹³¹, J. Iascai⁷⁷, K. Ide²²⁹, R. Iden²¹⁶, A. Ierardi^{44,45}, S. Ikeda¹⁴⁶, H. Imafuku⁴², Y. Inoue¹⁴⁰,
 G. Iorio⁹¹, P. Iosif^{48,184}, M. H. Iqbal³⁴, J. Irwin⁸⁶, R. Ishikawa²²⁹, M. Isi^{190,191}, K. S. Isleif²³⁰, Y. Itoh^{204,231},
 M. Iwaya²⁰³, B. R. Iyer²⁴, C. Jacquet¹⁰⁰, P.-E. Jacquet¹²¹, T. Jacquot⁴¹, S. J. Jadhav²³², S. P. Jadhav¹⁵⁴, M. Jain¹³²,
 T. Jain²²³, A. L. James¹¹, K. Jani¹⁴³, J. Janquart¹⁵, N. N. Janthaler²³², S. Jaraba²³³, P. Jaranowski²³⁴, R. Jaime⁹⁸,
 W. Javed³³, A. Jennings², M. Jensen², W. Jia³⁵, J. Jiang¹⁴⁹, H.-B. Jin^{235,236}, G. R. Johns¹²², N. A. Johnson⁴⁶,
 M. C. Johnston²¹², R. Johnston⁸⁶, N. Johny^{8,9}, D. H. Jones³⁴, D. I. Jones²⁰⁹, R. Jones⁸⁶, H. E. Jose⁷⁷, P. Joshi⁷,
 S. K. Joshi⁷⁹, G. Joubert⁵⁶, J. Ju²³⁷, L. Ju⁷², K. Jung²³⁸, J. Junker³⁴, V. Juste¹¹¹, H. B. Kabagöz^{35,63}, T. Kajita²³⁹,
 I. Kaku²⁰⁴, V. Kalogera⁹⁶, M. Kalomenopoulos²¹², M. Kamiizumi⁵⁰, N. Kanda^{204,231}, S. Kandhasamy⁷⁹,
 G. Kang²⁴⁰, N. C. Kannichel⁶, J. B. Kanner¹¹, S. A. KantiMahanty¹⁸, S. J. Kapadia⁷⁹, D. P. Kapasi⁵⁴, M. Karthikeyan¹³²,
 M. Kasprzak¹¹, H. Kato¹⁵¹, T. Kato²⁰³, E. Katsavounidis³⁵, W. Katzman⁶³, R. Kaushik¹⁰³, K. Kawabe², R. Kawamoto²⁰⁴,
 D. Keitel⁹⁸, L. J. Kemperman¹¹⁵, J. Kennington⁷, F. A. Kerkow¹⁸, R. Kesharwani⁷⁹, J. S. Key²⁴¹, R. Khadela^{8,9},
 S. Khadka⁸⁹, S. S. Khadkikar⁷, F. Y. Khalili¹⁰⁸, F. Khan^{8,9}, T. Khanam¹⁶², M. Khursheed¹⁰³, N. M. Khusid^{190,191},
 W. Kiendrebeogo^{113,242}, N. Kijbunchoo¹¹⁵, C. Kim²⁴³, J. C. Kim²⁴⁴, K. Kim²⁴⁵, M. H. Kim²³⁷, S. Kim²⁴⁶,

- 3

- B. G. Patterson³³, K. Paul¹⁰⁶, S. Paul⁷⁷, E. Payne¹¹, T. Pearce³³, M. Pedraza¹¹, A. Pele¹¹, F. E. Peña Arellano²⁸⁷, X. Peng¹¹⁸, Y. Peng⁵⁷, S. Penn²⁸⁸, M. D. Penuliar⁵⁴, A. Perego^{74,75}, Z. Pereira¹³², C. Périgois^{91,92,289}, G. Perna⁹¹, A. Perreca^{44,74,75}, J. Perret²⁰, S. Perriès⁵⁶, J. W. Perry^{37,107}, D. Pesios²⁵⁰, S. Peters¹⁶⁵, S. Petraccia²⁰⁵, C. Petrillo⁷⁶, H. P. Pfeiffer¹, H. Pham⁶³, K. A. Pham¹⁸, K. S. Phukon¹¹⁸, H. Phurailatpam²¹⁸, M. Piarulli¹⁰⁰, L. Piccari^{38,39}, O. J. Piccinni³⁴, M. Pichot¹¹³, M. Piendibene^{80,81}, F. Piergiovanni^{60,61}, L. Pierini³⁸, G. Pierra³⁸, V. Pierro^{131,290}, M. Pietrzak⁹⁵, M. Pillas¹⁶⁵, F. Pilo⁸⁰, L. Pinard¹⁷⁵, I. M. Pinto^{32,131,290,291}, M. Pinto⁶², B. J. Piotrkowski¹⁰, M. Pirello², M. D. Pitkin^{86,223}, A. Placidi⁵¹, E. Placidi^{38,39}, M. L. Planas⁹⁸, W. Plastino^{22,211}, C. Plunkett³⁵, R. Poggiani^{80,81}, E. Polini³⁵, J. Pomper^{80,81}, L. Pompili¹, J. Poon²¹⁸, E. Porcelli³⁷, E. K. Porter²⁰, C. Posnansky⁷, R. Poulton⁶², J. Powell¹⁵⁴, G. S. Prabhu⁷⁹, M. Pracchia¹⁶⁵, B. K. Pradhan⁷⁹, T. Pradier⁶⁴, A. K. Prajapati⁹³, K. Prasai²⁹², R. Prasanna²³², P. Prasia⁷⁹, G. Pratten¹¹⁸, G. Principe^{48,184}, G. A. Prodi^{74,75}, P. Prosperi⁸⁰, P. Proposito^{21,22}, A. C. Providence⁶⁵, A. Puecher¹, J. Pullin¹², P. Puppo³⁸, M. Pürer¹⁶³, H. Qi¹⁶, J. Qin³⁴, G. Quémener^{116,173}, V. Quetschke¹⁶⁴, P. J. Quinonez⁶⁵, N. Qutob⁵⁷, R. Rading²³⁰, I. Rainho¹³⁷, S. Raja¹⁰³, C. Rajan¹⁰³, B. Rajbhandari¹¹⁰, K. E. Ramirez⁶³, F. A. Ramis Vidal⁹⁸, M. Ramos Arevalo¹⁶⁴, A. Ramos-Buades^{37,98}, S. Ranjan⁵⁷, K. Ransom⁶³, P. Rapagnani^{38,39}, B. Ratto⁶⁵, A. Ravichandran¹³², A. Ray⁹⁶, V. Raymond³³, M. Razzano^{80,81}, J. Read⁵⁴, T. Regimbau³¹, S. Reid⁵⁵, C. Reissel³⁵, D. H. Reitze¹¹, A. I. Renzini^{11,126}, B. Revenu^{41,293}, A. Revilla Peña⁸², R. Reyes¹⁸⁰, L. Ricca¹⁵, F. Ricci^{38,39}, M. Ricci^{38,39}, A. Ricciardone^{80,81}, J. Rice⁷⁸, J. W. Richardson²¹⁰, M. L. Richardson¹¹⁵, A. Rijal⁶⁵, K. Riles⁹⁰, H. K. Riley³³, S. Rinaldi²⁶⁹, J. Rittmeyer⁹⁷, C. Robertson²²⁸, F. Robinet⁴¹, M. Robinson², A. Rocchi²², L. Rolland³¹, J. G. Rollins¹¹, A. E. Romano²⁹⁴, R. Romano^{3,4}, A. Romero³¹, I. M. Romero-Shaw²²³, J. H. Romie⁶³, S. Ronchini⁷, T. J. Rooke¹¹⁵, L. Rosa^{4,32}, T. J. Rosauer²¹⁰, C. A. Rose⁵⁷, D. Rosińska¹²⁴, M. P. Ross⁵³, M. Rossello-Sastre⁹⁸, S. Rowan⁸⁶, S. K. Roy^{190,191}, S. Roy¹⁵, D. Rozza^{126,127}, P. Ruggi⁶², N. Ruhama²³⁸, E. Ruiz Morales^{207,295}, K. Ruiz-Rocha¹⁴³, S. Sachdev⁵⁷, T. Sadecki², P. Saffarieh^{37,107}, S. Safi-Harb¹⁶⁸, M. R. Sah¹³, S. Saha¹⁴¹, T. Sainrat⁶⁴, S. Sajith Menon^{38,39,214}, K. Sakai²⁹⁶, Y. Sakai²⁷¹, M. Sakellariadou⁶⁷, S. Sakon⁷, O. S. Salafia^{126,127,158}, F. Salces-Carcoba¹¹, L. Salcomi⁶², M. Saleem¹⁴⁷, F. Salemi^{38,39}, M. Salle³⁷, S. U. Salunkhe⁷⁹, S. Salvador^{172,173}, C. Salvatore^{38,39}, A. Salvarese¹⁴⁷, A. Samajdar^{37,71}, A. Sanchez², E. J. Sanchez¹¹, L. E. Sanchez¹¹, N. Sanchis-Gual¹³⁷, J. R. Sanders¹⁸¹, E. M. Sanger¹, F. Santoliquido^{44,45}, F. Sarandrea²⁸, T. R. Saravanan⁷⁹, N. Sarin⁶, P. Sarkar^{8,9}, A. Sasli²⁵⁰, P. Sassi^{51,76}, B. Sassolas¹⁷⁵, R. Sato²²⁶, S. Sato¹⁵¹, Yukino Sato¹⁵¹, Yu Sato¹⁵¹, O. Sauter⁴⁶, R. L. Savage², T. Sawada⁵⁰, H. L. Sawant⁷⁹, S. Sayah¹⁷⁵, V. Scacco^{21,22}, D. Schaetzl¹¹, M. Scheel¹⁴⁸, A. Schiebelbein¹⁸⁹, M. G. Schiowski⁷⁸, P. Schmidt¹¹⁸, S. Schmidt⁷¹, R. Schnabel⁹⁷, M. Schneewind^{8,9}, R. M. S. Schofield⁷⁷, K. Schouteden¹⁰⁹, B. W. Schulte^{8,9}, B. F. Schutz^{8,9,33}, E. Schwartz²⁹⁷, M. Scialpi²⁹⁸, J. Scott⁸⁶, S. M. Scott³⁴, R. M. Sedas⁶³, T. C. Seetharamu⁸⁶, M. Seglar-Arroyo⁴³, Y. Sekiguchi²⁹⁹, D. Sellers⁶³, N. Sembo²⁰⁴, A. S. Sengupta³⁰⁰, E. G. Seo⁸⁶, J. W. Seo¹⁰⁹, V. Sequino^{4,32}, M. Serra³⁸, A. Sevrin¹⁸⁷, T. Shaffer², U. S. Shah⁵⁷, M. A. Shaikh²⁶⁰, L. Shao³⁰¹, A. K. Sharma⁹⁸, Preeti Sharma¹², Prianka Sharma¹⁰³, Ritwik Sharma¹⁸, S. Sharma Chaudhary¹⁰⁵, P. Shawhan¹²⁵, N. S. Shcheblanov^{262,302}, E. Sheridan¹⁴³, Z.-H. Shi¹⁴¹, M. Shikauchi⁴², R. Shimomura³⁰³, H. Shinkai³⁰³, S. Shirke⁷⁹, D. H. Shoemaker³⁵, D. M. Shoemaker¹⁴⁷, R. W. Short², S. ShyamSundar¹⁰³, A. Sider¹⁵⁷, H. Siegel^{190,191}, D. Sigg², L. Silenzi^{36,37}, L. Silvestri^{39,169}, M. Simmonds¹¹⁵, L. P. Singer³⁰⁴, Amitesh Singh²¹⁵, Anika Singh¹¹, D. Singh²⁰⁶, N. Singh⁹⁸, S. Singh^{59,216}, A. M. Sintes⁹⁸, V. Sipala^{155,170}, V. Skliris³³, B. J. J. Slagmolen³⁴, D. A. Slater¹⁹⁹, T. J. Slaven-Blair⁷², J. Smetana¹¹⁸, J. R. Smith⁵⁴, L. Smith^{48,86,184}, R. J. E. Smith⁶, W. J. Smith¹⁴³, S. Soares de Albuquerque Filho⁶⁰, M. Soares- Santos¹⁸⁸, K. Somiya²¹⁶, I. Song¹⁴¹, S. Soni³⁵, V. Sordini⁵⁶, F. Sorrentino²⁹, H. Sotani³⁰⁵, F. Spada⁸⁰, V. Spagnuolo³⁷, A. P. Spencer⁸⁶, P. Spinicelli⁶², A. K. Srivastava⁹³, F. Stachurski⁸⁶, C. J. Stark¹²², D. A. Steer³⁰⁶, N. Steinle¹⁶⁸, J. Steinlechner^{36,37}, S. Steinlechner^{36,37}, N. Stergioulas²⁵⁰, P. Stevens⁴¹, M. StPierre¹⁶³, M. D. Strong¹², A. Strunk², A. L. Stuver^{102,320}, M. Suchenek⁹⁵, S. Sudhagar⁹⁵, Y. Sudo²²⁹, N. Suelmann⁹⁷, L. Suleiman⁵⁴, K. D. Sullivan¹², J. Sun²⁴⁰, L. Sun³⁴, S. Sunil⁹³, J. Suresh¹¹³, B. J. Sutton⁶⁷, P. J. Sutton³³, K. Suzuki²¹⁶, M. Suzuki²⁰³, B. L. Swinkels³⁷, A. Syx¹¹⁶, M. J. Szczepańczyk³⁰⁷, P. Szewczyk¹²⁴, M. Tacca³⁷, H. Tagoshi²⁰³, S. C. Tait¹¹, K. Takada²⁰³, H. Takahashi²⁷¹, R. Takahashi²⁵, A. Takamori⁴², S. Takano³⁰⁸, H. Takeda^{309,310}, K. Takeshita²¹⁶, I. Takimoto Schmiegelow^{44,45}, M. Takou-Ayaoh⁷⁸, C. Talbot¹²⁹, M. Tamaki²⁰³, N. Tamanini¹⁰⁰, D. Tanabe¹⁴⁰, K. Tanaka⁵⁰, S. J. Tanaka²²⁹, S. Tanioka³³, D. B. Tanner⁴⁶, W. Tanner^{8,9}, L. Tao²¹⁰, R. D. Tapia⁷, E. N. Tapia San Martín³⁷, C. Taranto^{21,22}, A. Taruya³¹¹, J. D. Tasson¹⁵², J. G. Tau¹¹⁰, D. Tellez⁵⁴, R. Tenorio⁹⁸, H. Themann¹⁸⁰, A. Theodoropoulos¹³⁷, M. P. Thirugnanasambandam⁷⁹, L. M. Thomas¹¹, M. Thomas⁶³, P. Thomas², J. E. Thompson²⁰⁹, S. R. Thondapu¹⁰³, K. A. Thorne⁶³, E. Thrane⁶, J. Tissino^{44,45}, A. Tiwari⁷⁹, Pawan Tiwari⁴⁴, Praveer Tiwari¹⁹⁴, S. Tiwari¹⁸⁸, V. Tiwari¹¹⁸, M. R. Todd⁷⁸, M. Toffano⁹¹, A. M. Toivonen¹⁸, K. Toland⁸⁶, A. E. Tolley⁷³, T. Tomaru²⁵, V. Tommasini¹¹, T. Tomura⁵⁰, H. Tong⁶, C. Tong-Yu¹⁴⁰, A. Torres-Forne^{137,138}, C. I. Torrie¹¹, I. Tosta e Melo³¹², E. Tournefier³¹, M. Trad Nery¹¹³, K. Tran¹²², A. Trapananti^{51,52}, R. Travaglini¹⁶⁷, F. Travasso^{51,52}, G. Traylor⁶³, M. Trevor¹²⁵, M. C. Tringali⁶², A. Tripathi⁹⁰, G. Troian^{48,184}, A. Trovato^{48,184}, L. Trozzo⁴, R. J. Trudeau¹¹, T. Tsang³³, S. Tsuchida³¹³, L. Tsukada²¹², K. Turbang^{23,187}, M. Turconi¹¹³, C. Turski⁹⁴, H. Ubach^{82,83}, T. Uchiyama⁵⁰, R. P. Udall¹¹, T. Uehara³¹⁴, K. Ueno⁴², V. Undheim²⁷⁶, L. E. Uronen²¹⁸, T. Ushiba⁵⁰, M. Vacatello^{80,81}, H. Vahlbruch^{8,9}, N. Vaidya¹¹, G. Vajente¹¹, A. Vajpeyi⁶, J. Valencia⁹⁸

M. Valentini^{37,107}, S. A. Vallejo-Peña²⁹⁴, S. Vallerio²⁸, V. Valsan¹⁰, M. van Dael^{37,315}, E. Van den Bossche¹⁸⁷, J. F. J. van den Brand^{36,37,107}, C. Van Den Broeck^{37,71}, M. van der Sluys^{37,71}, A. Van de Walle⁴¹, J. van Dongen^{37,107}, K. Vandra¹⁰², M. VanDyke¹¹⁹, H. van Haevermaet²³, J. V. van Heijningen^{37,107}, P. Van Hove⁶⁴, J. Vanier²⁵⁸, M. VanKeuren¹⁰⁴, J. Vanosky², N. van Remortel²³, M. Vardaro^{36,37}, A. F. Vargas¹²³, V. Varma¹³², A. N. Vazquez⁸⁹, A. Vecchio¹¹⁸, G. Vedovato⁹², J. Veitch⁸⁶, P. J. Veitch¹¹⁵, S. Venikoudis¹⁵, R. C. Venterea¹⁸, P. Verdier⁵⁶, M. Vereecken¹⁵, D. Verkindt³¹, B. Verma¹³², Y. Verma¹⁰³, S. M. Vermeulen¹¹, F. Vetrano⁶⁰, A. Veutro^{38,39}, A. Vicere^{60,61}, S. Vidyant⁷⁸, A. D. Viets⁸⁸, A. Vijaykumar¹⁸⁹, A. Vilka¹¹⁰, N. Villanueva Espinosa¹³⁷, V. Villa-Ortega¹⁷⁷, E. T. Vincent⁵⁷, J.-Y. Vinet¹¹³, S. Viret⁵⁶, S. Vitale³⁵, H. Vocca^{51,76}, D. Voigt⁹⁷, E. R. G. von Reis², J. S. A. von Wrangel^{8,9}, W. E. Vossius²³⁰, L. Vujeva¹³⁹, S. P. Vyatchanin¹⁰⁸, J. Wack¹¹, L. E. Wade¹⁰⁴, M. Wade¹⁰⁴, K. J. Wagner¹¹⁰, L. Wallace¹¹, E. J. Wang⁸⁹, H. Wang²¹⁶, J. Z. Wang⁹⁰, W. H. Wang¹⁶⁴, Y. F. Wang¹, G. Waratkar¹⁹⁴, J. Warner², M. Was³¹, T. Washimi²⁵, N. Y. Washington¹¹, D. Watarai⁴², B. Weaver², S. A. Webster⁸⁶, N. L. Weickhardt⁹⁷, M. Weinert^{8,9}, A. J. Weinstein¹¹, R. Weiss^{35,321}, L. Wen⁷², K. Wette³⁴, J. T. Whelan¹¹⁰, B. F. Whiting⁴⁶, C. Whittle¹¹, E. G. Wickens⁷³, D. Wilken^{8,9}, A. T. Wilkin²¹⁰, B. M. Williams¹¹⁹, D. Williams⁸⁶, M. J. Williams⁷³, N. S. Williams¹, J. L. Willis¹¹, B. Willke^{8,9}, M. Wils¹⁰⁹, L. Wilson¹⁰⁴, C. W. Winborn¹⁰⁵, J. Winterflood⁷², C. C. Wipf¹¹, G. Woan⁸⁶, J. Woehler^{36,37}, N. E. Wolfe³⁵, H. T. Wong¹⁴⁰, I. C. F. Wong^{109,218}, K. Wong¹⁸⁹, T. Wouters^{37,71}, J. L. Wright², M. Wright^{71,86}, B. Wu⁷⁸, C. Wu¹⁴¹, D. S. Wu^{8,9}, H. Wu¹⁴¹, K. Wu¹¹⁹, Q. Wu⁵³, Y. Wu⁹⁶, Z. Wu¹⁰⁰, E. Wuchner⁵⁴, D. M. Wysocki¹⁰, V. A. Xu²⁰⁶, Y. Xu⁹⁸, N. Yadav²⁸, H. Yamamoto¹¹, K. Yamamoto¹⁵¹, T. S. Yamamoto⁴², T. Yamamoto⁵⁰, R. Yamazaki²²⁹, T. Yan¹¹⁸, K. Z. Yang¹⁸, Y. Yang¹⁴⁵, Z. Yarbrough¹², J. Yebana⁹⁸, S.-W. Yeh¹⁴¹, A. B. Yelikar¹⁴³, X. Yin³⁵, J. Yokoyama^{42,316}, T. Yokozawa⁵⁰, S. Yuan⁷², H. Yuzurihara⁵⁰, M. Zanolin⁶⁵, M. Zeeshan¹¹⁰, T. Zelenova⁶², J.-P. Zendri⁹², M. Zeoli¹⁵, M. Zerrad⁴⁰, M. Zevin⁹⁶, L. Zhang¹¹, N. Zhang⁵⁷, R. Zhang¹⁴⁹, T. Zhang¹¹⁸, C. Zhao⁷², Yue Zhao¹⁶¹, Yuhang Zhao²⁰, Z.-C. Zhao³¹⁷, Y. Zheng¹⁰⁵, H. Zhong¹⁸, H. Zhou⁷⁸, H. O. Zhu⁷², Z.-H. Zhu^{317,318}, A. B. Zimmerman¹⁴⁷, L. Zimmermann⁵⁶, M. E. Zucker^{11,35}, and J. Zweizig¹¹,

(The LIGO Scientific Collaboration, the Virgo Collaboration, and the KAGRA Collaboration)

- ¹ Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-14476 Potsdam, Germany
- ² LIGO Hanford Observatory, Richland, WA 99352, USA
- ³ Dipartimento di Farmacia, Università di Salerno, I-84084 Fisciano, Salerno, Italy
- ⁴ INFN, Sezione di Napoli, I-80126 Napoli, Italy
- ⁵ University of Warwick, Coventry CV 4 7AL, UK
- ⁶ OzGrav, School of Physics & Astronomy, Monash University, Clayton 3800, Victoria, Australia
- ⁷ The Pennsylvania State University, University Park, PA 16802, USA
- ⁸ Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-30167 Hannover, Germany
- ⁹ Leibniz Universität Hannover, D-30167 Hannover, Germany
- ¹⁰ University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA
- ¹¹ LIGO Laboratory, California Institute of Technology, Pasadena, CA 91125, USA
- ¹² Louisiana State University, Baton Rouge, LA 70803, USA
- ¹³ Tata Institute of Fundamental Research, Mumbai 400005, India
- ¹⁴ Centre de Physique Théorique, Aix-Marseille Université, Campus de Luminy, 163 Av. de Luminy, 13009 Marseille, France
- ¹⁵ Université catholique de Louvain, B-1348 Louvain-la-Neuve, Belgium
- ¹⁶ Queen Mary University of London, London E1 4NS, UK
- ¹⁷ University of California, Davis, Davis, CA 95616, USA
- ¹⁸ University of Minnesota, Minneapolis, MN 55455, USA
- ¹⁹ Instituto Nacional de Pesquisas Espaciais, 12227-010 São José dos Campos, São Paulo, Brazil
- ²⁰ Université Paris Cité, CNRS, Astroparticule et Cosmologie, F-75013 Paris, France
- ²¹ Università di Roma Tor Vergata, I-00133 Roma, Italy
- ²² INFN, Sezione di Roma Tor Vergata, I-00133 Roma, Italy
- ²³ Universiteit Antwerpen, 2000 Antwerpen, Belgium
- ²⁴ International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bengaluru 560089, India
- ²⁵ Gravitational Wave Science Project, National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan
- ²⁶ Advanced Technology Center, National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan
- ²⁷ Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, D-07743 Jena, Germany
- ²⁸ INFN Sezione di Torino, I-10125 Torino, Italy
- ²⁹ INFN, Sezione di Genova, I-16146 Genova, Italy
- ³⁰ Dipartimento di Fisica, Università degli Studi di Genova, I-16146 Genova, Italy
- ³¹ Univ. Savoie Mont Blanc, CNRS, Laboratoire d'Annecy de Physique des Particules - IN2P3, F-74000 Annecy, France
- ³² Università di Napoli "Federico II," I-80126 Napoli, Italy
- ³³ Cardiff University, Cardiff CF24 3AA, UK
- ³⁴ OzGrav, Australian National University, Canberra, Australian Capital Territory 0200, Australia
- ³⁵ LIGO Laboratory, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- ³⁶ Maastricht University, 6200 MD Maastricht, The Netherlands
- ³⁷ Nikhef, 1098 XG Amsterdam, The Netherlands
- ³⁸ INFN, Sezione di Roma, I-00185 Roma, Italy
- ³⁹ Università di Roma "La Sapienza," I-00185 Roma, Italy
- ⁴⁰ Aix Marseille Univ, CNRS, Centrale Med, Institut Fresnel, F-13013 Marseille, France
- ⁴¹ Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France
- ⁴² University of Tokyo, Tokyo, 113-0033, Japan
- ⁴³ Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Campus UAB, E-08193 Bellaterra (Barcelona), Spain
- ⁴⁴ Gran Sasso Science Institute (GSSI), I-67100 L'Aquila, Italy

- ⁴⁵ INFN, Laboratori Nazionali del Gran Sasso, I-67100 Assergi, Italy
- ⁴⁶ University of Florida, Gainesville, FL 32611, USA
- ⁴⁷ Dipartimento di Scienze Matematiche, Informatiche e Fisiche, Università di Udine, I-33100 Udine, Italy
- ⁴⁸ INFN, Sezione di Trieste, I-34127 Trieste, Italy
- ⁴⁹ Tecnológico de Monterrey, Escuela de Ingeniería y Ciencias, 64849, Monterrey, Nuevo León, Mexico
- ⁵⁰ Institute for Cosmic Ray Research, KAGRA Observatory, The University of Tokyo, 238 Higashi-Mozumi, Kamioka-cho, Hida City, Gifu 506-1205, Japan
- ⁵¹ INFN, Sezione di Perugia, I-06123 Perugia, Italy
- ⁵² Università di Camerino, I-62032 Camerino, Italy
- ⁵³ University of Washington, Seattle, WA 98195, USA
- ⁵⁴ California State University Fullerton, Fullerton, CA 92831, USA
- ⁵⁵ SUPA, University of Strathclyde, Glasgow G1 1XQ, UK
- ⁵⁶ Université Claude Bernard Lyon 1, CNRS, IP2I Lyon/IN2P3, UMR 5822, F-69622 Villeurbanne, France
- ⁵⁷ Georgia Institute of Technology, Atlanta, GA 30332, USA
- ⁵⁸ Royal Holloway, University of London, London TW20 0EX, UK
- ⁵⁹ Astronomical course, The Graduate University for Advanced Studies (SOKENDAI), 2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan
- ⁶⁰ Università degli Studi di Urbino “Carlo Bo,” I-61029 Urbino, Italy
- ⁶¹ INFN, Sezione di Firenze, I-50019 Sesto Fiorentino, Firenze, Italy
- ⁶² European Gravitational Observatory (EGO), I-56021 Cascina, Pisa, Italy
- ⁶³ LIGO Livingston Observatory, Livingston, LA 70754, USA
- ⁶⁴ Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France
- ⁶⁵ Embry-Riddle Aeronautical University, Prescott, AZ 86301, USA
- ⁶⁶ Dipartimento di Fisica “E.R. Caianiello,” Università di Salerno, I-84084 Fisciano, Salerno, Italy
- ⁶⁷ King’s College London, University of London, London WC2R 2LS, UK
- ⁶⁸ Korea Institute of Science and Technology Information, Daejeon 34141, Republic of Korea
- ⁶⁹ International College, Osaka University, 1-1 Machikaneyama-cho, Toyonaka City, Osaka 560-0043, Japan
- ⁷⁰ Accelerator Laboratory, High Energy Accelerator Research Organization (KEK), 1-1 Oho, Tsukuba City, Ibaraki 305-0801, Japan
- ⁷¹ Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University, 3584 CC Utrecht, The Netherlands
- ⁷² OzGrav, University of Western Australia, Crawley, Western Australia 6009, Australia
- ⁷³ University of Portsmouth, Portsmouth, PO1 3FX, UK
- ⁷⁴ Università di Trento, Dipartimento di Fisica, I-38123 Povo, Trento, Italy
- ⁷⁵ INFN, Trento Institute for Fundamental Physics and Applications, I-38123 Povo, Trento, Italy
- ⁷⁶ Università di Perugia, I-06123 Perugia, Italy
- ⁷⁷ University of Oregon, Eugene, OR 97403, USA
- ⁷⁸ Syracuse University, Syracuse, NY 13244, USA
- ⁷⁹ Inter-University Centre for Astronomy and Astrophysics, Pune 411007, India
- ⁸⁰ INFN, Sezione di Pisa, I-56127 Pisa, Italy
- ⁸¹ Università di Pisa, I-56127 Pisa, Italy
- ⁸² Institut de Ciències del Cosmos (ICCUB), Universitat de Barcelona (UB), c. Martí i Franquès, 1, 08028 Barcelona, Spain
- ⁸³ Departament de Física Quàntica i Astrofísica (FQA), Universitat de Barcelona (UB), c. Martí i Franquès, 1, 08028 Barcelona, Spain
- ⁸⁴ Institut d’Estudis Espacials de Catalunya, c. Gran Capità, 2-4, 08034 Barcelona, Spain
- ⁸⁵ Dipartimento di Medicina, Chirurgia e Odontoiatria “Scuola Medica Salernitana,” Università di Salerno, I-84081 Baronissi, Salerno, Italy
- ⁸⁶ IGR, University of Glasgow, Glasgow G12 8QQ, UK
- ⁸⁷ HUN-REN Wigner Research Centre for Physics, H-1121 Budapest, Hungary
- ⁸⁸ Concordia University Wisconsin, Mequon, WI 53097, USA
- ⁸⁹ Stanford University, Stanford, CA 94305, USA
- ⁹⁰ University of Michigan, Ann Arbor, MI 48109, USA
- ⁹¹ Università di Padova, Dipartimento di Fisica e Astronomia, I-35131 Padova, Italy
- ⁹² INFN, Sezione di Padova, I-35131 Padova, Italy
- ⁹³ Institute for Plasma Research, Bhat, Gandhinagar 382428, India
- ⁹⁴ Universiteit Gent, B-9000 Gent, Belgium
- ⁹⁵ Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, 00-716, Warsaw, Poland
- ⁹⁶ Northwestern University, Evanston, IL 60208, USA
- ⁹⁷ Universität Hamburg, D-22761 Hamburg, Germany
- ⁹⁸ IAC3–IEEC, Universitat de les Illes Balears, E-07122 Palma de Mallorca, Spain
- ⁹⁹ Aix-Marseille Université, Université de Toulon, CNRS, CPT, Marseille, France
- ¹⁰⁰ Laboratoire des 2 Infinis—Toulouse (L2IT-IN2P3), F-31062 Toulouse Cedex 9, France
- ¹⁰¹ Università di Siena, Dipartimento di Scienze Fisiche, della Terra e dell’Ambiente, I-53100 Siena, Italy
- ¹⁰² Villanova University, Villanova, PA 19085, USA
- ¹⁰³ RRCAT, Indore, Madhya Pradesh 452013, India
- ¹⁰⁴ Kenyon College, Gambier, OH 43022, USA
- ¹⁰⁵ Missouri University of Science and Technology, Rolla, MO 65409, USA
- ¹⁰⁶ Indian Institute of Technology Madras, Chennai 600036, India
- ¹⁰⁷ Department of Physics and Astronomy, Vrije Universiteit Amsterdam, 1081 HV Amsterdam, The Netherlands
- ¹⁰⁸ Lomonosov Moscow State University, Moscow 119991, Russia
- ¹⁰⁹ Katholieke Universiteit Leuven, Oude Markt 13, 3000 Leuven, Belgium
- ¹¹⁰ Rochester Institute of Technology, Rochester, NY 14623, USA
- ¹¹¹ Université libre de Bruxelles, 1050 Bruxelles, Belgium
- ¹¹² Bar-Ilan University, Ramat Gan, 5290002, Israel
- ¹¹³ Université Côte d’Azur, Observatoire de la Côte d’Azur, CNRS, Artemis, F-06304 Nice, France
- ¹¹⁴ University of British Columbia, Vancouver, BC V6T 1Z4, Canada
- ¹¹⁵ OzGrav, University of Adelaide, Adelaide, South Australia 5005, Australia
- ¹¹⁶ Centre national de la recherche scientifique, 75016 Paris, France
- ¹¹⁷ Univ Rennes, CNRS, Institut FOTON - UMR 6082, F-35000 Rennes, France
- ¹¹⁸ University of Birmingham, Birmingham B15 2TT, UK
- ¹¹⁹ Washington State University, Pullman, WA 99164, USA
- ¹²⁰ Cornell University, Ithaca, NY 14850, USA

- ¹²¹ Laboratoire Kastler Brossel, Sorbonne Université, CNRS, ENS-Université PSL, Collège de France, F-75005 Paris, France
- ¹²² Christopher Newport University, Newport News, VA 23606, USA
- ¹²³ OzGrav, University of Melbourne, Parkville, Victoria 3010, Australia
- ¹²⁴ Astronomical Observatory Warsaw University, 00-478 Warsaw, Poland
- ¹²⁵ University of Maryland, College Park, MD 20742, USA
- ¹²⁶ Università degli Studi di Milano-Bicocca, I-20126 Milano, Italy
- ¹²⁷ INFN, Sezione di Milano-Bicocca, I-20126 Milano, Italy
- ¹²⁸ Université de Lyon, Université Claude Bernard Lyon 1, CNRS, Institut Lumière Matière, F-69622 Villeurbanne, France
- ¹²⁹ University of Chicago, Chicago, IL 60637, USA
- ¹³⁰ University of Arizona, Tucson, AZ 85721, USA
- ¹³¹ INFN, Sezione di Napoli, Gruppo Collegato di Salerno, I-80126 Napoli, Italy
- ¹³² University of Massachusetts Dartmouth, North Dartmouth, MA 02747, USA
- ¹³³ Niels Bohr Institute, Copenhagen University, 2100 København, Denmark
- ¹³⁴ Universidad de Guadalajara, 44430, Guadalajara, Jalisco, Mexico
- ¹³⁵ Istituto di Astrofisica e Planetologia Spaziali di Roma, 00133 Roma, Italy
- ¹³⁶ Colorado State University, Fort Collins, CO 80523, USA
- ¹³⁷ Departamento de Astronomía y Astrofísica, Universitat de València, E-46100 Burjassot, València, Spain
- ¹³⁸ Observatori Astronòmic, Universitat de València, E-46980 Paterna, València, Spain
- ¹³⁹ Niels Bohr Institute, University of Copenhagen, 2100 København, Denmark
- ¹⁴⁰ National Central University, Taoyuan City 320317, Taiwan
- ¹⁴¹ National Tsing Hua University, Hsinchu City 30013, Taiwan
- ¹⁴² OzGrav, Charles Sturt University, Wagga Wagga, New South Wales 2678, Australia
- ¹⁴³ Vanderbilt University, Nashville, TN 37235, USA
- ¹⁴⁴ University of Chinese Academy of Sciences/International Centre for Theoretical Physics Asia-Pacific, Beijing 100190, People's Republic of China
- ¹⁴⁵ Department of Electrophysics, National Yang Ming Chiao Tung University, 101 University St., Hsinchu, Taiwan
- ¹⁴⁶ Kamioka Branch, National Astronomical Observatory of Japan, 238 Higashi-Mozumi, Kamioka-cho, Hida City, Gifu 506-1205, Japan
- ¹⁴⁷ University of Texas, Austin, TX 78712, USA
- ¹⁴⁸ CaRT, California Institute of Technology, Pasadena, CA 91125, USA
- ¹⁴⁹ Northeastern University, Boston, MA 02115, USA
- ¹⁵⁰ Dipartimento di Ingegneria Industriale (DIIN), Università di Salerno, I-84084 Fisciano, Salerno, Italy
- ¹⁵¹ Faculty of Science, University of Toyama, 3190 Gofuku, Toyama City, Toyama 930-8555, Japan
- ¹⁵² Carleton College, Northfield, MN 55057, USA
- ¹⁵³ University of Szeged, Dóm tér 9, Szeged 6720, Hungary
- ¹⁵⁴ OzGrav, Swinburne University of Technology, Hawthorn VIC 3122, Australia
- ¹⁵⁵ INFN Cagliari, Physics Department, Università degli Studi di Cagliari, Cagliari 09042, Italy
- ¹⁵⁶ Università degli Studi di Cagliari, Via Università 40, 09124 Cagliari, Italy
- ¹⁵⁷ Université Libre de Bruxelles, Brussels 1050, Belgium
- ¹⁵⁸ INAF, Osservatorio Astronomico di Brera sede di Merate, I-23807 Merate, Lecco, Italy
- ¹⁵⁹ Departamento de Matemáticas, Universitat de València, E-46100 Burjassot, València, Spain
- ¹⁶⁰ Montana State University, Bozeman, MT 59717, USA
- ¹⁶¹ The University of Utah, Salt Lake City, UT 84112, USA
- ¹⁶² Johns Hopkins University, Baltimore, MD 21218, USA
- ¹⁶³ University of Rhode Island, Kingston, RI 02881, USA
- ¹⁶⁴ The University of Texas Rio Grande Valley, Brownsville, TX 78520, USA
- ¹⁶⁵ Université de Liège, B-4000 Liège, Belgium
- ¹⁶⁶ DIFA—Alma Mater Studiorum Università di Bologna, Via Zamboni, 33 - 40126 Bologna, Italy
- ¹⁶⁷ Istituto Nazionale di Fisica Nucleare - Sezione di Bologna, viale Carlo Berti Pichat 6/2 - 40127 Bologna, Italy
- ¹⁶⁸ University of Manitoba, Winnipeg, MB R3T 2N2, Canada
- ¹⁶⁹ INFN-CNAF—Bologna, Viale Carlo Berti Pichat, 6/2, 40127, Bologna BO, Italy
- ¹⁷⁰ Università degli Studi di Sassari, I-07100 Sassari, Italy
- ¹⁷¹ INFN, Laboratori Nazionali del Sud, I-95125 Catania, Italy
- ¹⁷² Université de Normandie, ENSICAEN, UNICAEN, CNRS/IN2P3, LPC Caen, F-14000 Caen, France
- ¹⁷³ Laboratoire de Physique Corpusculaire Caen, 6 Blvd. du maréchal Juin, F-14050 Caen, France
- ¹⁷⁴ The University of Sheffield, Sheffield S10 2TN, UK
- ¹⁷⁵ Université Claude Bernard Lyon 1, CNRS, Laboratoire des Matériaux Avancés (LMA), IP21 Lyon / IN2P3, UMR 5822, F-69622 Villeurbanne, France
- ¹⁷⁶ Università di Firenze, Sesto Fiorentino I-50019, Italy
- ¹⁷⁷ IGFAE, Universidade de Santiago de Compostela, E-15782 Santiago de Compostela, Spain
- ¹⁷⁸ Dipartimento di Scienze Matematiche, Fisiche e Informatiche, Università di Parma, I-43124 Parma, Italy
- ¹⁷⁹ INFN, Sezione di Milano-Bicocca, Gruppo Collegato di Parma, I-43124 Parma, Italy
- ¹⁸⁰ California State University, Los Angeles, Los Angeles, CA 90032, USA
- ¹⁸¹ Marquette University, Milwaukee, WI 53233, USA
- ¹⁸² Perimeter Institute, Waterloo, ON N2L 2Y5, Canada
- ¹⁸³ Corps des Mines, Mines Paris, Université PSL, 60 Bd Saint-Michel, 75272 Paris, France
- ¹⁸⁴ Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy
- ¹⁸⁵ Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Lagrange, F-06304 Nice, France
- ¹⁸⁶ National Center for Nuclear Research, 05-400 Świerk-Otwock, Poland
- ¹⁸⁷ Vrije Universiteit Brussel, 1050 Brussel, Belgium
- ¹⁸⁸ University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland
- ¹⁸⁹ Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, ON M5S 3H8, Canada
- ¹⁹⁰ Stony Brook University, Stony Brook, NY 11794, USA
- ¹⁹¹ Center for Computational Astrophysics, Flatiron Institute, New York, NY 10010, USA
- ¹⁹² Montclair State University, Montclair, NJ 07043, USA
- ¹⁹³ HUN-REN Institute for Nuclear Research, H-4026 Debrecen, Hungary
- ¹⁹⁴ Indian Institute of Technology Bombay, Powai, Mumbai 400 076, India
- ¹⁹⁵ Centro de Física das Universidades do Minho e do Porto, Universidade do Minho, PT-4710-057 Braga, Portugal
- ¹⁹⁶ Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

- ¹⁹⁷ CNR-SPIN, I-84084 Fisciano, Salerno, Italy
- ¹⁹⁸ Scuola di Ingegneria, Università della Basilicata, I-85100 Potenza, Italy
- ¹⁹⁹ Western Washington University, Bellingham, WA 98225, USA
- ²⁰⁰ SUPA, University of the West of Scotland, Paisley PA1 2BE, UK
- ²⁰¹ Barry University, Miami Shores, FL 33168, USA
- ²⁰² Eötvös University, Budapest 1117, Hungary
- ²⁰³ Institute for Cosmic Ray Research, KAGRA Observatory, The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba 277-8582, Japan
- ²⁰⁴ Department of Physics, Graduate School of Science, Osaka Metropolitan University, 3-3-138 Sugimoto-cho, Sumiyoshi-ku, Osaka City, Osaka 558-8585, Japan
- ²⁰⁵ University of Sannio at Benevento, I-82100 Benevento, Italy and INFN, Sezione di Napoli, I-80100 Napoli, Italy
- ²⁰⁶ University of California, Berkeley, CA 94720, USA
- ²⁰⁷ Instituto de Física Teórica UAM-CSIC, Universidad Autónoma de Madrid, 28049 Madrid, Spain
- ²⁰⁸ Laboratoire d'Acoustique de l'Université du Mans, UMR CNRS 6613, F-72085 Le Mans, France
- ²⁰⁹ University of Southampton, Southampton SO17 1BJ, UK
- ²¹⁰ University of California, Riverside, Riverside, CA 92521, USA
- ²¹¹ Dipartimento di Ingegneria Industriale, Elettronica e Meccanica, Università degli Studi Roma Tre, I-00146 Roma, Italy
- ²¹² University of Nevada, Las Vegas, Las Vegas, NV 89154, USA
- ²¹³ University of Nottingham NG7 2RD, UK
- ²¹⁴ Ariel University, Ramat HaGolan St 65, Ariel, Israel
- ²¹⁵ The University of Mississippi, University, MS 38677, USA
- ²¹⁶ Graduate School of Science, Institute of Science Tokyo, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan
- ²¹⁷ Institute of Physics, Academia Sinica, 128 Sec. 2, Academia Rd., Nankang, Taipei 11529, Taiwan
- ²¹⁸ The Chinese University of Hong Kong, Shatin, NT, Hong Kong
- ²¹⁹ American University, WA, DC 20016, USA
- ²²⁰ Dipartimento di Fisica, Università degli Studi di Milano, Via Celoria 16, I-20133, Milano, Italy
- ²²¹ INFN, sezione di Milano, Via Celoria 16, I-20133, Milano, Italy
- ²²² Department of Applied Physics, Fukuoka University, 8-19-1 Nanakuma, Jonan, Fukuoka City, Fukuoka 814-0180, Japan
- ²²³ University of Cambridge, Cambridge CB2 1TN, UK
- ²²⁴ University of Lancaster, Lancaster LA1 4YW, UK
- ²²⁵ College of Industrial Technology, Nihon University, 1-2-1 Izumi, Narashino City, Chiba 275-8575, Japan
- ²²⁶ Faculty of Engineering, Niigata University, 8050 Ikarashi-2-no-cho, Nishi-ku, Niigata City, Niigata 950-2181, Japan
- ²²⁷ Department of Physics, Tamkang University, No. 151, Yingzhuan Rd., Danshui Dist., New Taipei City 25137, Taiwan
- ²²⁸ Rutherford Appleton Laboratory, Didcot OX11 0DE, UK
- ²²⁹ Department of Physical Sciences, Aoyama Gakuin University, 5-10-1 Fuchinobe, Sagami-hara City, Kanagawa 252-5258, Japan
- ²³⁰ Helmut Schmidt University, D-22043 Hamburg, Germany
- ²³¹ Nambu Yoichiro Institute of Theoretical and Experimental Physics (NITEP), Osaka Metropolitan University, 3-3-138 Sugimoto-cho, Sumiyoshi-ku, Osaka City, Osaka 558-8585, Japan
- ²³² Directorate of Construction, Services & Estate Management, Mumbai 400094, India
- ²³³ Observatoire Astronomique de Strasbourg, 11 Rue de l'Université, 67000 Strasbourg, France
- ²³⁴ Faculty of Physics, University of Białystok, 15-245 Białystok, Poland
- ²³⁵ National Astronomical Observatories, Chinese Academy of Sciences, 20A Datun Rd., Chaoyang Dist., Beijing, People's Republic of China
- ²³⁶ School of Astronomy and Space Science, University of Chinese Academy of Sciences, 20A Datun Rd., Chaoyang Dist., Beijing, People's Republic of China
- ²³⁷ Sungkyunkwan University, Seoul 03063, Republic of Korea
- ²³⁸ Department of Physics, Ulsan National Institute of Science and Technology (UNIST), 50 UNIST-gil, Ulju-gun, Ulsan 44919, Republic of Korea
- ²³⁹ Institute for Cosmic Ray Research, The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba 277-8582, Japan
- ²⁴⁰ Chung-Ang University, Seoul 06974, Republic of Korea
- ²⁴¹ University of Washington Bothell, Bothell, WA 98011, USA
- ²⁴² Laboratoire de Physique et de Chimie de l'Environnement, Université Joseph KI-ZERBO, 9GH2+3V5, Ouagadougou, Burkina Faso
- ²⁴³ Ewha Womans University, Seoul 03760, Republic of Korea
- ²⁴⁴ National Institute for Mathematical Sciences, Daejeon 34047, Republic of Korea
- ²⁴⁵ Korea Astronomy and Space Science Institute, Daejeon 34055, Republic of Korea
- ²⁴⁶ Department of Astronomy and Space Science, Chungnam National University, 9 Daehak-ro, Yuseong-gu, Daejeon 34134, Republic of Korea
- ²⁴⁷ Institute of Particle and Nuclear Studies (IPNS), High Energy Accelerator Research Organization (KEK), 1-1 Oho, Tsukuba City, Ibaraki 305-0801, Japan
- ²⁴⁸ Division of Science, National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan
- ²⁴⁹ Nagoya University, Nagoya, 464-8601, Japan
- ²⁵⁰ Department of Physics, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece
- ²⁵¹ Bard College, Annandale-On-Hudson, NY 12504, USA
- ²⁵² Technical University of Braunschweig, D-38106 Braunschweig, Germany
- ²⁵³ Institute of Mathematics, Polish Academy of Sciences, 00656 Warsaw, Poland
- ²⁵⁴ Astronomical Observatory, Jagiellonian University, 31-007 Kraków, Poland
- ²⁵⁵ Department of Physics and Astronomy, University of Padova, Via Marzolo, 8-35151 Padova, Italy
- ²⁵⁶ Sezione di Padova, Istituto Nazionale di Fisica Nucleare (INFN), Via Marzolo, 8-35131 Padova, Italy
- ²⁵⁷ Department of Physics, Nagoya University, ES building, Furocho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan
- ²⁵⁸ Université de Montréal/Polytechnique, Montreal, Quebec H3T 1J4, Canada
- ²⁵⁹ Indian Institute of Science Education and Research, Kolkata, Mohanpur, West Bengal 741252, India
- ²⁶⁰ Seoul National University, Seoul 08826, Republic of Korea
- ²⁶¹ Department of Computer Simulation, Inje University, 197 Inje-ro, Gimhae, Gyeongsangnam-do 50834, Republic of Korea
- ²⁶² NAVIER, École des Ponts, Univ Gustave Eiffel, CNRS, Marne-la-Vallée, France
- ²⁶³ Gravitational Wave Science Project, National Astronomical Observatory of Japan (NAOJ), Mitaka City, Tokyo 181-8588, Japan
- ²⁶⁴ Department of Physics, National Cheng Kung University, No.1, University Rd., Tainan City 701, Taiwan
- ²⁶⁵ St. Thomas University, Miami Gardens, FL 33054, USA
- ²⁶⁶ Scuola Normale Superiore, I-56126 Pisa, Italy
- ²⁶⁷ Institució Catalana de Recerca i Estudis Avançats, E-08010 Barcelona, Spain
- ²⁶⁸ Institut de Física d'Altes Energies, E-08193 Barcelona, Spain
- ²⁶⁹ Institut für Theoretische Astrophysik, Zentrum für Astronomie Heidelberg, Universität Heidelberg, Albert-Ueberle-Str. 2, 69120 Heidelberg, Germany
- ²⁷⁰ Institució Catalana de Recerca i Estudis Avançats (ICREA), Passeig de Lluís Companys, 23, 08010 Barcelona, Spain

- ²⁷¹ Research Center for Space Science, Advanced Research Laboratories, Tokyo City University, 3-3-1 Ushikubo-Nishi, Tsuzuki-Ku, Yokohama, Kanagawa 224-8551, Japan
- ²⁷² Tsinghua University, Beijing 100084, People's Republic of China
- ²⁷³ Institut des Hautes Études Scientifiques, F-91440 Bures-sur-Yvette, France
- ²⁷⁴ Faculty of Law, Ryukoku University, 67 Fukakusa Tsukamoto-cho, Fushimi-ku, Kyoto City, Kyoto 612-8577, Japan
- ²⁷⁵ Phenikaa Institute for Advanced Study (PIAS), Phenikaa University, Yen Nghia, Ha Dong, Hanoi, Vietnam
- ²⁷⁶ University of Stavanger, 4021 Stavanger, Norway
- ²⁷⁷ Physics Program, Graduate School of Advanced Science and Engineering, Hiroshima University, 1-3-1 Kagamiyama, Higashihiroshima City, Hiroshima 739-8526, Japan
- ²⁷⁸ GRAPPA, Anton Pannekoek Institute for Astronomy and Institute for High-Energy Physics, University of Amsterdam, 1098 XH Amsterdam, The Netherlands
- ²⁷⁹ University College London, London WC1E 6BT, UK
- ²⁸⁰ Observatoire de Paris, 75014 Paris, France
- ²⁸¹ Laboratoire Univers et Théories, Observatoire de Paris, 92190 Meudon, France
- ²⁸² Graduate School of Science and Technology, Niigata University, 8050 Ikarashi-2-no-cho, Nishi-ku, Niigata City, Niigata 950-2181, Japan
- ²⁸³ University of Maryland, Baltimore County, Baltimore, MD 21250, USA
- ²⁸⁴ CSIR-Central Glass and Ceramic Research Institute, Kolkata, West Bengal 700032, India
- ²⁸⁵ Consiglio Nazionale delle Ricerche—Istituto dei Sistemi Complessi, I-00185 Roma, Italy
- ²⁸⁶ Department of Astronomy, Yonsei University, 50 Yonsei-Ro, Seodaemun-Gu, Seoul 03722, Republic of Korea
- ²⁸⁷ Department of Physics, University of Guadalajara, Av. Revolución 1500, Colonia Olímpica C.P. 44430, Guadalajara, Jalisco, Mexico
- ²⁸⁸ Hobart and William Smith Colleges, Geneva, NY 14456, USA
- ²⁸⁹ INAF, Osservatorio Astronomico di Padova, I-35122 Padova, Italy
- ²⁹⁰ Dipartimento di Ingegneria, Università del Sannio, I-82100 Benevento, Italy
- ²⁹¹ Museo Storico della Fisica e Centro Studi e Ricerche “Enrico Fermi”, I-00184 Roma, Italy
- ²⁹² Kennesaw State University, Kennesaw, GA 30144, USA
- ²⁹³ Subatech, CNRS/IN2P3—IMT Atlantique—Nantes Université, 4 rue Alfred Kastler BP 20722 44307 Nantes CÉDEX 03, France
- ²⁹⁴ Universidad de Antioquia, Medellín, Colombia
- ²⁹⁵ Departamento de Física—ETSIDI, Universidad Politécnica de Madrid, 28012 Madrid, Spain
- ²⁹⁶ Department of Electronic Control Engineering, National Institute of Technology, Nagaoka College, 888 Nishikataai, Nagaoka City, Niigata 940-8532, Japan
- ²⁹⁷ Trinity College, Hartford, CT 06106, USA
- ²⁹⁸ Dipartimento di Fisica e Scienze della Terra, Università degli Studi di Ferrara, Via Saragat, 1, 44121, Ferrara FE, Italy
- ²⁹⁹ Faculty of Science, Toho University, 2-2-1 Miyama, Funabashi City, Chiba 274-8510, Japan
- ³⁰⁰ Indian Institute of Technology, Palaj, Gandhinagar, Gujarat 382355, India
- ³⁰¹ Kavli Institute for Astronomy and Astrophysics, Peking University, Yiheyuan Rd. 5, Haidian Dist., Beijing 100871, People's Republic of China
- ³⁰² Laboratoire MSME, Cité Descartes, 5 Blvd. Descartes, Champs-sur-Marne, 77454, Marne-la-Vallée Cedex 2, France
- ³⁰³ Faculty of Information Science and Technology, Osaka Institute of Technology, 1-79-1 Kitayama, Hirakata City, Osaka 573-0196, Japan
- ³⁰⁴ NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA
- ³⁰⁵ Faculty of Science and Technology, Kochi University, 2-5-1 Akebono-cho, Kochi-shi, Kochi 780-8520, Japan
- ³⁰⁶ Laboratoire de Physique de l'École Normale Supérieure, ENS, (CNRS, Université PSL, Sorbonne Université, Université Paris Cité), F-75005 Paris, France
- ³⁰⁷ Faculty of Physics, University of Warsaw, Ludwika Pasteura 5, 02-093 Warszawa, Poland
- ³⁰⁸ Laser Interferometry and Gravitational Wave Astronomy, Max Planck Institute for Gravitational Physics, Callinstrasse 38, 30167 Hannover, Germany
- ³⁰⁹ The Hakubi Center for Advanced Research, Kyoto University, Yoshida-honmachi, Sakyou-ku, Kyoto City, Kyoto 606-8501, Japan
- ³¹⁰ Department of Physics, Kyoto University, Kita-Shirakawa Oiwake-cho, Sakyou-ku, Kyoto City, Kyoto 606-8502, Japan
- ³¹¹ Yukawa Institute for Theoretical Physics (YITP), Kyoto University, Kita-Shirakawa Oiwake-cho, Sakyou-ku, Kyoto City, Kyoto 606-8502, Japan
- ³¹² University of Catania, Department of Physics and Astronomy, Via S. Sofia, 64, 95123, Catania CT, Italy
- ³¹³ National Institute of Technology, Fukui College, Geshi-cho, Sabae-shi, Fukui 916-8507, Japan
- ³¹⁴ Department of Communications Engineering, National Defense Academy of Japan, 1-10-20 Hashirimizu, Yokosuka City, Kanagawa 239-8686, Japan
- ³¹⁵ Eindhoven University of Technology, 5600 MB Eindhoven, The Netherlands
- ³¹⁶ Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU), WPI, The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba 277-8583, Japan
- ³¹⁷ Department of Astronomy, Beijing Normal University, Xijiekouwai St. 19, Haidian Dist., Beijing 100875, People's Republic of China
- ³¹⁸ School of Physics and Technology, Wuhan University, Bayi Rd. 299, Wuchang Dist., Wuhan, Hubei, 430072, People's Republic of China

Received 2026 March 26; revised 2026 April 15; accepted 2026 April 16; published 2026 June 15

Abstract

We present results from directed searches for continuous gravitational waves from a sample of 15 nearby supernova remnants, likely hosting young neutron star candidates, using data from the first eight months of the fourth observing run (O4) of the LIGO–Virgo–KAGRA Collaboration. The analysis employs five pipelines: four semicoherent methods—the Band-Sampled-Data directed pipeline, Weave, and two Viterbi pipelines (single- and dual-harmonic)—and PyStoch, a cross-correlation-based pipeline. These searches cover wide frequency bands and do not assume prior knowledge of the targets' ephemerides. No evidence of a signal is found from any of the 15 sources. We set 95% confidence-level upper limits on the intrinsic strain amplitude, with the most stringent constraints reaching $\sim 4 \times 10^{-26}$ near 300 Hz for the nearby source G266.2–1.2 (Vela Jr.). We also derive limits on neutron star ellipticity and r -mode amplitudes for the same source, with the best constraints reaching $\lesssim 10^{-7}$

³¹⁹ Deceased, 2026 March.

³²⁰ Deceased, 2024 September.

³²¹ Deceased, 2025 August.



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

and $\lesssim 10^{-5}$, respectively, at frequencies above 400 Hz. For frequencies above 200 Hz, these results represent the most sensitive wide-band directed searches for continuous gravitational waves from supernova remnants to date.

Unified Astronomy Thesaurus concepts: Neutron stars (1108); Supernova remnants (1667); Gravitational waves (678); Gravitational wave astronomy (675)

1. Introduction

Continuous gravitational waves (CWs) are long-duration, nearly monochromatic signals expected from nonaxisymmetric, rapidly rotating neutron stars (NSs). While transient gravitational waves (GWs) have been directly detected from numerous compact binary mergers (B. P. Abbott et al. 2019a; R. Abbott et al. 2021a, 2023, 2024, 2025; S. Olsen et al. 2022; A. H. Nitz et al. 2023), CWs remain undetected. These signals arise from time-varying mass quadrupole moments caused, for example, by small deviations from perfect axial symmetry in a spinning NS. Even minor imperfections in the star’s spherical shape can generate CWs. However, such signals are expected to be several orders of magnitude weaker in amplitude than those from mergers (see K. Riles 2023; K. Wette 2023; O. J. Piccinni 2022 for recent reviews). Detecting CWs would represent a major breakthrough in astrophysics, offering direct insights into NS structure and dynamics. This includes constraints on asymmetries and crustal deformations, the equation of state of ultradense matter, energy loss mechanisms, and magnetic field configurations (see, e.g., M. Sieniawska & M. Bejger 2019; B. Haskell et al. 2022; F. Gittins 2024; B. J. Owen 2026 for reviews). Moreover, CW observations could uncover hidden NSs or pulsars, invisible to traditional astronomical searches, which often rely on detecting electromagnetic pulsations. A GW detection provides an independent channel to reveal these hidden objects, expanding our view of the Galactic NS population. Though no CW detections have yet been reported, current searches are yielding significant upper limits on strain amplitude, ellipticity, and r -mode oscillation amplitudes. These limits are guided by theoretical predictions of r -mode behavior and relativistic frequency estimates from models of NSs with realistic equations of state (e.g., W. C. G. Ho & D. Lai 2000; A. Idrisy et al. 2015).

As such, young core-collapse supernova remnants (SNRs), known or expected to house NSs, represent excellent targets for the search for CWs. In the absence of pulsations, an NS may manifest itself as a central compact object (CCO) detected in X-ray wavelengths near the center of its hosting SNR, and typified by the NS discovered by the Chandra X-ray Observatory (CXO) in the young SNR Cassiopeia A (Cas A; G. G. Pavlov et al. 2000; A. De Luca 2017). Other manifestations of young NSs include a pointlike source powering a pulsar wind nebula (PWN), the synchrotron bubble inflated by the relativistic wind of the NS as it interacts with its surrounding SNR or interstellar medium, and typically observed in radio and/or X-ray energies (see, e.g., B. M. Gaensler & P. O. Slane 2006). Young NSs typically spin down rapidly due to strong electromagnetic and GW torques (B. Knispel & B. Allen 2008), complicating long-duration CW searches. Previous continuous GW searches for SNRs have employed various strategies with different trade-offs. Interesting sources like Cas A and SNR 1987A were targeted for the first time in the fifth science run of initial LIGO (J. Abadie et al. 2010; L. Sun et al. 2016), while other results for SNR 1987A can be found in the more recent O2 (B. J. Owen et al. 2022) and O3 (B. J. Owen et al. 2024; C. Salvatore et al.

2025) searches, which used data from the second and third Advanced LIGO–Virgo observing runs, respectively. Other, more extensive searches, including additional targets along with Cas A, were performed in S6 (J. Aasi et al. 2015a) and O1–O2 data (B. P. Abbott et al. 2019b; J. Ming et al. 2019; L. Lindblom & B. J. Owen 2020; M. Millhouse et al. 2020; M. A. Papa et al. 2020; J. Ming et al. 2022). Searches in O3 data (R. Abbott et al. 2021b, 2022a; J. Wang & K. Riles 2024), were characterized by being the broadest and most extensive performed to date (including a set of 15 sources in R. Abbott et al. 2021b), reaching sensitive results for the well-known sources Cas A and Vela Jr. in the most sensitive frequency range (J. Wang & K. Riles 2024). Competitive results were achieved by the latest deep searches performed using the Einstein@Home project (J. Ming et al. 2024, 2025; J. Morales et al. 2025). In particular, J. Ming et al. (2025) reported a $\sim 27\%$ false alarm probability candidate surviving for the source G347.3–0.5, which is also analyzed in this work by three of our pipelines. However, we did not find any compatible candidate using the search configurations adopted here. Previous results from the same project can be found in J. Ming et al. (2019), M. A. Papa et al. (2020), and J. Ming et al. (2022).

In this search, we apply five search pipelines to look for CWs from 15 young to middle-aged (i.e., $< 10^5$ yr) SNRs using data from the O4 run of the LIGO–Virgo–KAGRA collaboration. In particular, only the first eight months (O4a, beginning on 2023 May 24 and concluding on 2024 January 16) of data from LIGO Hanford and LIGO Livingston detectors were analyzed. All these pipelines were previously used in O3 or O3a analysis (R. Abbott et al. 2021b, 2022a; C. Salvatore et al. 2025). These include the directed Band-Sampled-Data (BSD) pipeline (O. J. Piccinni et al. 2018), the single-harmonic Viterbi (SHV) and dual-harmonic Viterbi (DHV) methods (L. Sun et al. 2018, 2019), and the Weave (K. Wette et al. 2018) semi-coherent searches, as well as the cross-correlation analysis by PyStoch (A. Ain et al. 2018). Together, the pipelines provide complementary parameter-space coverage and sensitivity to different emission scenarios.

The paper is organized as follows. Section 2 describes the SNR targets and their astrophysical properties, while in Section 3, we briefly discuss the signal models used in this search. Section 4 outlines the search methods and parameter space explored. Section 5 presents the results, including upper limits and astrophysical implications. Conclusions are summarized in Section 6. Appendices A–E detail technical information on the pipelines, the candidate’s follow-up and postprocessing, and the procedures used to set upper limits for BSD, PyStoch, SHV, DHV, and Weave, respectively.

2. Targets

2.1. Methodology

The SNR targets, summarized in Table 1, are selected from the Green catalog of radio SNRs (D. A. Green 2025) and

Table 1
List of 15 SNRs in This Work in Ascending Order of Galactic Longitude (for Galactic Remnants), and Associated CCOs or PWNe

Source	Compact Object or PWN Association	Age (kyr)	Distance (kpc)	R.A. (hh:mm:ss)	Decl. (deg:arcmin: arcsec)	Pipeline
G1.9+0.3	...	0.15–0.22	8.34–8.5	17:48:46.9	−27:10:16	SHV
G15.9+0.2	CCO CXOU J181852.0–150213	1–3	5–16	18:18:52.1	−15:02:14	SHV
G18.9–1.1	PWN, CXOU J182913.1–125113	2.8–6.1	1.6–4.7	18:29:13.1	−12:51:13	DHV, SHV
G39.2–0.3/3C 396	PWN, CXOU J190404.7+052711.8	4.7–5.2	6.2–9.5	19:04:04.7	5:27:12	BSD, DHV, SHV
G65.7+1.2/DA 495	PWN, CXOU J195217.04 +292552.5	3–155	1–3.6	19:52:17.0	29:25:53	BSD, DHV, SHV
G93.3+6.9/DA 530	PWN candidate, CXOU J205214+551722	3.2–7	4.4–8.3	20:52:14.0	55:17:22	BSD, DHV, SHV
G111.7–2.1/Cas A	CCO CXOUJ232327.9+584842	0.32–0.35	3.3–3.7	23:23:27.9	58:48:42	SHV,PyStoch, Weave
G189.1+3.0/IC 443	PWN G189.23+2.90	3–30	0.7–2.0	06:17:05.3	22:21:27	BSD, DHV, SHV
G266.2–1.2/Vela Jr.	CCO CXOU J085201.4–461753	2.4–5.1	0.5–1.3	08:52:01.4	−46:17:53	BSD, DHV, SHV PyStoch, Weave
G291.0–0.1/MSH 11–62	PWN G291.02–0.11, CXOU J111148.6 –603926	1.3–10	3–10	11:11:48.6	−60:39:26	BSD, SHV
G330.2+1.0	CCO CXOU J160103.1–513353	1–15	5–10	16:01:03.1	−51:33:5	SHV,PyStoch, Weave
G347.3–0.5/ RX J1713.7–3946	CCO 1WGA J1713.4–3949	1–1.7	0.9–1.1	17:13:28.3	−39:49:53	SHV
G350.1–0.3	CCO candidate XMMU J172054.5–372652	≤0.7	4.5–9	17:20:54.5	−37:26:52	BSD, DHV, SHV
G353.6–0.7	CCO XMMU J173203.3–344518	2.4–27	2.4–6.1	17:32:03.3	−34:45:18	
G279.7–31.9/SNR 1987A	NS or PWN	0.038	50	05:35:28.0	−69:16:13	PyStoch

Note. All are Galactic SNRs except for the last source, SNR 1987A, in the Large Magellanic Cloud. The first source, G1.9+0.3, is commonly believed to be a Type Ia SNR; however, it was included given the uncertainties in SNR classification and given that it is the youngest known SNR in the Milky Way galaxy. R.A. and decl. are in J2000 coordinates. CXOU (XMMU) refers to J2000 coordinates of the associated NS candidate source detected in X-rays with Chandra (XMM-Newton). All 5 pipelines were applied to the Vela Jr. SNR (G266.2–1.2); whereas for the other SNRs, only selected pipelines were applied. See Section 2 for details.

SNRcat, the online high-energy catalog of SNRs hosted by the University of Manitoba³²² (G. Ferrand & S. Safi-Harb 2012), with their distance and age estimates regularly updated. The youngest objects in this sample are considered to be among the most promising candidates for CWs since their expected high spin-down implies a potentially large energy budget for radiation in GWs.

The target selection is primarily based on a sample of core-collapse SNRs known to host isolated NSs or NS candidates. Their position has been inferred from the presence of a PWN or a pointlike X-ray source (a CCO or CCO candidate), both of which are located within or near the center of their hosting SNRs. The embedded NS candidates have not yet been discovered as pulsating sources, and as such, we do not know their characteristic ages. However, their ages are estimated from their hosting SNRs.

A CW search in the ~ 10 Hz–2 kHz (the sensitive range of Earth-based interferometer detectors) would reveal indirectly the presence of objects with spin periods in the ~ 1 –200 ms range since the CW frequency is (depending on the emission mechanism) approximately twice the spin frequency of the hidden star, and does not typically exceed that value (see Section 3). Such a search could help reveal the fastest and youngest spinning NSs. This range of rotation period values overlaps with the known pulsation periods of many young rotation-powered pulsars, as well as that of the three known pulsars in the CCOs family (with spin periods of 105, 112, and 424 ms; E. V. Gotthelf et al. 2013).

2.2. Target List and Comparison to O3a

The list of targeted SNRs is essentially the same as that chosen for O3a (R. Abbott et al. 2021b), with some updates summarized next. First, SNR 1987A (G279.7–31.9) was added, given it is believed to have now entered the remnant stage (i.e., the supernova blast wave is interacting with the circumstellar medium; A. P. Ravi et al. 2024) and shows evidence of a long-sought compact object or another powering engine (E. Greco et al. 2021; C. Fransson et al. 2024). This is also the youngest and only target in our sample with a well-known age and distance (being located in the Large Magellanic Cloud).

The previously considered target, G354.4+0.0 (also known as G354.46+0.07), is removed from our O4a list since it is no longer believed to be an SNR, but rather an H II region (N. Hurley-Walker et al. 2019). G1.9+0.3 is kept in our O4a list despite it being commonly believed to be a type Ia remnant (K. J. Borkowski et al. 2013). This is because a core-collapse origin is not completely ruled out, given the SNR’s location close to the Galactic center (e.g., S. P. Reynolds et al. 2008) and the known uncertainties in typing young SNRs from X-ray ejecta fits to nucleosynthesis yields (e.g., K. J. Borkowski et al. 2013; C. Braun et al. 2023; C. Treyturiuk et al. 2026). Being the youngest known Galactic SNR, G1.9+0.3 is potentially a promising target for CWs search, which may provide new insight into its nature.

Furthermore, the age and distance estimates shown in Table 1 have been updated since O3a to reflect the values in SNRcat (2025 version), including the most recent studies of these objects. We note that for Vela Jr., our distance upper limit is consistent with the most recent accurate distance

³²² <http://snrcat.physics.umanitoba.ca>

measurement of 1.41 ± 0.14 kpc (J. Suherli et al. 2026). These estimates affect the search parameters differently for each pipeline, as detailed in the following sections.

Of the 15 SNRs in Table 1, at least eight targets are searched using three different pipelines, while the remaining SNRs are searched using one or two pipelines. This is because each pipeline covers different parts of the parameter space, corresponding to slightly different assumptions of the characteristic age of the system. Furthermore, two remnants, G291.0–0.1 (MSH 11–62) and G330.2+1.0, already searched for in O3a by the SHV method (see Section 4.4), are added as targets in the BSD analysis (Section 4.2). G291.0–0.1 is a plerionic composite (i.e., a composite type SNR harboring a PWN) powered by a pulsar candidate (CXOU J111148.6–603926), and G330.2+1.0 hosts an X-ray emitting CCO (CXOU J160103.1–513353). These are young to middle-aged remnants of ages 1.3–10 kyr and 1–15 kyr, respectively.

3. Signal and Emission Models

A rotating NS with nonzero equatorial ellipticity has a time-varying mass quadrupole moment and emits CWs. The amplitude of this emission is determined by the star’s internal structure, including factors such as the nuclear equation of state, crustal rigidity, and magnetic field configuration (K. Glampedakis & L. Gualtieri 2018). Young NSs, in particular, are expected to retain relatively large deformations acquired at birth (B. Knispel & B. Allen 2008), although these asymmetries typically decay over time due to thermal, magnetic, and viscous relaxation processes (P. Haensel et al. 1990; O. Y. Gnedin et al. 2001; A. Y. Potekhin et al. 2015).

As the NS spins, it gradually loses angular momentum through several mechanisms, including GW emission. This energy loss causes a reduction in its spin (and GW) frequency, which can be modeled as ³²³

$$\dot{f} \propto -f^n, \quad n = \frac{\ddot{f}}{f^2}, \quad (1)$$

where the *braking index* n is a dimensionless parameter that describes the physical mechanism driving the NS’s energy loss. In practice, the spin-down evolution may be more complex than the simple power-law behavior described by Equation (1). The total energy loss may involve simultaneous contributions from both gravitational and electromagnetic (e.g., magnetic dipole) radiation. Some of the pipelines in this search look for signals with braking indices in the 2–7 range, as in J. Abadie et al. (2010), where the lowest $n = 2$ value is compatible with the braking indices of nearly all known pulsars (O. Hamil et al. 2015). In the case of spin-down dominated entirely by mass quadrupoles GW emission (as in the so-called *gravitar* scenario) $n = 5$, or $n = 7$ if the emission arises from *r*-mode oscillations (C. Palomba 2005). For magnetic dipole radiation, the braking index takes the value $n = 3$. The age of the NS can be estimated as

$$t_{\text{age}} = -\frac{f}{(n-1)\dot{f}} \left[1 - \left(\frac{f}{f_i} \right)^{n-1} \right] \Big|_{f \ll f_i} \approx -\frac{f}{(n-1)\dot{f}}, \quad (2)$$

³²³ Given the proportionality between the GW frequency and the star’s spin frequency f_* , we use f to denote either quantity in the formulae above. We will specify the relation between the GW frequency and f_* whenever it is needed.

where f_i is the spin frequency at birth. For older sources, where $f \ll f_i$, the expression is simply a function of f and $\dot{f}$ for a fixed braking index n .

The expected GW frequency depends on the emission mechanism and the star’s spinning frequency f_* : thermoelastic or magnetic quadrupoles emit at f_* or $2f_*$ (G. Ushomirsky et al. 2000; C. Cutler 2002; P. D. Lasky & A. Melatos 2013), while *r*-modes emit at approximately $4f_*/3$ (N. Andersson 1998; B. J. Owen et al. 1998; S. Caride et al. 2019; A. Idrisy et al. 2015; F. Gittins & N. Andersson 2023). Additional contributions may arise from superfluid dynamics within the NS, potentially leading to dual-harmonic emission scenarios ($f_* + 2f_*$; D. I. Jones 2010) or even to emission at higher-order harmonics of the spin frequency (A. Melatos et al. 2015; T. Cheunchitra et al. 2024). The dual-harmonic signal model, with components at f_* and $2f_*$, is given by (P. Jaranowski et al. 1998; L. Sun et al. 2019)

$$h_{2+}(t) = \frac{1}{2} h_0 (1 + \cos^2 \iota) \sin^2 \theta \cos \Phi(t), \quad (3)$$

$$h_{2\times}(t) = h_0 \cos \iota \sin^2 \theta \sin \Phi(t), \quad (4)$$

$$h_{1+}(t) = \frac{1}{8} h_0 \sin 2\iota \sin 2\theta \sin \frac{\Phi(t)}{2}, \quad (5)$$

$$h_{1\times}(t) = \frac{1}{4} h_0 \sin \iota \sin 2\theta \cos \frac{\Phi(t)}{2}, \quad (6)$$

where ι is the source inclination angle, θ is the wobble angle between the star’s rotation axis and its principal axis of the moment of inertia, and $\Phi(t)$ is the GW phase (at $2f_*$) observed at the detector.³²⁴ Standard CW searches target the simpler “triaxial star, principal-axis” model ($\theta = \pi/2$), where the h_1 terms vanish, leaving only the h_2 component at $2f_*$. The measured detector strain is then $h(t) = F_+(t; \alpha, \delta, \psi) h_{2+}(t) + F_\times(t; \alpha, \delta, \psi) h_{2\times}(t)$, with F_+ , $F_\times$ the antenna patterns for sky location (α, δ) and polarization angle ψ . The phase of the GW signal is primarily determined by the intrinsic evolution of the source’s spin frequency, characterized by its frequency and derivatives ($f, \dot{f}, \ddot{f}, \dots$) at a chosen reference time t_0 (with respect to the solar system barycenter):

$$\Phi(t) = \phi_0 + 2\pi \left[f(t - t_0) + \frac{\dot{f}}{2}(t - t_0)^2 + \frac{\ddot{f}}{6}(t - t_0)^3 + \dots \right], \quad (7)$$

where ϕ_0 is a constant phase. In addition to this intrinsic evolution, the observed phase is modulated by Doppler variations due to the relative motion between the source and the detector, as well as by the relativistic Einstein and Shapiro delays. In this search, we do not explicitly account for glitches; we assume that none of the targets have glitched during the considered observing time. We note that in addition to secular spin-down, random spin variations (timing noise or spin wandering) can affect the signal phase evolution (N. Namkham et al. 2019; A. Parthasarathy et al. 2019; M. E. Lower et al. 2020), driven by complex internal and magnetospheric processes (J. M. Cordes & G. Greenstein 1981; A. Melatos &

³²⁴ Here, we do not consider the scenarios with emissions at additional frequencies, e.g., when precession and triaxiality of the star are also considered (M. Zimmermann & E. Szednits 1979; C. Van Den Broeck 2005; P. D. Lasky & A. Melatos 2013).




Figure 1. Parameter space covered by each of the five pipelines for G266.2 -1.2 (Vela Jr.).

B. Link 2014). These complications motivate the use of semi-coherent methods (see Section 4), which coherently integrate short data segments (of duration T_{coh}) and combine the results incoherently across the full observing span. In the absence of electromagnetic measurements of $(f, \dot{f}, \dots)$, the maximum strain h_0^{max} emitted by a rotating NS (assuming its spin-down is fully dominated by GW emission) is inferred from the indirect, age-based spin-down limit for a source located at a distance D (K. Wette et al. 2008):

$$h_0^{\text{max}} = 2.27 \times 10^{-24} \left(\frac{1 \text{ kpc}}{D} \right) \left(\frac{1 \text{ kyr}}{t_{\text{age}}} \frac{I_{zz}}{10^{38} \text{ kg m}^2} \right)^{1/2}, \quad (8)$$

where I_{zz} is the NS’s moment of inertia.

4. Methods

This section briefly describes each of the five pipelines applied in this search, the parameter space investigated, and the emission mechanism considered. Pipeline details, along with the description of the postprocessing of the selected candidates, are reported in the Appendices. Each pipeline has a different approach when selecting the parameter space for the search. For comparison, we report in Figure 1 the values of the frequency f and first-order spin-down/up $\dot{f}$ parameter spanned by each search for Vela Jr., the only source targeted by all five pipelines. The different $(f, \dot{f})$ ranges covered by each pipeline arise from the distinct choices of signal emission models and from the computational constraints of each method, as described in Sections 4.2–4.6. PyStoch and Weave also include the second-order spin-down parameter in the signal template as indicated in Appendices B.1 and E.1, respectively.

4.1. Data

All the pipelines use data from the LIGO Hanford and Livingston detectors (J. Aasi et al. 2015b) in this search. The analysis covers the data collected between 2023 May 24, 15:00 UTC and 2024 January 16, 16:00 UTC—the period referred to as O4a (D. Ganapathy et al. 2023; W. Jia et al. 2024; A. G. Abac et al. 2025a; E. Capote et al. 2025; S. Soni et al. 2025). All the pipelines use data collected when the detectors

are in the science observing mode (E. Goetz & K. Riles 2024). The low-latency, cleaned, analysis-ready, calibrated data—C00 frames (channel names: H1:GDS-CALIB_STRAIN_CLEAN_AR and L1:GDS-CALIB_STRAIN_CLEAN_AR; A. D. Viets et al. 2018; A. D. Viets 2019; L. Sun et al. 2020; G. Vajente et al. 2020; LIGO-Virgo-KAGRA Collaboration et al. 2025; M. Wade et al. 2025) have been used to generate the different input files for each pipeline. All the inputs have passed through a glitch-gating procedure (F. Acernese et al. 2009; O. J. Piccinni et al. 2018; D. Davis et al. 2024). The time-domain BSD files, which are generated from the short fast Fourier transform (FFT) database files (P. Astone et al. 2005), a collection of short-time Fourier transform of duration 1024 s, have been cleaned using a double-gating procedure described in O. J. Piccinni et al. (2018) and F. Acernese et al. (2009). PyStoch begins its analysis using folded Stochastic Intermediate Data (SID), which are ultimately derived from the frames files. The SID consists of cross-correlated spectra and cross-detector noise estimates computed with a coherence time of 192 s and frequency resolution of $1/32$ Hz (A. I. Renzini et al. 2023). These data are then folded into a single sidereal day, resulting in the *folded* SID (A. Ain et al. 2018). The remaining pipelines start their analyses from the 30 minute short Fourier transforms (SFTs; LIGO Scientific Collaboration 2022). The SHV and DHV pipelines used the G02 version of gated data (D. Davis et al. 2024) while the Weave method used the earlier G01 release, which had spurious gates near boundaries of buffered data segments, a bug found to have negligible influence on search sensitivity. The lists of instrumental artefacts used to exclude some contaminated frequency bands are reported in E. Goetz et al. (2024).

4.2. BSD

The BSD pipeline, adapted for the SNR case (O. J. Piccinni et al. 2018; R. Abbott et al. 2021b), is a directed semi-coherent search method based on the Frequency–Hough (FH) transform (F. Antonucci et al. 2008; P. Astone et al. 2014a). Previous applications of this pipeline targeting the Galactic center during O2 and O3 are reported in O. J. Piccinni et al. (2020) and R. Abbott et al. (2022b), respectively. In this analysis, we target a subset of six sources, already searched for by the same pipeline in O3a (R. Abbott et al. 2021b), with the addition of two new targets: G291.0–0.1 and G330.2 +1.0. The full pipeline details can be found in R. Abbott et al. (2021b) and references therein. The signal model used in this search neglects the contributions from the second-order spin-down parameter (assumed always to be contained in a single $\dot{f}_0$ bin). The first-order spin-down range is defined as $-f_{\text{max}}/t_{\text{age}} \leq \dot{f} \leq 0.1 f_{\text{max}}/t_{\text{age}}$, with f_{max} the upper frequency of each 10 Hz band. We search both negative and positive spin-down to allow for possible spin-up scenarios. When available, we use the youngest reliable age estimate for each remnant from SNRcat’s 2025 version (G. Ferrand & S. Safi-Harb 2012). The source age used in the search also determines the range of second-order spin-down values that must be covered. However, this range will be larger for higher frequencies, determining de facto a limit on the maximum frequency used in this search (see Appendix A). Conversely, if the maximum frequency is fixed, the resulting second-order spin-down range places a lower bound on the source age (see, e.g., Equation (B2) of R. Abbott et al. 2021b). In this O4a analysis, we set a maximum frequency of 600 Hz, which

corresponds to a minimum source age of 3.8 kyr. The list of targets and corresponding parameter-space configurations is summarized in Appendix A.1.

4.3. PyStoch

The PyStoch pipeline (A. Ain et al. 2018) employs a Python-based tool built on the radiometer method, typically used to map the stochastic GW background across the sky. It analyzes cross-correlated and folded data from the LIGO Hanford and LIGO Livingston detectors in the [20–1726] Hz band. A previous search using O3 data can be found in C. Salvadore et al. (2025). In this search, PyStoch estimates the optimal signal-to-noise ratio as a function of the frequency for a specific sky location. This study focuses on four well-known SNRs: SNR 1987A, Vela Jr., Cas A, and G347.3–0.5; see Table 1. The explored parameter space for each target is summarized in Appendix B.1. The lower and upper bounds delineate the limiting cases of the physical scenarios being analyzed, i.e.,

$$-\frac{f}{t_{\text{age}}} \leq \dot{f} \leq -\frac{f}{6t_{\text{age}}}, \quad 2\frac{\dot{f}^2}{f} \leq \ddot{f} \leq 7\frac{\dot{f}^2}{f}, \quad (9)$$

including the frequency distribution parameter in Equation (B1).

4.4. Viterbi (Single Harmonic)

The SHV method implements a hidden Markov model (HMM) to efficiently track CW signals from NSs exhibiting secular spin down and spin wandering (S. Suvorova et al. 2016; L. Sun et al. 2018). Previous searches targeting young SNRs using this algorithm were conducted on Advanced LIGO’s data from the second observing run (O2; M. Millhouse et al. 2020; D. Beniwal et al. 2022) and the first half of the third LIGO–Virgo–KAGRA observing run (O3a; R. Abbott et al. 2021b). In this analysis, we search for 14 targets. For a given SNR with distance $D_{\min} \leq D \leq D_{\max}$ and age $t_{\text{age},\min} \leq t_{\text{age}} \leq t_{\text{age},\max}$, we define the searchable parameter space according to four criteria described in detail in Appendix C.1.

4.5. Viterbi (Dual Harmonic)

The DHV method (L. Sun et al. 2019) is a semi-coherent tracking algorithm designed to follow two frequency components simultaneously, improving sensitivity to signals emitted at both the stellar rotation frequency f_* and its second harmonic, $2f_*$. Other methods in this paper assume the conventional signal model in which the GWs are emitted only at a frequency twice the spin frequency of the star, $2f_*$. In addition, we assume that the signal frequency evolution is dominated by the star’s secular spin-down and approximated by a negatively biased random walk. The star’s unknown spin-down rate, which can vary over time, is covered in a range between zero and the maximum allowed spin-down rate $\dot{f}_{\text{max}}$, determined by the search configuration (see Appendix D). The list of targets and parameter space covered is given in Appendix D.1. Two search configurations are used for relatively older and younger targets. A minimum assumed characteristic age of the target, t_{age} , is selected to be either 20 kyr or 5 kyr, based on the estimated ages of the remnants (see Table 1). To mitigate the impact of noise artifacts, particularly due to combining dual harmonics, we impose a lower-

frequency cutoff at 50 Hz for f_* in all targets. This avoids contamination from instrumental lines that are more prevalent at lower frequencies. The maximum frequency is set by the selected t_{age} , i.e., $f_{\text{max}} = |\dot{f}_{\text{max}}|(n-1)t_{\text{age}}$. Assuming that the spin down is dominated by gravitational radiation, we take $n = 5$ (L. Sun et al. 2018; R. Abbott et al. 2021b).

4.6. Weave

The Weave pipeline is a template-based, semi-coherent search method based on the $\mathcal{F}$ -statistic (P. Jaranowski et al. 1998). The $\mathcal{F}$ -statistic computed for each segment analyzed is proportional to the square of the signal amplitude h_0^2 , maximized over h_0 , the unknown orientation angles ι and ψ , and the phase constant ϕ_0 . Thus, the search focuses on the parameters governing the phase evolution in Equation (7). Due to the high computational cost of template-based Weave searches, we limit our search to Cas A, Vela Jr., and G347.3–0.5 for GW signal frequency f ranges from 20 to 475 Hz. The search ranges for frequency derivatives are defined by assuming a power-law spin-down: $\dot{f} \propto -f^n$, with a braking index n ranging between 2 and 7. The maximum absolute values of $\dot{f}$ and $\ddot{f}$ at the lowest and highest search frequencies are reported in Appendix E.1.

5. Results

We do not find any statistically significant candidates from all our targets. All the candidates from each pipeline were discarded after the postprocessing and follow-up procedures described in Appendices A.2–E.2. None of the candidates surviving prior to the final steps and found with the reported search configurations seems to resemble the one listed in J. Ming et al. (2025). This may be expected, given the modest false alarm probability of 27% reported for the candidate. Nevertheless, to fully assess the nature of this candidate, more investigations using a fully coherent method on the full O4 dataset are needed.

In each case, the most significant outliers are consistent with instrumental artifacts or noise fluctuations. We report upper limits on the GW strain amplitude h_0 for each target and pipeline, including both single- and dual-harmonic models where applicable. These results are summarized in Section 5.1 and demonstrate a marked improvement over previous observing runs, particularly in frequency bands where the sensitivity of the detectors was optimal. For several remnants, the achieved upper limits surpass indirect spin-down constraints over wide frequency ranges. As discussed in Section 5.2, the upper-limit curves are translated into the ellipticity (i.e., the maximum degree of deformation the star can sustain) and, for the case of an emission happening via r -modes, into the upper bound on the r -mode amplitude parameter. Despite the nondetection, these results contribute to valuable constraints by placing meaningful bounds on the ellipticity and r -mode amplitude of the putative NSs. The upper limits for all pipelines and targets, as well as the notebook used to generate the plots, are available in Zenodo (LIGO Scientific Collaboration, Virgo Collaboration, and KAGRA Collaboration 2026).³²⁵

³²⁵ DOI:10.5281/zenodo.19567772




Figure 2. Upper Limits for the source G266.2–1.2 (Vela Jr.) obtained for the pipelines assuming the single-harmonic emission scenario. The dual-harmonic results for the same source are shown in Figure 12.

5.1. Upper Limits

In the following, we report the 95% confidence-level upper limits for each pipeline. These limits are obtained using different approaches, including injection-and-recovery methods or semianalytic estimates that were spot-checked against injections (see Appendices A.3–E.3). A common figure of merit for quantifying search sensitivity is known as the sensitivity depth $\mathcal{D}$ (B. Behnke et al. 2015; C. Dreissigacker et al. 2018):

$$\mathcal{D}(f) \equiv \frac{\sqrt{S_h(f)}}{h_0^{95\%}}, \quad (10)$$

where $\sqrt{S_h(f)}$ is an estimate of the effective strain amplitude spectral noise density. This quantity is independently estimated for each pipeline, as the input data used may differ between methods. We quantify the sensitivity depth for each pipeline.

In the following, we briefly describe the methodology used to set these upper limits in each pipeline. As a reference, in Figure 2, we report the results for Vela Jr., the only common target among all the pipelines. We note, however, that DHV results are not reported, as the interpretation of these upper limits for a dual-harmonic scenario is not directly comparable with that of a single-harmonic emission model. The upper limit from the sources investigated by DHV can be found in Figure 12, where the best Vela Jr. upper limit is 2.12×10^{-25} reached at 280 Hz.

BSD results. All the candidates surviving from the BSD pipeline are consistent with noise fluctuations (see Appendix

A.2), and no evidence of a CW is found in the LIGO O4a dataset. Constraints on the GW strain amplitude are derived using a semianalytical approach previously applied in O. J. Piccinni et al. (2020) and R. Abbott et al. (2022b). We corroborate the results by computing the 95% confidence-level upper limits on h_0 using artificial signals injected in randomly selected 1 Hz subbands. Results for all the targets are reported in Appendix A.3. For all the analyzed targets, the best strain sensitivity exceeds the indirect limits even when assuming the worst-case scenario (i.e., a very old source at the highest distance, as shown in Table 1), except for G65.7+1.2 and G330.2+1.0. When considering the best-case scenario, corresponding to young and nearby sources, all the targets exceed the indirect age-based limit for frequencies above 50 Hz, except in highly contaminated frequency bands.

The best achieved result at 95% confidence level is $h_0 = 5.94 \times 10^{-26}$ at around 250 Hz for the nearby source Vela Jr. The sensitivity depth for Vela Jr. at 200 Hz is $59.1 \text{ Hz}^{-1/2}$, while a maximum value of $81.8 \text{ Hz}^{-1/2}$ is reached for the lowest frequency band for the same source, as expected by the use of a longer coherence time. The lowest sensitivity depth among all targets at 200 Hz is achieved for G39.2–0.3 with $45.8 \text{ Hz}^{-1/2}$.

PyStoch results. None of the candidates survived the two follow-up procedures described in Appendix B.2. Subsequently, 95% confidence-level upper limits on the strain amplitude are computed across a range of frequency resolutions. The computation of the upper limits is performed within a Bayesian framework, incorporating prior distributions over relevant source parameters such as the inclination angle,

polarization angle, and calibration uncertainty. The final result is a marginalized posterior distribution for h_0 , from which the upper limit at a given confidence level (e.g., 95%) is extracted. The most stringent upper limits are obtained using the default resolution of $1/32$ Hz. The results for the whole frequency band are reported in Appendix B.3. The best upper limits are obtained using the default resolution, across the 20–1726 Hz band for each target in O4a. As expected, the search sensitivity degrades with increasing frequency bin combination; the results obtained directly from the default bin resolution yield tighter upper limits, while broader bin combinations are crucial to improve the signal-to-noise ratio of a signal spread over a broader frequency interval. The latter case, due to error propagation, will still be worse than the case of a narrower signal. The best upper limit is achieved for Vela Jr. at ~ 320 Hz, reaching 9.02×10^{-26} . We update the best upper limit on the SNR 1987A reaching 9.83×10^{-26} , beating previous results of B. J. Owen et al. (2024) and C. Salvadore et al. (2025). The sensitivity depth for Vela Jr. at the best (worst) frequency resolution is 36.5 (14.9) $\text{Hz}^{-1/2}$. The best sensitivity depth at the best frequency resolution is achieved for Cas A at 1693 Hz with $41.6 \text{ Hz}^{-1/2}$ and for G347.3–0.5 at 1161 Hz with $16.4 \text{ Hz}^{-1/2}$, with the latter being too low to reach the estimated strain of the candidate reported by J. Ming et al. (2025).

SHV results. All candidates surviving the veto steps of the SHV pipeline are consistent with noise fluctuations, and no evidence of CWs is found in the LIGO O4a dataset. None of them survived the follow-up procedures described in Appendix C.2. Constraints on the GW strain amplitude are derived using a scaling relation described in Appendix C.3, supported by numerical upper-limit calculations based on software injections. The results are shown in Appendix C.3. Across all analyzed targets, we state here the best upper limit achieved at 95% confidence level $h_0^{95\%} = 6.14 \times 10^{-26}$ at $f = 164$ Hz from G65.7+1.2. The best depth at 200 Hz across all sources is $51.7 \text{ Hz}^{-1/2}$ (G65.7+1.2), while Vela Jr. reaches a depth of 28.2 at the same frequency. The depth reached by SVH is not enough to probe the candidate in J. Ming et al. (2025). Furthermore, the search setup used for G347.3–0.5 is not able to produce candidates with first-order spin-down parameters compatible with the candidate.

DHV results. Without finding evidence of CWs in the DHV search (see Appendix D.2), we take a frequentist approach and estimate the 95% confidence strain sensitivity, denoted by $h_0^{95\%}$, in a frequency subband, and extend this estimate to the whole frequency band using an analytical scaling as described in Appendix D.3. Note that the strain sensitivity quoted in this pipeline is based on a different signal model from other pipelines (see Equations (3)–(6)). In this signal model, the strain sensitivity depends on the star’s wobble angle and orientation. Thus, we consider a specific scenario, representing the worst-case for single-band searches, where tracking the two frequency bands simultaneously offers the most significant sensitivity improvement compared to a single-band search. This scenario assumes $\theta = 45$ deg and $\cos \iota = 0$, corresponding to linearly polarized signals at both f_* and $2f_*$ (L. Sun et al. 2019). In Appendix D.3, the estimated $h_0^{95\%}$ in all subbands is reported. The best $h_0^{95\%}$ values over the full frequency band searched are 1.82×10^{-25} at $f_* = 213.25$ Hz and 2.12×10^{-25} at $f_* = 144.25$ Hz, for $T_{\text{coh}} = 12$ hr and 9 hr, respectively. These results are obtained from randomized sky

positions and hence apply to all sources using the same T_{coh} .³²⁶ The sensitivity depths for this pipeline are $13.9 \text{ Hz}^{-1/2}$ for $T_{\text{coh}} = 12$ hr and $12.4 \text{ Hz}^{-1/2}$ for $T_{\text{coh}} = 9$ hr.

Weave results. Using Weave, we find no significant evidence for a CW signal from the CCOs in the SNRs Cas A, Vela Jr., and G347.3–0.5 (see Appendix E.2). Given the absence of a detection, we determine 95% confidence-level frequentist upper limits on strain amplitude $h_0^{95\%}$ for each 1 Hz band, excluding saturated 0.1 Hz subbands. The upper-limit results for all the targets are reported in Appendix E.3, including the detailed procedure to compute them. The resulting frequency-dependent sensitivity depths have values at 200 Hz of $\mathcal{D}_{\text{CasA}} = 87.0 \text{ Hz}^{-1/2}$, $\mathcal{D}_{\text{Vela Jr.}} = 93.2 \text{ Hz}^{-1/2}$, and $\mathcal{D}_{\text{G347.3}} = 93.1 \text{ Hz}^{-1/2}$. In comparison with J. Ming et al. (2025), although our results are more sensitive at higher frequencies, our sensitivity at the candidate frequency reported by J. Ming et al. (2025) is lower. Consequently, our search does not reach the quoted strain estimated for this candidate, which is consistent with its nondetection.

For all the results, we observe an improvement with respect to O3a searches, compatible with the longer observation period in O4a (eight months compared to six months in O3a) and an improved detector sensitivity (of at least a factor of 1.5). Further improvements are observed for SHV, where longer coherence times and an adapted search setup have been applied. For Weave, an additional improvement with respect to O3a is due to the lower detection statistic threshold used to define initial candidates for follow-up.

5.2. Astrophysical Constraints

The GW strain amplitude can be converted into the fiducial ellipticity of the NS, ϵ (P. Jaranowski et al. 1998), and the r -mode amplitude parameter α (if an r -mode emission is assumed, B. J. Owen 2010). Given $h_0^{95\%}$, we constrain the ellipticity of the NS in terms of the GW frequency f via

$$\epsilon = 9.46 \times 10^{-5} \left(\frac{h_0}{10^{-24}} \right) \left(\frac{D}{1 \text{ kpc}} \right) \left(\frac{100 \text{ Hz}}{f} \right)^2, \quad (11)$$

assuming that the moment of inertia with respect to the rotation axis (I_{zz} for a perpendicular biaxial rotor) is 10^{38} kg m^2 . Here, we have considered the standard scenario of a single-harmonic emission at $2f_*$, valid for all the pipelines (Figures 3–6) except for DHV. For the specific dual-harmonic scenario, the GW emission is at both f_* and $2f_*$; hence, we can write Equation (11) in terms of the spin frequency of the star f_* ,

$$\epsilon = 2.36 \times 10^{-5} \left(\frac{h_0}{10^{-24}} \right) \left(\frac{D}{1 \text{ kpc}} \right) \left(\frac{100 \text{ Hz}}{f_*} \right)^2. \quad (12)$$

Figure 7 shows the constraints on ϵ as a function of f_* obtained for the DHV search. These constraints are converted from $h_0^{95\%}$ and are obtained for a specific scenario with source properties $\theta = 45$ deg and $\cos \iota = 0$. Therefore, the limits on ϵ in Figure 7 are not directly comparable to the results obtained in other conventional searches where $\theta = 90$ deg and the emission is only at $2f_*$ and reported in Figures 3–6.

³²⁶ The sensitivity is dominated by the length of T_{coh} (for a fixed T_{obs}) rather than the sky position of the source in the DHV pipeline (R. Abbott et al. 2021b).




Figure 3. Equatorial ellipticity (left) and r -mode amplitude α (right) with 95% confidence for all the BSD targets. Filled circles show the ellipticity or r -mode amplitude computed using the minimum estimated source distance $D_{\min}$. Shaded regions indicate the full range of values obtained when varying the source distance from $D_{\min}$ to $D_{\max}$ (see Table 1). The limits are derived from the upper-limit curves in Figure 8.




Figure 4. Equatorial ellipticity (left) and r -mode amplitude α (right) with 95% confidence for all targets studied in the PyStoch search. The filled circles indicate the minimum $\epsilon^{95\%}$ for each SNR, and the shaded regions span the full range of ellipticity values obtained by varying the distance between $D_{\min}$ and $D_{\max}$ (see Table 1).

We can also convert $h_0^{95\%}$ to limits on the amplitude of r -mode oscillations via (B. J. Owen 2010):

$$\alpha \simeq 0.028 \left(\frac{h_0}{10^{-24}} \right) \left(\frac{D}{1 \text{ kpc}} \right) \left(\frac{100 \text{ Hz}}{f} \right)^3. \quad (13)$$

Figures 3–6 show the constraints derived from the BSD, PyStoch, SHV, and Weave 95% confidence-level upper limits for the ellipticity (left) and the r -mode amplitude (right). Solid

curves correspond to $D = D_{\min}$, while shaded regions indicate the constraints obtained for distances in the range $D_{\min} \leq D \leq D_{\max}$. Distances are taken from Table 1.

We do not infer r -mode amplitudes for the DHV pipeline since the signal model adopted in the dual-harmonic search cannot be interpreted as current quadrupole emission from an r -mode.

The strongest ellipticity and r -mode constraints are obtained at the highest frequencies. Indeed, for all sources and targets,




Figure 5. Equatorial ellipticity (left) and r -mode amplitude α (right) with 95% confidence for all the SHV targets. The filled circles indicate the minimum $\epsilon^{95\%}$ and $\alpha^{95\%}$ for each SNR, and the shaded regions span the full range of ellipticity and r -mode amplitude values obtained by varying the distance between $D_{\min}$ and $D_{\max}$ (see Table 1).




Figure 6. Equatorial ellipticity and r -mode amplitude α upper limits (95% confidence level) from the Weave searches in each 1 Hz band for Cas A (blue), Vela Jr. (orange), and G347.3–0.5 (green), derived from the strain amplitude upper limits shown in Figure 13. The shaded region indicates the resulting ellipticity/ r -mode amplitude range calculated from the full estimated distance range for each SNR (see Table 1), and the solid dots correspond to the minimum distance estimate.

we are able to reach values below $\epsilon < 10^{-5}$ at the high end of the frequency band. Despite uncertainties in the source distances, we are approaching or exceeding the rough theoretical upper limit for normal NSs (N. K. Johnson-McDaniel & B. J. Owen 2013). The closest source, G266.2–1.2 (Vela Jr.), dramatically surpasses this limit, with ellipticity constraints reaching below $\epsilon \approx 10^{-7}$ across all pipelines.

These constraints are, however, model dependent: uncertainties in the stellar geometry and composition, including the

internal equation of state, the NS mass, the moment of inertia, and the magnetic field strength, can significantly affect the inferred values, introducing uncertainties of up to a factor of approximately 2–3. Despite these limitations, the improved detector sensitivities allow us to place tighter limits than previous O3 searches over a broader parameter space. In particular, for SNR 1987A, the constraints improve upon the O3 modeled search results in (B. J. Owen et al. 2024) by a factor of ~ 1.5 , consistent with the different integration time




Figure 7. Constraints on the NS ellipticity with 95% confidence from the DHV search as a function of f_* for (a) $T_{\text{coh}} = 12$ hr and (b) $T_{\text{coh}} = 9$ hr, converted from the $h_0^{95\%}$ values in Figure 12. The shaded region indicates the resulting ellipticity range calculated from the full estimated distance range for each SNR (see Table 1), and the solid dots correspond to the minimum distance estimate.

and the improved detector sensitivity used in this search. The most stringent constraints on the r -mode amplitude obtained above ~ 200 Hz strongly surpass the theoretical prediction level of $\alpha \sim 10^{-3}$ expected from nonlinear saturation mechanisms (R. Bondarescu et al. 2009), reaching values of $\alpha \sim 10^{-6}$ at the highest frequencies covered by the SHV pipeline.

6. Conclusions

In this work, we have presented the results of a comprehensive multipipeline search for CWs in SNRs potentially hosting young NSs, using data from the first eight months of the O4 LIGO–Virgo–KAGRA run. Our analysis employed complementary pipelines—BSD, PyStoch, SHV, DHV, and Weave—each targeting distinct signal models, parameter space, and offering varied sensitivity trade-offs. No evidence of a CW signal is found; all candidates are consistent with noise or eliminated by follow-up analyses.

We place new upper limits on the GW strain amplitude h_0 at 95% confidence level, beating previous O3 results. The BSD pipeline sets limits across 1 Hz subbands from 10 to 600 Hz, surpassing indirect energy conservation limits in several sources, notably for G65.7+1.2, G189.1+3.0, and Vela Jr.; this limit is beaten by a factor of ~ 20 , 25, 50, respectively, assuming the most optimistic scenario of a close-by and young source. However, these limits are surpassed even in the worst-case configuration for six out of eight sources in the BSD search. PyStoch results, spanning 20–1726 Hz at varying frequency resolutions, present the results from the search for CW emission from the SNR 1987A using an unmodeled approach, updating the limits found in the O3 search (C. Salvatore et al. 2025). The Viterbi pipelines provide constraints that incorporate spin wandering effects (single-harmonic) and simultaneous emission at f_* and $2f_*$ (dual-harmonic). Single-harmonic upper limits reach $h_0^{95\%} = 6.14 \times 10^{-26}$ at $f = 164.00$ Hz from G65.7+1.2. Dual-harmonic upper limits reach

$h_0^{95\%} = 1.82 \times 10^{-25}$ at $f_* = 213.25$ Hz (12 hr coherence) and 2.12×10^{-25} at $f_* = 144.25$ Hz (9 hr coherence), consistent with analytic expectations. The Weave pipeline’s high-resolution follow-up confirms all initial candidates are consistent with noise or instrumental artifacts reaching the best $h_0^{95\%}$ limit on the signal strain $\approx 4.33 \times 10^{-26}$ near 283 Hz for Cas A, $\approx 3.92 \times 10^{-26}$ near 266 Hz for Vela Jr., and $\approx 3.85 \times 10^{-26}$ near 378 Hz for G347.3–0.5.

Together, these results represent a significant advance in search sensitivity compared with previous searches for CWs from young NSs in nearby SNRs. These results provide the most stringent upper limits to date across the entire analyzed frequency band for 12 out of the 15 sources. For Cas A, Vela Jr., and G347.3–0.5, targets of recent deep searches (J. Ming et al. 2025; J. Morales et al. 2025), our results show improved limits above 200 Hz within the frequency band explored by Weave. While no signals are detected, the improved upper limits place stronger constraints on NS deformation and energy loss models. Future observing runs with longer datasets and enhanced detector sensitivity, combined with multipipeline strategies demonstrated here, will further improve detection prospects.

Acknowledgments

This material is based upon work supported by NSF’s LIGO Laboratory, which is a major facility fully funded by the National Science Foundation. The authors also gratefully acknowledge the support of the Science and Technology Facilities Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen/Germany for support of the construction of Advanced LIGO and the construction and operation of the GEO 600 detector. Additional support for Advanced LIGO was provided by the Australian Research Council. The authors gratefully acknowledge the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Centre National de la Recherche Scientifique (CNRS), and the Netherlands Organization for Scientific

Research (NWO) for the construction and operation of the Virgo detector and the creation and support of the EGO consortium. The authors also gratefully acknowledge research support from these agencies as well as by the Council of Scientific and Industrial Research of India, the Department of Science and Technology, India, the Science & Engineering Research Board (SERB), India, the Ministry of Human Resource Development, India, the Spanish Agencia Estatal de Investigación (AEI), the Spanish Ministerio de Ciencia, Innovación y Universidades, the European Union NextGenerationEU/PRTR (PRTR-C17.I1), the ICSC—Centro Nazionale di Ricerca in High Performance Computing, Big Data and Quantum Computing, funded by the European Union Next-GenerationEU, the Comunitat Autònoma de les Illes Balears through the Conselleria d'Educació i Universitats, the Conselleria d'Innovació, Universitats, Ciència i Societat Digital de la Generalitat Valenciana and the CERCA Programme Generalitat de Catalunya, Spain, the Polish National Agency for Academic Exchange, the National Science Centre of Poland and the European Union—European Regional Development Fund; the Foundation for Polish Science (FNP), the Polish Ministry of Science and Higher Education, the Swiss National Science Foundation (SNSF), the Russian Science Foundation, the European Commission, the European Social Funds (ESF), the European Regional Development Funds (ERDF), the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, the Hungarian Scientific Research Fund (OTKA), the French Lyon Institute of Origins (LIO), the Belgian Fonds de la Recherche Scientifique (FRS-FNRS), Actions de Recherche Concertées (ARC) and Fonds Wetenschappelijk Onderzoek—Vlaanderen (FWO), Belgium, the Paris Île-de-France Region, the National Research, Development and Innovation Office of Hungary (NKFIH), the National Research Foundation of Korea, the Natural Sciences and Engineering Research Council of Canada (NSERC), the Canadian Foundation for Innovation (CFI), the Brazilian Ministry of Science, Technology, and Innovations, the International Center for Theoretical Physics South American Institute for Fundamental Research (ICTP-SAIFR), the Research Grants Council of Hong Kong, the National Natural Science Foundation of China (NSFC), the Israel Science Foundation (ISF), the US-Israel Binational Science Fund (BSF), the Leverhulme Trust, the Research Corporation, the National Science and Technology Council (NSTC), Taiwan, the United States Department of Energy, and the Kavli Foundation. The authors gratefully acknowledge the support of the NSF, STFC, INFN, and CNRS for the provision of computational resources.

This work was supported by MEXT, the JSPS Leading-edge Research Infrastructure Program, JSPS Grant-in-Aid for Specially Promoted Research 26000005, JSPS Grant-in-Aid for Scientific Research on Innovative Areas 2402: 24103006, 24103005, and 2905: JP17H06358, JP17H06361, and JP17H06364, JSPS Core-to-Core Program A. Advanced Research Networks, JSPS Grants-in-Aid for Scientific Research (S) 17H06133 and 20H05639, JSPS Grant-in-Aid for Transformative Research Areas (A) 20A203: JP20H05854, the joint research program of the Institute for Cosmic Ray Research, University of Tokyo, the National Research Foundation (NRF), the Computing Infrastructure Project of the Global Science experimental Data hub Center (GSDC) at KISTI, the Korea Astronomy and Space Science Institute

(KASI), the Ministry of Science and ICT (MSIT) in Korea, Academia Sinica (AS), the AS Grid Center (ASGC) and the National Science and Technology Council (NSTC) in Taiwan under grants including the Science Vanguard Research Program, the Advanced Technology Center (ATC) of NAOJ, and the Mechanical Engineering Center of KEK.

Additional acknowledgments for support of individual authors may be found in the following document: <https://dcc.ligo.org/LIGO-M2300033/public>. For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) license to any Author Accepted Manuscript version arising. We request that citations to this article use “A. G. Abac et al. (LIGO–Virgo–KAGRA Collaboration), ...” or similar phrasing, depending on journal convention. This article has been assigned document number LIGO-P2500356.

We thank Jing Ming and colleagues for feedback.

Appendix A Details on the BSD Search

A.1. Search Implementation

The input data of the BSD semi-coherent search consists of a complex-valued subsampled time series, the BSD files (O. J. Piccinni et al. 2018). For each target, the detector’s time series is partially corrected for the Doppler effect, assuming the known sky location of the source. This correction is repeated for each 1 Hz subband using its central frequency (as detailed in O. J. Piccinni et al. 2020). From these partially corrected time series, we construct time–frequency peakmaps by extracting local maxima above a given threshold in the equalized spectrum (P. Astone et al. 2005). The peakmaps are then fed into the FH transform, which maps the time–frequency peaks onto the source’s frequency and spin-down plane ($f_0, \dot{f}_0$) at a given reference time t_0 . The second-order spin-down parameter $\ddot{f}_0$ is always assumed equal to zero. The first-order spin-down range is defined as $-f_{\max}/t_{\text{age}} \leq \dot{f} \leq 0.1f_{\max}/t_{\text{age}}$, with $f_{\max}$ the upper frequency of each 10 Hz band. Given that the second-order spin-down is neglected, and we fix a maximum frequency of 600 Hz for all the targets, the minimum age of the source will be 3.8 kyr (see, e.g., Equation (B2) of R. Abbott et al. 2021b).

Each FH map is built with a resolution equal to the template grid spacing as (R. Abbott et al. 2021b):

$$\delta f_{\text{FH}} = \frac{1}{T_{\text{coh}} K_f}, \quad \delta \dot{f}_{\text{FH}} = \frac{1}{T_{\text{coh}} T_{\text{obs}} K_{\dot{f}}}, \quad (\text{A1})$$

where T_{coh} is the coherence time, T_{obs} the observational time span, and $K_f = 10$ and $K_{\dot{f}} = 2$ are the overresolution factors following P. Astone et al. (2014a). Since the coherence time scales with the maximum frequency as $T_{\text{coh}} \propto 1/\sqrt{f_{\max}}$, the grid resolution varies across frequency bands. The coherence time used varies from 17.8 hr at the lowest frequencies to 3.3 hr at 600 Hz. We search both negative and positive spin-down to allow for possible spin-up scenarios. The list of targets and search configurations is listed in Table 2.

As in previous searches, candidates with the highest FH number count are selected from a final map built by summing all monthly FH maps for a given source and frequency range. Candidates are selected independently from each detector, and coincident candidates are then passed to the postprocessing and follow-up steps. We additionally apply a selection threshold on the critical ratio (CR), ρ_{CR} , computed from the

Table 2
Sources Searched in the BSD Analysis (Section 4.2) and the Parameter Space Covered

Source ^a	Minimum t_{age} (kyr)	T_{coh} (hr) (@ 100 Hz)	f (Hz)	$\dot{f}$ (Hz s ⁻¹) (@ 100 Hz)
G65.7+1.2, G189.1+3.0, G266.2-1.2,	3.8	8	[10, 600]	$[-9.18 \times 10^{-10}, 9.18 \times 10^{-11}]$
G353.6-0.7, G93.3+6.9,	...	...	...	...
G291.0-0.1, G330.2+1.0	...	...	...	...
G39.2-0.3	4.7	8	[10, 600]	$[-7.42 \times 10^{-10}, 7.42 \times 10^{-11}]$

Note. The coherence time and the spin-down/up range scale with the maximum frequency in each 10 Hz frequency band. For each source, we report the coherence time, T_{coh} , and spin-down/up range for $\dot{f}$, used for the frequency band [90, 100] Hz where $f_{\text{max}}=100$ Hz.

^a G291.0-0.1 and G330.2+1.0 are two newly added sources in the BSD O4a search, with respect to the BSD O3a search.

FH number count map. For each source, $\rho_{\text{CR,thr}}$ is set to the mean plus one standard deviation of the CR distribution, after excluding candidates overlapping known lines (E. Goetz et al. 2024) or with inconsistent detector significance. The thresholds, which are separately computed for each detector/target, range from 4.74 to 4.87. Detailed information on the postprocessing and follow-up stage can be found in Appendix A.2.

Coincidences between candidates from LIGO Hanford ($f_0^{\text{H}}, \dot{f}_0^{\text{H}}$) and Livingston ($f_0^{\text{L}}, \dot{f}_0^{\text{L}}$) are identified using the metric:

$$d = \sqrt{\left(\frac{f_0^{\text{H}} - f_0^{\text{L}}}{\delta f_{\text{FH}}}\right)^2 + \left(\frac{\dot{f}_0^{\text{H}} - \dot{f}_0^{\text{L}}}{\delta \dot{f}_{\text{FH}}}\right)^2}, \quad (\text{A2})$$

with $d_{\text{thr}} = 4$ as the coincidence threshold (see P. Astone et al. 2014a). All surviving candidates undergo a postprocessing and follow-up step.

A.2. Postprocessing and Follow-up

After applying the known-lines veto, the consistency veto, and a CR threshold selection, ~ 50 candidates remain for each target (exact numbers are reported in Table 3).

For each of these candidates, we apply a veto based on the 5-vector statistics (P. Astone et al. 2010, 2014b), which provides an alternative estimate of the signal-to-noise ratio and CR after applying Doppler and spin-down corrections. These corrections are implemented through a heterodyne phase demodulation based on the candidate's estimated frequency

f_0 and spin-down $\dot{f}_0$. The corrected time series is obtained by multiplying the original data by an exponential factor $\exp[-i\Phi(t)]$, where $\Phi(t)$ describes the assumed phase evolution of the signal. A true astrophysical signal is expected to show an increase in significance after correction, scaling roughly with the fourth root of the coherence time. To test this, we compute the 5-vector statistics using two coherence times: $T_{\text{sid}} = 86164.0905$ s (one sidereal day), and $T_{4\text{sid}} = 4T_{\text{sid}}$. For each candidate, we compute their significance ρ_i^{C} and ρ_i^{NC} , where $i = \text{CR}$, snr denotes either the CR or the signal-to-noise ratio, and superscripts “C” and “NC” refer to corrected and uncorrected data, respectively. We retain only those candidates for which the corrected significance improves, i.e., $\rho_i^{\text{C}} - 0.95\rho_i^{\text{NC}} > 0$, ensuring a minimum 5% tolerance. Simulation studies confirm that this veto has a false dismissal probability below 10% as long as a tolerance of 5% is assumed. At this stage, ~ 10 candidates per target survived, and we applied a second consistency veto using the 5-vector statistics, which further discarded some candidates.

Among the candidates that survived these vetos, we applied the cumulative-significance veto, as described in Section A.1.5 of the O3a SNR paper (R. Abbott et al. 2021b). This procedure further reduced the candidate list, eliminating the majority of remaining outliers and leaving only a few exceptions (reported in Table 4).

To veto the candidates based on the cumulative-significance veto, we examined the trends of the CR, signal-to-noise ratio, and the 5-vector S -statistics for each candidate, computed both in the corrected and uncorrected cases. Candidates that exhibited a clear reduction in significance after correction

Table 3
Number of Candidates Surviving the Different Selection Criteria of the BSD Postprocessing Stage

First Level Selection	G39.2	G65.7	G93.3	G189.1	G266.2	G291.0	G330.2	G353.6
H	389064	468268	468093	467946	468036	468170	468171	467863
L	388524	467260	467250	467420	467375	467408	467425	467597
Coincidences	2276	2728	2716	2617	2688	2634	2613	2700
Veto Stage								
Known lines	1919	2315	2310	2204	2268	2260	2241	2291
CR Consistency	1916	2311	2310	2203	2264	2255	2236	2288
CR Threshold	39	55	61	44	47	42	44	36
5-vector veto	4	5	15	10	10	10	9	9
Second consistency (5-vector)	3	4	7	9	6	9	7	8
Cumulative-significance veto	1	1	...	...	3	1	1	1

Note. The top part of the table reports the first level of candidates selected from each detector (H or L) and those found in coincidence between the two. This is followed by the number of candidates surviving the different veto steps.

Table 4

Surviving Candidates from the BSD Pipeline after Vetos in Appendix A.2

Source	f_0 (Hz)	$\dot{f}_0$ (Hz s ⁻¹)	ρ_{CR} (FH)
G39.2−0.3	582.6910443	$-4.07153112 \times 10^{-10}$	5.39
G65.7+1.2	155.4805487	$-1.00577202 \times 10^{-9}$	5.28
G266.2−1.2	212.3495104	$-1.61918580 \times 10^{-9}$	5.34
...	365.5448170	$-3.8870222216 \times 10^{-10}$	4.97
...	450.3867653	$-3.4974123560 \times 10^{-9}$	5.53
G291.0−0.1	563.7719252	$-1.899359832 \times 10^{-9}$	5.08
G330.2+1.0	108.3626199	$-7.987104272 \times 10^{-11}$	4.98
G353.6−0.7	30.00352801	$-3.263776543 \times 10^{-10}$	6.73

Note. The candidates investigated were excluded after comparing the corrected and uncorrected peakmap and the spectra around the candidate in each detector. For each source with a surviving associated candidate, we report the original mean CR from the FH map, the candidate frequency f_0 , and the spin-down $\dot{f}_0$ at the time of the coincidences.

were removed, as were those with very low 5-vector CR values, specifically those below 3, corresponding to a false alarm probability bigger than $\sim 1\%$. In a few cases, candidates met the first two criteria but not the third; these were tagged as low-significance outliers and retained for completeness, though flagged accordingly. The list of candidates that survived up to the cumulative-significance veto step is reported in Table 4.

Following this quantitative filtering, we performed a series of visual inspections to assess the remaining candidates. This included examination of the peakmap, both before and after correction, along with the corresponding histogram, as well as inspection of the power spectral density with and without correction. These visual checks provided an additional qualitative assessment of candidate behavior under correction and helped further refine the final selection. None of the candidates was confirmed as astrophysical.

A.3. Upper Limits

We set constraints on the GW strain amplitude using a well-established procedure previously applied in O. J. Piccinni et al. (2020) and R. Abbott et al. (2022b). The 95% confidence-level upper-limit estimate is given by (P. Astone et al. 2014a; C. Palomba 2025):

$$h_0^{95\%} \approx \frac{\mathcal{B}}{N^{1/4}} \sqrt{\frac{S_n(f)}{T_{\text{coh}}}} \sqrt{\rho_{\text{CR,max}} - \sqrt{2} \operatorname{erfc}^{-1}(2\Gamma)}, \quad (\text{A3})$$

where $\rho_{\text{CR,max}}$ is the maximum CR in each band. We indicate with N the effective number of FFTs used for the search and with $\mathcal{B}$ a parameter dependent on the source sky location and the averaged signal polarization. The sensitivity curve, i.e., the minimum detectable strain, is obtained by substituting $\rho_{\text{CR,max}}$ in Equation (A3) with the CR threshold used for candidate selection $\rho_{\text{CR,thr}}$. The resulting $h_0^{95\%}(f)$ upper limits and sensitivity curves are shown in Figure 8. We corroborate the results by computing the 95% confidence-level upper limits on h_0 using artificial signals injected in randomly selected 1 Hz subbands: six subbands in the 10–600 Hz frequency range for all the targets. For each subband, $h_0^{95\%}$ is derived using a frequentist approach by injecting 50 synthetic signals per amplitude level. The injections are performed assuming the




Figure 8. Estimated 95% confidence-level upper limit obtained from the BSD search.

known sky position of each target. Spin-down, inclination angle ($\cos \iota$), and polarization angle (ψ) are drawn from uniform distributions. We repeat the injection-and-recovery process using different values of h_0 . The detection efficiency for each amplitude is then computed as the fraction of successfully recovered signals. A sigmoidal fit to the detection efficiency curve yields the $h_0^{95\%}$ value corresponding to 95% detection efficiency.

To obtain a more robust estimate, the factor $\mathcal{B}$ has been calculated as the average of the empirically derived prefactors from the injections. This approach ensures that the results remain robust to potential deviations. The final upper-limit curve is determined by selecting the more conservative (larger) value between the curves of each detector.

Appendix B Details on the PyStoch Search

B.1. Search Implementation

In this search, PyStoch estimates the optimal signal-to-noise ratio as a function of the frequency for a specific sky location using the folded datasets as input. The targets and parameter space investigated by PyStoch are listed in Table 5.

The frequencies corresponding to instrumental lines are excluded from this computation and are reported in a notch list (A. G. Abac et al. 2025b). The signal-to-noise ratio for each frequency bin is defined as $\frac{Y}{\sigma_Y}$, i.e., the ratio between the cross-correlation statistics Y and the corresponding standard deviation σ_Y (A. Ain et al. 2018).

Since GW radiometer searches are unmodeled and typically do not incorporate spin evolution, targeted narrow-band stochastic GW background analyses often employ a running average to combine the signal-to-noise ratio across adjacent frequency bins. This approach helps account for potential frequency drifts of the source over the observation period (B. P. Abbott et al. 2017, 2019c; R. Abbott et al. 2021c),

Table 5
PyStoch Search Setup for Cas A, G347.3–0.5, Vela Jr., and SNR 1987A, Assuming a Given Source Age

Source	t_{age} (yr)	$ \Delta f_0 $ (Hz)	$ \dot{f}_0 $ (Hz s ⁻¹)	$\ddot{f}_0$ (Hz s ⁻²)
G111.7–2.1/Cas A	330	[0.007, 3.677]	$[3.48 \times 10^{-10}, 1.80 \times 10^{-7}]$	$[4.23 \times 10^{-20}, 3.75 \times 10^{-17}]$
G347.3–0.5/RX J1713.7–3946	1600	[0.001, 0.698]	$[6.59 \times 10^{-11}, 3.41 \times 10^{-8}]$	$[1.52 \times 10^{-21}, 1.35 \times 10^{-18}]$
G266.2–1.2/Vela Jr.	700	[0.003, 1.590]	$[1.51 \times 10^{-10}, 7.82 \times 10^{-8}]$	$[7.98 \times 10^{-21}, 7.08 \times 10^{-18}]$
SNR 1987A	38	[0.058, 7.460]	$[2.85 \times 10^{-9}, 3.68 \times 10^{-7}]$	$[2.83 \times 10^{-18}, 3.93 \times 10^{-16}]$

Note. The corresponding ranges of $|\Delta f_0|$, $\dot{f}_0$, $\ddot{f}_0$ are also reported.

reducing the likelihood that signal power spreads across multiple bins and is therefore lost. To account for the frequency drift resulting from NS spin-down, we estimate the expected signal broadening, Δf_0 , for each target as:

$$\Delta f_0 = \underbrace{-\frac{f_0}{t_{\text{age}}(n-1)}}_{\dot{f}_0} T_{\text{obs}} + \frac{1}{2} \underbrace{\frac{n}{f_0} \left(\frac{f_0}{t_{\text{age}}(n-1)} \right)^2}_{\ddot{f}_0} T_{\text{obs}}^2, \quad (\text{B1})$$

where T_{obs} denotes the duration of the O4a observing run. This spread corresponds to the optimal frequency resolution expected for a given signal.

For the first time, we implement PyStoch with a customized binning strategy tailored to signals from young NSs in SNRs. Starting from the default 1/32 Hz resolution, adjacent frequency bins are adaptively merged to match the expected signal broadening Δf_0 for each target (see Equation (B1)), aiming to keep any potential CW signal fully contained within at least one frequency bin (despite possible frequency drift). To further improve sensitivity and reduce the risk of signal power being diluted across multiple bins, we apply a sliding-window combination of adjacent bins. This approach, repeated for multiple window sizes, enhances coherence detection across a range of frequency scales while preserving the original frequency grid for interpretation. Simulations presented in C. Salvatore et al. (2025) indicate that combining bins beyond 1.5 Hz introduces noise clustering effects and reduces performance, establishing an empirical maximum width for effective binning.

Candidates are selected based on the simultaneous fulfillment of two criteria: (i) a signal-to-noise ratio exceeding 4.5, which corresponds to a p -value of 10 %, and (ii) a bin width consistent with the expected signal broadening Δf_0 , computed from Equation (B1) based on the braking indices compatible with the candidate frequency.³²⁷

The statistical significance of the outliers is quantified via p -values, representing the probability that a measured

signal-to-noise ratio arises from random noise fluctuations. These values are estimated through Monte Carlo simulations using Gaussian realizations consistent with the noise characteristics of the data.

For each frequency bin, simulated distributions of the detection statistic are generated and processed through the same bin combination strategy described previously. The maximum signal-to-noise ratio from each realization is recorded, and the resulting ensemble is used to construct an empirical mapping between signal-to-noise ratio values and their corresponding p -values via linear interpolation. Frequency bins with p -values $\leq 10\%$ are considered statistically significant, as they are unlikely to arise from random noise fluctuations.

As a result, only a limited number of promising candidates were identified for Vela Jr. and Cas A, while no significant candidate was observed for SNR 1987A and G347.3–0.5. These candidates were followed up using two independent methods: the FH transform and the 5-vector resampling technique (see Section B.2).

B.2. Postprocessing and Follow-up

The list of candidates passed to the follow-up stage is reported in Table 6.

We stress that the selected candidates were subjected to follow-up analyses using two dedicated CW search pipelines: the FH method (P. Astone et al. 2014a) and the 5-vector resampling method (F. Amicucci et al. 2025; A. Singhal et al. 2019). No surviving candidates were confirmed by the 5-vector resampling search. For Cas A, the FH pipeline identified coincident candidates, which are defined as those lying within a distance of less than 3 in the search parameter space, when short coherence times (64 and 128 s) were used. However, these coincidences did not persist at longer coherence times (512 s), suggesting they were likely due to noise or transient features.³²⁸ Finally, neither of the two follow-up methods identified statistically significant candidates for Vela Jr.

³²⁷ Assume we observe an outlier at a frequency f_0 with a frequency distribution Δf_0 (i.e., by combining a certain number of frequency bins). If there exists an index $n \in [2, 7]$ such that, for a given candidate frequency f_0 , the corresponding value of Δf_0 (see Equation (B1)) matches the expected prediction, then the pair (f_0, n) is considered compatible with the theoretical frequency distribution.

³²⁸ The FH follow-up analysis considered only the first-order spin-down parameter, with the second-order term being neglected. While this choice results in a reduction in the detection sensitivity (i.e., CR) for the optimal FFT length, it was a conscious choice made to simplify the initial analysis. A fully optimized version of the code, capable of incorporating and accurately handling the second-order spin-down term, will be available in the near future.

Table 6
Candidates Identified for Each Source by PyStoch

Source	p -value	f_0 (Hz)	signal-to-noise ratio	Δf_0 (Hz)	n	$\dot{f}_0$ (Hz s ⁻¹)	$\ddot{f}_0$ (Hz s ⁻²)
G266.2–1.2/Vela Jr.	0.09	1168.29688	4.63	0.34375	4.2	-1.68×10^{-8}	1.01×10^{-18}
G111.7–2.1/Cas A	0.08	788.51563	4.65	0.59375	3.9	-2.91×10^{-8}	4.20×10^{-18}
G111.7–2.1/Cas A	0.07	788.51563	4.59	0.71875	3.4	-3.52×10^{-8}	5.35×10^{-18}

Note. The parameters are listed from left to right as follows: p -value, reference frequency at the start of O4a (f_0), signal-to-noise ratio, candidate frequency distribution (Δf_0), braking index (n), and spin-down parameters evaluated at the reference time (set at the O4a start, GPS time 1368980142 s).

B.3. Upper Limits

In the absence of a significant detection, upper limits on the CW strain amplitude are placed. The computation of upper limits is performed within a Bayesian framework, incorporating prior distributions over relevant source parameters such as the inclination angle, polarization angle, and calibration uncertainty. The final result is the marginalized posterior distribution for h_0 , from which the upper limit at a given confidence level (e.g., 95%) is extracted.

Due to the computational cost associated with evaluating the full marginalization for each frequency bin, an interpolation-

based approach is adopted. Upper limits are precomputed for a range of representative signal-to-noise ratio values and both circular and generic polarizations. The ratio between these cases, which depends only on the signal-to-noise ratio, allows for a rapid estimation of the marginalized upper limits across the entire frequency band. This procedure is validated against full posterior computations for a subset of bins. Results for each target are reported in Figure 9, while the best results are listed in Table 7.

Further technical and mathematical details of the p -value estimation and upper limits computation can be found in the dedicated stochastic analysis paper (A. G. Abac et al. 2025c).




Figure 9. Best (1/32 Hz, left) and worst (47/32 Hz, right) 95% confidence-level upper limit between 20 and 1726 Hz, computed with PyStoch for each target: SNR 1987A (orange), Vela Jr. (blue), G347.3–0.5 (green), Cas A (red).

Table 7
Best 95% Confidence-level Upper-limit Set by PyStoch with Corresponding Frequency for Each SNR

Source	Best $h_0^{95\%}$ ($\times 10^{-26}$)	f (Hz)
G111.7–2.1/Cas A	9.21	268.69
G347.3–0.5	9.78	332.53
G266.2–1.2/Vela Jr.	9.02	343.59
SNR 1987A	9.83	323.16

Appendix C

Details on SHV Search

C.1. Search Implementation

In this Appendix, we summarize the SHV pipeline used in Section 4.4. We refer the reader to Y. Bu & A. Melatos (2025), which includes the technical and full details of this method and a major update to the SHV pipeline since the O3a search (R. Abbott et al. 2021b).

We follow the Viterbi HMM formalism used in previous SHV searches (e.g., S. Suvorova et al. 2016; L. Sun et al. 2018). We consider a set of N_T observations at discrete times $t_0, t_1, \dots, t_{N_T-1}$ and N_Q possible hidden states $\{q_1, \dots, q_{N_Q}\}$, where each hidden state corresponds to a discrete GW frequency bin. The corresponding sequence of discrete states $\{q(t_0), q(t_1), \dots, q(t_{N_T-1})\}$, through which the system progresses, forms a Markov chain. The transition probability from time t_k to t_{k+1} is defined as $A_{q_k q_j} = P[q(t_{k+1}) = q_j | q(t_k) = q_k]$. We adopt a simple random-walk transition model in frequency, allowing the signal to remain in the same bin or wander up or down by one bin, during each time step with equal probability. We assume a uniform prior over the initial hidden state. The emission probability is computed from the coherent $\mathcal{F}$ -statistic in each segment of duration T_{coh} . Given the observation sequence, the most likely hidden-state sequence can be inferred by maximizing the most probable hidden-state path.

C.1.1. Parameter Choice

Here, we outline the recipe for selecting the searchable parameter domain in the SHV pipeline. The list of targets and the parameter domain for each target are listed in Table 8. For each target with distance $D_{\text{min}} \leq D \leq D_{\text{max}}$, age $t_{\text{age,min}} \leq t_{\text{age}} \leq t_{\text{age,max}}$, and braking index $n_{\text{min}} \leq n \leq n_{\text{max}}$, we define the search parameters T_{coh} and f via four criteria:

1. The frequency bin width δf is sufficiently small, such that f remains constant during each coherent segment, with

$$\delta f = \frac{1}{2T_{\text{coh}}} > |\dot{f}(f, n)| T_{\text{coh}}. \quad (\text{C1})$$

2. The $\mathcal{F}$ -statistic software ingests at least two SFTs (LIGO Scientific Collaboration 2022), with $T_{\text{coh}} \geq 2T_{\text{SFT}}$.
3. The signal amplitude h_0^{sig} exceeds the minimum detectable strain at 95% confidence for a semi-coherent search h_0^{est} , with $h_0^{\text{sig}} > h_0^{\text{est}}$.
4. The maximum strain implied by energy conservation, h_0^{max} , exceeds the signal amplitude, with $h_0^{\text{max}} > h_0^{\text{sig}}$.

Using Equation (1) to express $\dot{f}$ and adopting the standard late-time approximation

$$\dot{f}(t_{\text{age}}) \approx -\frac{f(t_{\text{age}})}{(n-1)t_{\text{age}}}, \quad (\text{C2})$$

the first criterion yields an upper bound on the coherence time:

$$T_{\text{coh}} < \left[\frac{2f}{(n-1)t_{\text{age}}} \right]^{-1/2}. \quad (\text{C3})$$

The second criterion provides a lower bound on T_{coh} . The third and fourth criteria are simultaneously satisfied when $h_0^{\text{max}} > h_0^{\text{est}}$, which gives a lower bound of the form

$$T_{\text{coh}} > (2.27 \times 10^{-24} \text{ kpc kyr}^2)^{-4} D^4 t_{\text{age}}^2 (n-1)^2 \Theta^4 S_n(f)^2 T_{\text{obs}}^{-1}, \quad (\text{C4})$$

where $\Theta = 35$ is an empirical statistical factor (K. Wette et al. 2008). The resulting overlap of constraints defines the searchable (f, T_{coh}) domain. For each frequency within this domain, we choose T_{coh} to maximize sensitivity, following the prescription illustrated in Figure 10 and summarized in Table 8.

Table 8
Sources in the SHV Analysis (Section 4.4) with Search Bands, Coherence Times T_{coh} , and Frequency Derivative Ranges $\dot{f}$

Source	f (Hz)	T_{coh} (hr)	T_{coh} (hr) (@ 100 Hz)	$\dot{f}$ (Hz s ⁻¹) (@ 100 Hz)
G18.9–1.1	[29, 2000]	[4.7, 39.2]	21.1	$[-8.67 \times 10^{-11}, 8.67 \times 10^{-11}]$
G39.2–0.3/3C 396	[55, 736]	[7.1, 26.3]	19.5	$[-1.01 \times 10^{-10}, 1.01 \times 10^{-10}]$
G65.7+1.2/DA 495	[21, 2000]	[23.7, 228.7]	106.4	$[-3.41 \times 10^{-12}, 3.41 \times 10^{-12}]$
G93.3+6.9/DA 530	[39, 1379]	[6.0, 36.3]	22.6	$[-7.55 \times 10^{-11}, 7.55 \times 10^{-11}]$
G189.1+3.0/IC 443	[21, 2000]	[10.4, 102.2]	46.8	$[-1.76 \times 10^{-11}, 1.76 \times 10^{-11}]$
G266.2–1.2/Vela Jr.	[20, 2000]	[4.3, 43.2]	19.3	$[-1.04 \times 10^{-10}, 1.04 \times 10^{-10}]$
G353.6–0.7	[30, 2000]	[9.9, 80.7]	44.4	$[-1.96 \times 10^{-11}, 1.96 \times 10^{-11}]$
G1.9+0.3	[32, 1607]	[1.0, 7.1]	4.0	$[-2.41 \times 10^{-9}, 2.41 \times 10^{-9}]$
G15.9+0.2	[34, 1915]	[3.3, 25.4]	14.8	$[-1.76 \times 10^{-10}, 1.76 \times 10^{-10}]$
G111.7–2.1/Cas A	[28, 2000]	[1.1, 9.5]	5.1	$[-1.48 \times 10^{-9}, 1.48 \times 10^{-9}]$
G291.0–0.1/MSH 11-62	[30, 2000]	[6.0, 48.9]	27.0	$[-5.29 \times 10^{-11}, 5.29 \times 10^{-11}]$
G330.2+1.0	[31, 2000]	[7.4, 59.2]	33.1	$[-3.52 \times 10^{-11}, 3.52 \times 10^{-11}]$
G347.3–0.5	[21, 2000]	[2.4, 24.2]	11.1	$[-3.13 \times 10^{-10}, 3.13 \times 10^{-10}]$
G350.1–0.3	[32, 1999]	[1.5, 12.7]	7.2	$[-7.44 \times 10^{-10}, 7.44 \times 10^{-10}]$

Note. The corresponding frequency derivative ranges are computed as $1/(2T_{\text{coh}}^2)$ using the @100 Hz coherence time (converted to seconds). The age and distance of each target are listed in Table 1. SHV does not specifically search the first-order spin-downs. The estimated spin-down reported is inferred from T_{coh} .




Figure 10. T_{coh} and f values that satisfy the four criteria defined in Appendix C.1.1. The shaded pink region satisfies criterion one, i.e., Equation (C3). The dotted gray region satisfies criterion two. The shaded blue region satisfies criteria three and four simultaneously. The purple domain indicates where all four criteria are satisfied, designated as the search region. For each f in the purple region, we choose T_{coh} according to the pink curve to maximize the search sensitivity.

C.1.2. Detection Threshold

We set detection thresholds by choosing an overall false alarm probability, $\alpha = 0.30$, for the ensemble of 14 targets. Across all targets, with each labeled by i , there are $N_{\text{band},i}$ subbands. Hence, the equivalent false alarm probability per subband for target i is $\alpha_{\text{band},i} = 1 - (1 - \alpha_{\text{target}})^{1/N_{\text{band},i}}$.

Following previous SHV searches, we determine the threshold for the Viterbi log-likelihood statistic $\mathcal{L}$ via Monte Carlo simulations in Gaussian noise with the O4a duty cycle. The likelihood threshold $\mathcal{L}_{\text{th}}$ is defined as the $1 - \alpha_{\text{band},i}$ th percentile of the $\mathcal{L}$ distribution. The resulting thresholds are well described by a power-law dependence on coherence time, $\mathcal{L}_{\text{th}} = C_{\text{band}} T_{\text{coh}}^{p_{\text{band}}}$ which we use to evaluate thresholds across the search space. Any path with $\mathcal{L} > \mathcal{L}_{\text{th}}$ is followed up using the procedure in Appendix C.2. For a detailed

analysis of the individual targets, see Y. Bu & A. Melatos (2025).

C.2. Postprocessing and Follow-up

After conducting the SHV search, we apply a standard follow-up procedure comprising a known-line veto, an interferometer veto, a sky-position veto, and a subband veto (see R. Abbott et al. 2021b). The number of candidates remaining after each stage is summarized in Table 9. A total of six candidates survive all automated vetoes; their parameters are listed in Table 10.

We examine these survivors using single-interferometer statistics and off-target sky maps. None displays the expected on-target morphology or cross-detector consistency of a true signal. We therefore conclude that all six outliers are consistent with instrumental artifacts and that no SHV candidate survives follow-up.

Table 9
Survivor Counts for Each Target of SHV with $\mathcal{L} > \mathcal{L}_{\text{th}}$ and at the End of All Four Postprocessing Stages

Veto Stage	G111	G65	G1	G291	G266	G189	G330	G353	G347	G18	G93	G350	G15	G39
Above threshold	435	368	317	316	313	294	294	293	265	264	261	258	213	109
Known-lines veto	31	60	27	29	30	33	37	28	24	22	25	21	25	8
Interferometer veto	1	14	1	1	3	4	2	1	2	2	2	1	1	...
Sky position veto	1	14	1	1	3	4	1	1	1	2	2	...	1	...
Subband veto	1	3	1	...	1	...	...	...	...	...	...	...	...	...

Note. We shorten the names of the targets for simplicity, as G347 for G347.3–0.5, etc. The full names are reported in Table 1.

Table 10
Candidates Surviving All Vetos from the SHV Search

Source	f_0 (Hz)	$\mathcal{L}_{\text{th}}$	$\mathcal{L}$	$\mathcal{L}_{\text{H1}}$	$\mathcal{L}_{\text{L1}}$	$\mathcal{L}_{\text{off}} - \text{target}$
G111.7–2.1/Cas A	1169.00	27861.43	35500.86	36564.78	42308.51	35078.42
G266.2–1.2/Vela Jr.	213.31	3177.03	3777.69	3339.62	5149.54	8557.62
G65.7+1.2/DA 495	108.76	418.83	430.26	442.83	419.67	432.77
G65.7+1.2/DA 495	137.83	472.39	483.77	474.44	477.31	478.21
G65.7+1.2/DA 495	139.95	474.62	491.81	568.67	484.56	545.42
G1.9+0.3	1161.08	34800.84	45007.50	54777.89	35121.50	45144.90

Note. Each row lists a candidate’s target name, start frequency f_0 of the Viterbi path, threshold log-likelihood $\mathcal{L}_{\text{th}}$, combined detection statistic $\mathcal{L}$, single-interferometer log-likelihoods $\mathcal{L}_{\text{H1}}$ and $\mathcal{L}_{\text{L1}}$ from the interferometer veto, and off-target log-likelihood $\mathcal{L}_{\text{off}} - \text{target}$ from the sky-position veto.

C.3. Upper Limits

We determine the upper limits using all targets. For each target, we choose two subbands within the original search that have no known lines and return fewer than two unique paths with $\mathcal{L} > \mathcal{L}_{\text{th}}$, with $f \simeq f_{\text{min}}$ and $f \simeq f_{\text{max}}$, where f_{min} and f_{max} denote the minimum and maximum frequency of the target’s search band. The chosen frequencies are shown in Y. Bu & A. Melatos (2025), along with T_{coh} and $\mathcal{L}_{\text{th}}$ from the original search.

By recording the fraction of injected signals recovered (i.e., those with $\mathcal{L} > \mathcal{L}_{\text{th}}$) for each h_0 , we estimate the detection efficiency $\epsilon(h_0)$. We then use logistic regression to fit a sigmoidal model to $\epsilon(h_0)$, and solve $\epsilon(h_0^{95\%}) = 0.95$ to obtain the 95% confidence upper limit $h_0^{95\%}$ in the given subband. We calculate $a = h_0^{95\%}/h_0^{\text{est}}$ for each subband. We assume that the sensitivity only depends on T_{coh} and neglect other factors such as sky position. To obtain the upper limit across the full search band, we define $h_0^{95\%}(T_{\text{coh}}) = a(T_{\text{coh}}) h_0^{\text{est}}$. To verify that the full-band scaled upper limit $h_{0,\text{UL}}^{95\%}$ is reliable, we compare it with the upper limit obtained from injections $h_{0,\text{inj}}^{95\%}$. For each target and each injection, we compute the ratio $R \equiv h_{0,\text{inj}}^{95\%}/h_{0,\text{UL}}^{95\%}$. For each target, we then define a rescaling constant per target $k = \max(R)$ over injections for that target. If $k > 1.1$, the entire upper-limit curve for that target is rescaled as $h_{0,\text{UL}}^{95\%} \rightarrow k h_{0,\text{UL}}^{95\%}$. See Y. Bu & A. Melatos (2025) for the values of k for each target. We present the upper-limit curve for all targets in Figure 11.

The SHV pipeline models the frequency evolution as a random walk, which is flexible enough to accommodate secular spin down and stochastic spin wandering. Consequently, the resulting upper limits are not directly comparable to the pipelines in this paper, which model the frequency evolution as a Taylor series.




Figure 11. Upper-limit curves for all targets searched by the SHV pipeline. The scatter points show the sensitivity estimate $h_0^{95\%}(T_{\text{coh}}) = a(T_{\text{coh}})h_0^{\text{est}}$ across the searched frequency band.

Appendix D

Details on the DHV Search

D.1. Search Implementation

The HMM structure used in the dual-harmonic search largely follows the standard formulation described in Appendix C, with specific modifications outlined in this Appendix. The list of targets and parameter space covered is listed in Table 11.

Table 11

Targets Searched in the DHV Analysis (Section 4.5) and the Parameter Space Covered

Source	Minimum t_{age} (kyr)	T_{coh} (hr)	f_* (Hz)	$\dot{f}_*$ (Hz s ⁻¹)
G353.6−0.7	20	12	[50, 338]	[−1.34 × 10 ^{−10} , 0]
G189.1+3.0	...	...	...	...
G65.7+1.2	...	...	...	...
G266.2−1.2	5	9	[50, 150]	[−2.38 × 10 ^{−10} , 0]
G93.3+6.9	...	...	...	...
G18.9−1.1	...	...	...	...
G39.2−0.3	...	...	...	...

In this analysis, two frequency components are tracked simultaneously. The signal power in the i th frequency bin, f_i , is computed by the sum of two $\mathcal{F}$ -statistic components computed in dual-harmonic frequency bins, $\mathcal{F}_1(f_i) + \mathcal{F}_2(2f_i)$. The GW signal for each frequency component is assumed to be monochromatic over T_{coh} .

We choose a coherence-time interval T_{coh} such that the following condition is satisfied:

$$\left| \int_t^{t+T_{\text{coh}}} dt' \dot{f}_*(t') \right| < \delta f_1, \quad (\text{D1})$$

where $\delta f_1 = 1/(4T_{\text{coh}})$ is the frequency bin width in the $\mathcal{F}_1$ statistic. For the $\mathcal{F}_2$ statistic, we adopt a bin width of $\delta f_2 = 2\delta f_1 = 1/(2T_{\text{coh}})$. The choice of bin widths ensures that the signal drifts by no more than one bin in both $\mathcal{F}_1$ and $\mathcal{F}_2$ across adjacent coherent segments. Thus, the $\dot{f}_*$ range covered in the search (listed in Table 11) depends on T_{coh} , i.e., $|\dot{f}_*| \in [0, 1/(4T_{\text{coh}}^2)]$.

The emission probability for each time step of length T_{coh} is computed as (P. Jaranowski et al. 1998; L. Sun et al. 2019):

$$\begin{aligned} \ln L_{o(t_k)q(t_k)} &= \ln P[o(t_k) | f_i \leq f_*(t_k) < f_i + \delta f_1] \\ &= \mathcal{F}_1(f_i) + \mathcal{F}_2(2f_i), \end{aligned} \quad (\text{D2})$$

where f_i denotes the frequency in the i th bin.

In this work, we assume that the frequency evolution is dominated by the secular spin-down of the NS. Under this assumption, the transition probability matrix $A_{q_i q_i}$ reduces

to the form (L. Sun et al. 2019):

$$A_{q_i \rightarrow q_i} = A_{q_i q_i} = \frac{1}{2}, \quad (\text{D3})$$

with all other entries set to zero. A uniform prior distribution is applied to the initial state, $\Pi[q(t_0)] = N_Q^{-1}$.

The Viterbi score S is used as the detection statistic in this pipeline, representing the significance of the optimal Viterbi path in each subband relative to all other paths at the final step of the tracking (L. Sun et al. 2019).

Two different coherence times of $T_{\text{coh}} = 12$ hr and 9 hr are selected for three older sources and four younger sources, respectively (see Table 11). For parallelized computing, the full frequency band is divided into 1 Hz (1.5 Hz) subbands for $T_{\text{coh}} = 12$ hr ($T_{\text{coh}} = 9$ hr).³²⁹

We set detection thresholds corresponding to 1% false alarm probability at $S_{\text{th}} = 5.26$ for $T_{\text{coh}} = 12$ hr and $S_{\text{th}} = 5.15$ for $T_{\text{coh}} = 9$ hr, based on Monte Carlo simulations in Gaussian noise and validated using real O4a data.

After conducting the search for each target and identifying all the above-threshold candidates, we apply a standard follow-up procedure with a series of vetoes described in Section D.2.

D.2. Postprocessing and Follow-up

After a standard procedure of the known-line veto, interferometer veto, Doppler-shift veto, and sky-position veto (see details in R. Abbott et al. 2021b), 16 candidates remain, listed in Table 12. We pass these 16 survivors through the final stage of a coherence-time veto, and all of them are rejected. With the increased T_{coh} listed in the sixth column, all except one of the new scores fall below the original score. To be more conservative, we further check the two candidates with a new score below the original score but above S_{th} , marked by “*” and “†.” With the increased T_{coh} , a lower mean spin-down rate is returned for both candidates, allowing us to further increase the coherence time to $T_{\text{coh}} = 20$ hr. We find that the score further drops for both candidates. We conclude that they do not survive further scrutiny. One candidate, marked by “†,” returns with a higher score of 14.38, but the optimal path (at ≈ 260.79 Hz, intersecting a known instrumental line at Hanford) is distinct from the original one. By shifting the search subband to 259.7–260.7 Hz with $T_{\text{coh}} = 15$ hr, the score drops below S_{th} , and the new optimal path does not match the original. Thus, none of the remaining candidates survives the final check.

³²⁹ The detection statistic used in this analysis requires a much larger number of frequency bins in each subband than the total number of tracking steps (i.e., $2BT_{\text{coh}} \gg T_{\text{obs}}/T_{\text{coh}}$ with B being the bandwidth), such that the paths ending in different bins are mostly independent. Therefore, we extend the subband to 1.5 Hz for $T_{\text{coh}} = 9$ hr to satisfy this requirement.

Table 12
Final Candidates from the DHV Pipeline and the Coherence-time Veto Results (All Vetted)

Source	S	$2f_{\star\text{start}}$ (Hz)	$2f_{\star\text{end}}$ (Hz)	Mean $2f_{\star}$ (10^{-10} Hz s $^{-1}$)	New T_{coh} (hr)	New S	Paths Overlap?
G353.6–0.7	6.68	184.9413888893380	184.9388657411880	–1.23198	15	6.04 *	✓
	5.58	346.2777314809320	346.2748495364930	–1.40717	15	4.47	×
	5.32	586.6338310172640	586.6316666654170	–1.05679	15	5.01	✓
	5.59	641.9995486091329	641.9969907387680	–1.24894	15	4.52	×
G189.1+3.0	6.10	260.6886458319710	260.6859606467910	–1.31110	15	14.38 †	×
	6.16	272.0795138887320	272.0770949072550	–1.18112	15	5.77 ‡	✓
	5.53	382.4306828695180	382.4274074065610	–1.59932	15	4.49	×
	5.62	405.6810763875410	405.6783564801390	–1.32805	15	4.49	✓
	5.28	431.0656134257960	431.0625810183950	–1.48064	15	5.02	✓
	6.37	498.999999980210	498.9982060165430	–0.87595	15	4.47	×
	5.41	608.6361921283710	608.6331018505990	–1.50890	15	4.86	×
G65.7+1.2	5.90	396.2719791661280	396.2690393513190	–1.43543	15	4.17	×
	5.38	412.6593865727690	412.6568055542560	–1.26024	15	4.13	×
	5.56	441.6801273134690	441.6772337949560	–1.41282	15	4.00	×
	5.27	575.3298726845320	575.3267361104640	–1.53150	15	3.97	×
G18.9–1.1	5.28	117.3951234567360	117.3906481480940	–2.18517	12	3.94	×

Note. The first five columns list the source name, original score, estimated start and end $2f_{\star}$, and the mean $2f_{\star}$. Column 6 lists the new T_{coh} used in the coherence-time veto. The last two columns show the follow-up results: the new score obtained by increasing T_{coh} , and whether the new optimal path overlaps the original candidate path. The three new scores marked by “*”, “†”, and “‡” are above the original thresholds set in the search. Note that the pipeline returns estimated frequencies and spin-down rates corresponding to the $2f_{\star}$ component.

D.3. Upper Limits

In Figure 12, we show the upper limits for the DHV targets.

First, we derive the frequentist upper limit through injections in one sample subband, starting from 112.5 Hz for $f_{\star}$ and 225 Hz for $2f_{\star}$. A set of 200 synthetic signals is injected into the O4a data at random sky positions in that subband with a fixed h_0 . The corresponding detection rate is calculated. This process is repeated with different h_0 values with a step size of 1×10^{-26} . With all the injected h_0 values and the corresponding detection rates, $h_0^{95\%}$ is obtained through a sigmoidal fit. We find $h_0^{95\%} = 2.04 \times 10^{-25}$ and 2.29×10^{-25} for $T_{\text{coh}} = 12$ hr and 9 hr, respectively, in this sample band. The $h_0^{95\%}$ values obtained in the sample subband improve upon the O3a results by a factor of approximately 1.4 (R. Abbott et al. 2021b). Given that the detectable strain amplitude scales with the effective amplitude spectral density and inversely with the square root of the observation time, the sensitivity achieved in O4a is consistent with expectations based on the O3a results.

Next, we use these values obtained in the sample subband (starting from 112.5 Hz for $f_{\star}$ and 225 Hz for $2f_{\star}$) to analytically calculate $h_0^{95\%}$ in the full frequency band (blue dots), using the scaling (L. Sun et al. 2019)

$$h_0^{95\%}(f) \propto \left(\frac{S_n(f)S_n(2f)}{S_n(f) + S_n(2f)} \right)^{1/2}, \quad (\text{D4})$$

where S_n is the effective power spectral density calculated from the harmonic mean of the two detectors over all the 30 minute SFTs collected from 2023 October 1 to 31 (GPS time 1380153618–1382745618). Finally, to verify the analytical scaling, the injection procedure in the first step is repeated in 12 randomly selected subbands that span the full range. The




Figure 12. Estimated sensitivity $h_0^{95\%}$ obtained from the DHV search as a function of $f_{\star}$ for $T_{\text{coh}} = 12$ hr (blue) and $T_{\text{coh}} = 9$ hr (orange), assuming a specific scenario with source properties $\theta = 45$ deg and $\cos \iota = 0$ (signals at both $f_{\star}$ and $2f_{\star}$ are linearly polarized). These estimates are obtained from randomized sky positions and hence apply to all sources using the same T_{coh} .

empirically obtained $h_0^{95\%}$ in these sample subbands agree with the analytic estimates to $<2.1\%$.

Appendix E Details on the Weave Search

E.1. Search Implementation

The Weave software provides a systematic way to cover the parameter space in a template search, ensuring an acceptable loss of signal-to-noise ratio for true signals that fall between template points (K. Wette et al. 2018). It combines recent

advances in template placement, using an optimal parameter-space metric (K. Wette & R. Prix 2013; K. Wette 2015) and optimal template lattices (K. Wette 2014).

Due to the high computational cost of template-based Weave searches, we limit our search to Cas A, Vela Jr., and G347.3–0.5 for GW signal frequency f ranges from 20 to 475 Hz. Detector noise deteriorates badly below 20 Hz because of ground motion and is also badly degraded at much higher frequencies in bands contaminated by forests of “violin mode” frequencies, near 500 Hz and near integer multiples thereof. These frequencies correspond to vibration resonances of the 16 suspension fibers supporting the four primary mirrors in each interferometer. Similar but much less severe resonances in suspension fibers supporting the beam splitter mirrors contaminate narrow bands between 300 and 400 Hz. In this search, we focus on frequencies below the fundamental violin mode harmonics for two reasons: (1) young, but not newborn, isolated NSs are expected to rotate at lower frequencies; and (2) higher-frequency searches are computationally more demanding because of the increase in parameter-space volume. We choose a ceiling of 475 Hz because the region between 475 Hz and the lowest violin mode harmonic (just below 500 Hz) is contaminated by nonlinear interactions between the violin modes and other, lower-frequency lines, leading to frequency “shoulders” and sidebands.

Similar to previous searches (J. Abadie et al. 2010; J. Aasi et al. 2015a; B. P. Abbott et al. 2019b; R. Abbott et al. 2021b, 2022a; J. Wang & K. Riles 2024; D. H. T. Cheung et al. 2026), we define the search range by assuming a power-law spin-down: $\dot{f} \propto -f^n$, with a braking index n ranging between 2 and 7 to account for multiple potential spin-down contributions. Additionally, we follow R. Abbott et al. (2022a) and J. Wang & K. Riles (2024) and adopt a slightly more conservative approach, allowing the upper limit on $\dot{f}$ to reach zero as follows:

$$-\frac{f}{t_{\text{age}}} \leq \dot{f} \leq 0, \quad 2\frac{\dot{f}^2}{f} \leq \ddot{f} \leq 7\frac{\dot{f}^2}{f}. \quad (\text{E1})$$

Setting the upper limit on $\dot{f}$ to zero allows for time-dependent braking indices and uncertainties in the source’s effective age. Table 13 lists the ages of the targets used to configure the search. It also includes the maximum absolute values of $\dot{f}$ and $\ddot{f}$ at the lowest and highest search frequencies.

The Weave program (K. Wette et al. 2018) creates a template grid in the parameter space for each time segment that is appropriate to computing the $\mathcal{F}$ -statistic for a coherence time T_{coh} equal to the total observation period T_{obs} divided by N_{seg} . The spacing of the grid points in $(f, \dot{f}, \ddot{f})$ is set according

to a metric (K. Wette & R. Prix 2013; K. Wette 2015) that ensures a worst-case maximum mismatch (m_{coh}) defined by the fractional loss in $\mathcal{F}$ -statistic value due to a true signal not coinciding with a search template.

Separately, a much finer grid is defined for the entire observation period with respect to the midpoint of the observation period, with its own mismatch parameter $m_{\text{semi-coh}}$, analogous to m_{coh} . This parameter, similar to m_{coh} , is the average of the coherent mismatch values across all segments (K. Wette 2015). Its value is chosen based on a balance between sensitivity and computational cost. At initialization, the Weave pipeline creates a mapping between each point in the semi-coherent template grid and the nearest point in each of the coarser segment grids, accounting for frequency evolution. The semi-coherent detection statistic $2\tilde{\mathcal{F}}$ is calculated as the average of the $2\mathcal{F}$ values across the N_{seg} segments for each semi-coherent template based on this mapping:

$$2\tilde{\mathcal{F}} = \frac{1}{N_{\text{seg}}} \sum_{i=1}^{N_{\text{seg}}} 2\mathcal{F}_i. \quad (\text{E2})$$

For the searches presented here, we use the same mismatch parameters $m_{\text{coh}} = 0.1$ and $m_{\text{semi-coh}} = 0.2$ as used in R. Abbott et al. (2022a).

We divide the search jobs into 0.1 Hz bands of f , with further divisions in $\dot{f}$, as needed, to keep each job’s computational duration less than 24 hr, for practical reasons. Table 14 shows the $\dot{f}$ divisions width for different frequency bands for the searches. The discreteness of the $\dot{f}$ and $\ddot{f}$ steps leads, at low frequencies, to a coverage of the parameter space well beyond that described by Equation (E1), a region that includes, for example, the parameters of the candidate reported in J. Ming et al. (2025).

Each individual search job returns the $(f, \dot{f}, \ddot{f})$ values of the 1000 templates (“top list”) with the largest (“loudest”) $2\tilde{\mathcal{F}}$ in each 0.1 Hz bands with $\dot{f}$ divisions. We sum up the number of templates for each 1 Hz band and calculate the nominal threshold

$$2\tilde{\mathcal{F}}_{\text{thresh}} = \frac{1}{N_{\text{seg}}} \text{CDF}_{\chi^2}^{-1} \left(1 - \frac{1}{N_{\text{temp}}}, 4N_{\text{seg}} \right), \quad (\text{E3})$$

where $\text{CDF}_{\chi^2}^{-1}$ is the inverse cumulative distribution function of the χ^2 distribution with $4N_{\text{seg}}$ degrees of freedom, evaluated at $1 - 1/N_{\text{temp}}$ percentile, and N_{temp} is the number of templates in a 1 Hz band. This threshold is set based on a signal-free χ^2 distribution with four degrees of freedom per segment, such that the expected number of candidates is one per 1 Hz band in

Table 13
Age Used to Define the Weave Search Range, Search Range for $\dot{f}$, $\ddot{f}$ Values at 20 and 475 Hz, and T_{coh} of Initial Searches

	G111.7–2.1/Cas A	G266.2–1.2/Vela Jr.	G347.3–0.5
t_{age} (yr)	300	690	1600
$\dot{f}$ range (Hz s ^{−1}) @20 Hz	$[-2.1 \times 10^{-9}, 0]$	$[-9.2 \times 10^{-9}, 0]$	$[-4.0 \times 10^{-10}, 0]$
$\ddot{f}$ range (Hz s ^{−2}) @20 Hz	$[0, 1.6 \times 10^{-18}]$	$[0, 3.0 \times 10^{-19}]$	$[0, 5.6 \times 10^{-20}]$
$\dot{f}$ range (Hz s ^{−1}) @475 Hz	$[-5.0 \times 10^{-8}, 0]$	$[-2.2 \times 10^{-8}, 0]$	$[-9.4 \times 10^{-9}, 0]$
$\ddot{f}$ range (Hz s ^{−2}) @475 Hz	$[0, 3.7 \times 10^{-17}]$	$[0, 7.0 \times 10^{-18}]$	$[0, 1.3 \times 10^{-18}]$
T_{coh} (day)	5	7.5	7.5

Table 14Width of $\dot{f}$ Subranges for Weave’s Initial Search Jobs (0.1 Hz Subbands) to Keep Job Durations under 24 Computational Hours

Source	f (Hz)	$\dot{f}$ Width (Hz s ⁻¹)
G111.7–2.1/Cas A	20–200	1.5×10^{-9}
G111.7–2.1/Cas A	200–475	1×10^{-9}
G266.2–1.2/Vela Jr.	20–475	8×10^{-10}
G347.3–0.5	20–475	8×10^{-10}

Note. Each subband is limited to a top list of 1000 candidates.**Table 15**Parameters Defining the Analytic Threshold Function $2\tilde{\mathcal{F}}_{\text{thresh}}(f) = 2\tilde{\mathcal{F}}_0 f^a$ Applied to the $2\tilde{\mathcal{F}}$ Detection Statistic

	f (Hz)	$2\tilde{\mathcal{F}}_0$	a
G111.7–2.1/Cas A	20–475	7.11	0.0186
G266.2–1.2/Vela Jr.	20–475	7.85	0.0218
G347.3–0.5	20–475	7.71	0.0198

Gaussian noise. We then fit this threshold to the function $2\tilde{\mathcal{F}}_{\text{thresh}}(f) = 2\tilde{\mathcal{F}}_0 f^a$. The fitted parameters are listed in Table 15. Templates with $2\tilde{\mathcal{F}}$ values exceeding this threshold are considered candidates. However, non-Gaussian artifacts lead to much higher candidate counts in particular bands contaminated by instrumental line sources. In some cases, strong instrumental lines can cause all templates returned by a single search job to have $\mathcal{F}$ -statistic values exceeding the nominal threshold. This occurs for a subset of the 0.1 Hz band and the range of $\dot{f}$ searched. We refer to those cases as “saturated” as potentially interesting templates can be suppressed by the top-list cap.

E.2. Postprocessing and Follow-up

For nonsaturated subranges of individual 0.1 Hz bands, candidates exceeding the threshold $2\tilde{\mathcal{F}}_{\text{thresh}}(f)$ are followed up in a sequential procedure where at each step, the coherence time T_{coh} is increased (and hence the number of segments N_{seg} is decreased). Because the noncentrality parameter for the $2\tilde{\mathcal{F}}$ detection statistic scales approximately linearly with T_{coh} , one expects a similar percentage increase in the $2\tilde{\mathcal{F}}$. To determine the required ratio increase in $2\tilde{\mathcal{F}}$, We perform 50 software injections (signal simulations) with randomly chosen frequency parameters within each 1 Hz band and with $h_0 = h_{\text{sens}}^{95\%}$ that yields 95% detection efficiency (see Section E.3). We then follow up candidates with $2\tilde{\mathcal{F}}$ above the $2\tilde{\mathcal{F}}_{\text{thresh}}(f)$ threshold and include higher-order spin-down terms to better capture the signal, given the coherence time and spin-down order configuration for each follow-up stage, as shown in Table 16. The requirement for the increase in $2\tilde{\mathcal{F}}$ to recover the injected signals above the $2\tilde{\mathcal{F}}_{\text{thresh}}(f)$ threshold for all follow-up stages is listed in Table 17.

In addition, following up only the top 10 loudest survivors at later follow-up stages is sufficient to recover the injected signals with high efficiency. Therefore, although we evaluate

Table 16

Coherence Time and Spin-down Order Configuration for Different Weave Follow-up Stages

	G111.7–2.1/Cas A		G266.2–1.2/Vela Jr., G347.3–0.5	
1st follow-up T_{coh} (spin-down order) change	5 days "	10 days (2 " 2)	7.5 days "	15 days(2 " 2)
2nd follow-up	10 "	20(2 " 2)	15 "	30(2 " 2)
3rd follow-up	20 "	40(2 " 3)	30 "	60(2 " 3)
4th follow-up	-	-	60 "	120(3 " 4)
5th follow-up	-	-	120 "	240(4 " 4)

Table 17 $2\tilde{\mathcal{F}}$ Increase Requirement in Each Stage of Weave Candidate Follow-up

	G111.7–2.1/Cas A	G266.2–1.2/Vela Jr.	G347.3–0.5
1	32%	36%	35%
2	41%	49%	49%
3	62%	62%	60%
4	...	68%	65%
5	...	74%	70%

all candidates above the $2\tilde{\mathcal{F}}_{\text{thresh}}(f)$ threshold from the initial search stage, in the successive stages, only the top 10 loudest survivors per initial candidate are evaluated for further follow-up. We have verified that these threshold criteria preserve greater than 95% recovery efficiency for the injections, while allowing conservatively for an uncertainty in source R.A. and decl. of 10^{-5} radians ($\sim 2^\circ$). The loudest candidate in each saturated subrange is followed up using the same procedure to ensure that if the saturation were due to an unusually strong signal, it would not be suppressed by the top-list cap.

We further validated the follow-up procedure with four “hardware injections” that were injected in the LIGO detector during the run via modulation of interferometer mirror positions via auxiliary lasers (C. Biwer et al. 2017). We chose Injections 3, 6, 8, and 12, ranging from ~ 37 to ~ 189 Hz, which use the same simulation parameters as used in the O3 run (R. Abbott et al. 2021d), except for somewhat reduced amplitude. All signals were successfully recovered.

For Cas A/Vela Jr./G347.3–0.5, there were $\sim 1.3 \times 10^5 / 1.2 \times 10^5 / 6.9 \times 10^4$ candidates above the initial search threshold in bands that were not saturated (not excluded from consideration by severe instrumental artifacts). The parameters of the surviving candidates after the second follow-up stage and the complete lists of the saturated subbands can be found in D. Cheung (2025).

For Cas A, repeatedly doubling the coherence time to $T_{\text{coh}} = 40$ days eliminates all candidates. For G347.3–0.5, we eliminate all candidates in the final follow-up stage when $T_{\text{coh}} = 240$ days. On the other hand, two clusters of candidates for Vela Jr. survive the final follow-up stage ($T_{\text{coh}} = 240$ days) with source frequencies near 247 Hz. In this case, there is a prolonged period during the run when an instrumental line overlaps the nominal signal’s frequency trajectory versus time, a period during which there is a disproportionate contribution to the detection statistic. Hence, we do not consider either candidate cluster to be a credible astrophysical signal.

We conclude that there is no significant evidence in the Weave pipeline analysis for a CW signal from the CCOs in the SNRs Cas A, Vela Jr., and G347.3–0.5.

E.3. Upper Limits

In Figure 13, we report the 95% confidence-level frequentist upper limits on strain amplitude $h_0^{95\%}$ for all the targets searched by Weave. To obtain these upper limits, we conduct software injections of signals with varying amplitudes, drawing frequencies uniformly within each 1 Hz band (excluding 0.1 Hz subbands containing at least one $\dot{f}$ interval with saturated candidate top lists). The parameters $\cos\iota$, ψ , $\dot{f}$, and $\ddot{f}$ are also drawn uniformly. An injection is deemed detectable if its $2\tilde{\mathcal{F}}$ value, computed using Weave with the initial search setup, meets or exceeds $2\tilde{\mathcal{F}}_{\text{thresh}}(f)$ (assuming all candidates above this threshold corresponding to a true signal would be followed up and confirmed). This procedure is repeated for different signal amplitudes h_0 , and the detection fraction is fitted using a sigmoidal function

$$p(h_0) = \frac{1}{1 + \exp\left(\frac{a-h_0}{b}\right)}, \quad (\text{E4})$$

where $p(h_0)$ represents the fraction of detected injections for a given h_0 , and a and b are the parameters of the sigmoidal function. The number of injections is chosen to ensure that the statistical uncertainty in the estimated $h_0^{95\%}$ is below 5%.




Figure 13. GW strain amplitude upper limits (95% confidence level) in each 1 Hz band for the Cas A (blue), Vela Jr. (orange), and G347.3–0.5 (green) searches using Weave. All the upper limits presented here are below the age-based strain amplitude limit ($>10^{-24}$) for the three targets.

ORCID iDs

- A. G. Abac <https://orcid.org/0000-0003-4786-2698>
K. Ackley <https://orcid.org/0000-0002-8648-0767>
C. Adamcewicz <https://orcid.org/0000-0001-5525-6255>
S. Adhicary <https://orcid.org/0009-0004-2101-5428>
N. Adhikari <https://orcid.org/0000-0002-4559-8427>
R. X. Adhikari <https://orcid.org/0000-0002-5731-5076>
S. Afroz <https://orcid.org/0009-0004-4459-2981>
D. Agarwal <https://orcid.org/0000-0002-8735-5554>
M. Agathos <https://orcid.org/0000-0002-9072-1121>
O. D. Aguiar <https://orcid.org/0000-0002-2139-4390>
L. Aiello <https://orcid.org/0000-0003-2771-8816>
A. Ain <https://orcid.org/0000-0003-4534-4619>
P. Ajith <https://orcid.org/0000-0001-7519-2439>
T. Akutsu <https://orcid.org/0000-0003-0733-7530>
S. Albanesi <https://orcid.org/0000-0001-7345-4415>
A. Allocca <https://orcid.org/0000-0002-5288-1351>
P. A. Altin <https://orcid.org/0000-0001-8193-5825>
S. Alvarez-Lopez <https://orcid.org/0009-0003-8040-4936>
A. Amato <https://orcid.org/0000-0001-9557-651X>
F. Amicucci <https://orcid.org/0009-0005-2139-4197>
S. B. Anderson <https://orcid.org/0000-0003-2219-9383>
W. G. Anderson <https://orcid.org/0000-0003-0482-5942>
M. Andia <https://orcid.org/0000-0003-3675-9126>
M. Andrés-Carcasona <https://orcid.org/0000-0002-8738-1672>
T. Andrić <https://orcid.org/0000-0002-9277-9773>
S. Ansolli <https://orcid.org/0000-0002-5613-7693>
J. M. Antelis <https://orcid.org/0000-0003-3377-0813>
S. Antier <https://orcid.org/0000-0002-7686-3334>
S. K. Apple <https://orcid.org/0009-0007-4490-5804>
K. Arai <https://orcid.org/0000-0001-8916-8915>
A. Araya <https://orcid.org/0000-0002-6884-2875>
M. C. Araya <https://orcid.org/0000-0002-6018-6447>
M. Arca Sedda <https://orcid.org/0000-0002-3987-0519>
J. S. Areeda <https://orcid.org/0000-0003-0266-7936>
F. Armato <https://orcid.org/0000-0002-8856-8877>
S. Armstrong <https://orcid.org/0009-0009-4285-2360>
N. Arnaud <https://orcid.org/0000-0001-6589-8673>
M. Arogeti <https://orcid.org/0000-0001-5124-3350>
S. M. Aronson <https://orcid.org/0000-0001-7080-8177>
G. Ashton <https://orcid.org/0000-0001-7288-2231>
Y. Aso <https://orcid.org/0000-0002-1902-6695>
P. Astone <https://orcid.org/0000-0003-4981-4120>
F. Attadio <https://orcid.org/0009-0008-8916-1658>
F. Aubin <https://orcid.org/0000-0003-1613-3142>
K. AultONeal <https://orcid.org/0000-0002-6645-4473>
G. Avallone <https://orcid.org/0000-0001-5482-0299>
E. A. Avila <https://orcid.org/0009-0008-9329-4525>
S. Babak <https://orcid.org/0000-0001-7469-4250>
S. Bae <https://orcid.org/0000-0003-2429-3357>
S. Bagnasco <https://orcid.org/0000-0001-6062-6505>

- L. Baiotti  <https://orcid.org/0000-0003-0458-4288>
R. Bajpai  <https://orcid.org/0000-0003-0495-5720>
T. Baker  <https://orcid.org/0000-0001-5470-7616>
G. Baldi  <https://orcid.org/0000-0001-8963-3362>
N. Baldicchi  <https://orcid.org/0009-0009-8888-291X>
S. Banagiri  <https://orcid.org/0000-0001-7852-7484>
B. Banerjee  <https://orcid.org/0000-0002-8008-2485>
D. Bankar  <https://orcid.org/0000-0002-6068-2993>
P. Baral  <https://orcid.org/0000-0001-6308-211X>
M. Baratti  <https://orcid.org/0009-0003-5744-8025>
P. Barneo  <https://orcid.org/0000-0002-8883-7280>
F. Barone  <https://orcid.org/0000-0002-8069-8490>
B. Barr  <https://orcid.org/0000-0002-5232-2736>
L. Barsotti  <https://orcid.org/0000-0001-9819-2562>
M. Barsuglia  <https://orcid.org/0000-0002-1180-4050>
D. Barta  <https://orcid.org/0000-0001-6841-550X>
M. A. Barton  <https://orcid.org/0000-0002-9948-306X>
A. Basalaev  <https://orcid.org/0000-0001-5623-2853>
R. Bassiri  <https://orcid.org/0000-0001-8171-6833>
A. Basti  <https://orcid.org/0000-0003-2895-9638>
M. Bawaj  <https://orcid.org/0000-0003-3611-3042>
J. C. Bayley  <https://orcid.org/0000-0003-2306-4106>
A. C. Baylor  <https://orcid.org/0000-0003-0918-0864>
F. Beirnaert  <https://orcid.org/0000-0002-4003-7233>
M. Beijer  <https://orcid.org/0000-0002-4991-8213>
D. Belardinelli  <https://orcid.org/0000-0001-9332-5733>
A. S. Bell  <https://orcid.org/0000-0003-1523-0821>
L. Bellizzi  <https://orcid.org/0000-0002-2071-0400>
W. Benoît  <https://orcid.org/0000-0003-4750-9413>
I. Bentara  <https://orcid.org/0009-0000-5074-839X>
J. D. Bentley  <https://orcid.org/0000-0002-4736-7403>
S. Bera  <https://orcid.org/0000-0003-0907-6098>
F. Bergamin  <https://orcid.org/0000-0002-1113-9644>
B. K. Berger  <https://orcid.org/0000-0002-4845-8737>
S. Bernuzzi  <https://orcid.org/0000-0002-2334-0935>
M. Beroiz  <https://orcid.org/0000-0001-6486-9897>
D. Bersanetti  <https://orcid.org/0000-0002-7377-415X>
J. Betzwieser  <https://orcid.org/0000-0003-1533-9229>
D. Beveridge  <https://orcid.org/0000-0002-1481-1993>
G. Bevilacqua  <https://orcid.org/0000-0002-7298-6185>
N. Bevin  <https://orcid.org/0000-0002-4312-4287>
D. Bhattacharjee  <https://orcid.org/0000-0001-6623-9506>
S. Bhaumik  <https://orcid.org/0000-0001-8492-2202>
V. Biancalana  <https://orcid.org/0000-0002-1642-5391>
G. Billingsley  <https://orcid.org/0000-0002-4141-2744>
A. Binetti  <https://orcid.org/0000-0001-6449-5493>
S. Bini  <https://orcid.org/0000-0002-0267-3562>
O. Birnholtz  <https://orcid.org/0000-0002-7562-9263>
S. Biscoveanu  <https://orcid.org/0000-0001-7616-7366>
M. Bitossi  <https://orcid.org/0000-0002-9862-4668>
M.-A. Bizouard  <https://orcid.org/0000-0002-4618-1674>
J. K. Blackburn  <https://orcid.org/0000-0002-3838-2986>
N. Bode  <https://orcid.org/0000-0002-7101-9396>
G. Boileau  <https://orcid.org/0000-0002-3576-6968>
M. Boldrini  <https://orcid.org/0000-0001-9861-821X>
G. N. Bolingbroke  <https://orcid.org/0000-0002-7350-5291>
L. D. Bonavena  <https://orcid.org/0000-0002-2630-6724>
R. Bondarescu  <https://orcid.org/0000-0003-0330-2736>
F. Bondu  <https://orcid.org/0000-0001-6487-5197>
E. Bonilla  <https://orcid.org/0000-0002-6284-9769>
M. S. Bonilla  <https://orcid.org/0000-0003-4502-528X>
R. Bonnand  <https://orcid.org/0000-0001-5013-5913>
V. Boschi  <https://orcid.org/0000-0001-8665-2293>
Y. Bothra  <https://orcid.org/0000-0002-9380-6390>
P. R. Brady  <https://orcid.org/0000-0002-4611-9387>
M. Branchesi  <https://orcid.org/0000-0003-1643-0526>
T. Briant  <https://orcid.org/0000-0002-6013-1729>
E. Brockmueller  <https://orcid.org/0000-0002-1489-942X>
A. F. Brooks  <https://orcid.org/0000-0003-4295-792X>
M. L. Brozzetti  <https://orcid.org/0000-0002-5260-4979>
R. Bruntz  <https://orcid.org/0000-0002-0840-8567>
F. Buccì  <https://orcid.org/0000-0003-1726-3838>
O. Bulashenko  <https://orcid.org/0000-0003-1720-4061>
A. Buonanno  <https://orcid.org/0000-0002-5433-1409>
R. Buscicchio  <https://orcid.org/0000-0002-7387-6754>
C. Buy  <https://orcid.org/0000-0003-2872-8186>
G. S. Cabourn Davies  <https://orcid.org/0000-0002-4289-3439>
R. Cabrita  <https://orcid.org/0000-0003-0133-1306>
V. Cáceres-Barbosa  <https://orcid.org/0000-0001-9834-4781>
L. Cadonati  <https://orcid.org/0000-0002-9846-166X>
G. Cagnoli  <https://orcid.org/0000-0002-7086-6550>
C. Cahillane  <https://orcid.org/0000-0002-3888-314X>
S. R. Callos  <https://orcid.org/0000-0003-0639-9342>
G. Caneva Santoro  <https://orcid.org/0000-0002-2935-1600>
K. C. Cannon  <https://orcid.org/0000-0003-4068-6572>
E. Capocasa  <https://orcid.org/0000-0003-3762-6958>
E. Capote  <https://orcid.org/0009-0007-0246-713X>
G. Capurri  <https://orcid.org/0000-0003-0889-1015>
J. B. Carlin  <https://orcid.org/0000-0001-5694-0809>
M. Carpinelli  <https://orcid.org/0000-0002-8205-930X>
J. J. Carter  <https://orcid.org/0000-0001-8845-0900>
G. Carullo  <https://orcid.org/0000-0001-9090-1862>
J. Casanueva Diaz  <https://orcid.org/0000-0002-2948-5238>
C. Casentini  <https://orcid.org/0000-0001-8100-0579>
M. Cavaglià  <https://orcid.org/0000-0002-3835-6729>
R. Cavalieri  <https://orcid.org/0000-0001-6064-0569>
G. Cella  <https://orcid.org/0000-0002-0752-0338>
P. Cerdá-Durán  <https://orcid.org/0000-0003-4293-340X>
E. Cesarini  <https://orcid.org/0000-0001-9127-3167>
A. Chakraborty  <https://orcid.org/0009-0004-4937-4633>
P. Chakraborty  <https://orcid.org/0000-0002-0994-7394>
S. Chalathadka Subrahmanya  <https://orcid.org/0000-0002-9207-4669>
J. C. L. Chan  <https://orcid.org/0000-0002-3377-4737>
S. Chao  <https://orcid.org/0000-0003-3853-3593>
P. Charlton  <https://orcid.org/0000-0002-4263-2706>
E. Chassande-Mottin  <https://orcid.org/0000-0003-3768-9908>
C. Chatterjee  <https://orcid.org/0000-0001-8700-3455>
Debarati Chatterjee  <https://orcid.org/0000-0002-0995-2329>
Deep Chatterjee  <https://orcid.org/0000-0003-0038-5468>
S. Chaty  <https://orcid.org/0000-0002-5769-8601>
A. Chen  <https://orcid.org/0000-0001-9174-7780>
D. Chen  <https://orcid.org/0000-0003-1433-0716>
H. Y. Chen  <https://orcid.org/0000-0001-5403-3762>
Yitian Chen  <https://orcid.org/0000-0002-8664-9702>
P. Chessa  <https://orcid.org/0000-0001-9092-3965>
H. T. Cheung  <https://orcid.org/0000-0003-3905-0665>
F. Chiadini  <https://orcid.org/0000-0002-9339-8622>
A. Chincarini  <https://orcid.org/0000-0003-4094-9942>
M. L. Chiofalo  <https://orcid.org/0000-0002-6992-5963>
A. Chiummo  <https://orcid.org/0000-0003-2165-2967>

- S. Choudhary  <https://orcid.org/0000-0003-0949-7298>
 N. Christensen  <https://orcid.org/0000-0002-6870-4202>
 S. S. Y. Chua  <https://orcid.org/0000-0001-8026-7597>
 G. Ciani  <https://orcid.org/0000-0003-4258-9338>
 P. Cieciela  <https://orcid.org/0000-0002-5871-4730>
 M. Cieřlar  <https://orcid.org/0000-0001-8912-5587>
 M. Cifaldi  <https://orcid.org/0009-0007-1566-7093>
 J. A. Clark  <https://orcid.org/0000-0003-3243-1393>
 T. A. Clarke  <https://orcid.org/0000-0002-6714-5429>
 E. Codazzo  <https://orcid.org/0000-0001-7170-8733>
 P.-F. Cohadon  <https://orcid.org/0000-0003-3452-9415>
 S. Colace  <https://orcid.org/0009-0007-9429-1847>
 M. Colleoni  <https://orcid.org/0000-0002-7214-9088>
 S. Colloms  <https://orcid.org/0009-0009-9828-3646>
 A. Colombo  <https://orcid.org/0000-0002-7439-4773>
 L. Conti  <https://orcid.org/0000-0003-2731-2656>
 T. R. Corbitt  <https://orcid.org/0000-0002-5520-8541>
 I. Cordero-Carrión  <https://orcid.org/0000-0002-1985-1361>
 S. Corezzi  <https://orcid.org/0000-0002-3437-5949>
 N. J. Cornish  <https://orcid.org/0000-0002-7435-0869>
 A. Corsi  <https://orcid.org/0000-0001-8104-3536>
 M. W. Coughlin  <https://orcid.org/0000-0002-8262-2924>
 P. Couvares  <https://orcid.org/0000-0002-2823-3127>
 R. Coyne  <https://orcid.org/0000-0002-5243-5917>
 J. D. E. Creighton  <https://orcid.org/0000-0003-3600-2406>
 P. Cremonese  <https://orcid.org/0000-0001-6472-8509>
 J. R. Cudell  <https://orcid.org/0000-0002-2003-4238>
 T. J. Cullen  <https://orcid.org/0000-0001-8075-4088>
 A. Cumming  <https://orcid.org/0000-0003-4096-7542>
 E. Cuoco  <https://orcid.org/0000-0002-6528-3449>
 M. Cusinato  <https://orcid.org/0000-0003-4075-4539>
 L. V. Da Conceição  <https://orcid.org/0000-0002-5042-443X>
 T. Dal Canton  <https://orcid.org/0000-0001-5078-9044>
 S. Dal Pra  <https://orcid.org/0000-0002-1057-2307>
 G. Dálya  <https://orcid.org/0000-0003-3258-5763>
 B. D'Angelo  <https://orcid.org/0000-0001-9143-8427>
 S. Danilishin  <https://orcid.org/0000-0001-7758-7493>
 S. D'Antonio  <https://orcid.org/0000-0003-0898-6030>
 L. P. Dartez  <https://orcid.org/0000-0002-2216-0465>
 V. Dattilo  <https://orcid.org/0000-0002-8816-8566>
 D. Davis  <https://orcid.org/0000-0001-5620-6751>
 M. C. Davis  <https://orcid.org/0000-0001-7663-0808>
 P. Davis  <https://orcid.org/0009-0004-5008-5660>
 E. J. Daw  <https://orcid.org/0000-0002-3780-5430>
 M. Dax  <https://orcid.org/0000-0001-8798-0627>
 J. De Bolle  <https://orcid.org/0000-0002-5179-1725>
 J. Degallaix  <https://orcid.org/0000-0002-1019-6911>
 M. De Laurentis  <https://orcid.org/0000-0002-3815-4078>
 F. De Lillo  <https://orcid.org/0000-0003-4977-0789>
 S. Della Torre  <https://orcid.org/0000-0002-7669-0859>
 W. Del Pozzo  <https://orcid.org/0000-0003-3978-2030>
 F. De Marco  <https://orcid.org/0000-0002-5411-9424>
 F. De Matteis  <https://orcid.org/0000-0001-7860-9754>
 T. Dent  <https://orcid.org/0000-0003-1354-7809>
 A. Depasse  <https://orcid.org/0000-0003-1014-8394>
 R. De Pietri  <https://orcid.org/0000-0003-1556-8304>
 R. De Rosa  <https://orcid.org/0000-0002-4004-947X>
 C. De Rossi  <https://orcid.org/0000-0002-5825-472X>
 M. Desai  <https://orcid.org/0009-0003-4448-3681>
 R. DeSalvo  <https://orcid.org/0000-0002-4818-0296>
 A. Dhani  <https://orcid.org/0000-0001-9930-9101>
 M. C. Díaz  <https://orcid.org/0000-0002-7555-8856>
 M. Di Cesare  <https://orcid.org/0009-0003-0411-6043>
 T. Dietrich  <https://orcid.org/0000-0003-2374-307X>
 C. Di Fronzo  <https://orcid.org/0000-0002-2693-6769>
 M. Di Giovanni  <https://orcid.org/0000-0003-4049-8336>
 T. Di Girolamo  <https://orcid.org/0000-0003-2339-4471>
 J. Ding  <https://orcid.org/0000-0003-1693-3828>
 S. Di Pace  <https://orcid.org/0000-0001-6759-5676>
 I. Di Palma  <https://orcid.org/0000-0003-1544-8943>
 F. Di Renzo  <https://orcid.org/0000-0002-5447-3810>
 Divyajyoti  <https://orcid.org/0000-0002-2787-1012>
 A. Dmitriev  <https://orcid.org/0000-0002-0314-956X>
 Z. Doctort  <https://orcid.org/0000-0002-2077-4914>
 N. Doerksen  <https://orcid.org/0009-0002-3776-5026>
 L. D'Onofrio  <https://orcid.org/0000-0001-9546-5959>
 K. L. Dooley  <https://orcid.org/0000-0002-1636-0233>
 S. Doravari  <https://orcid.org/0000-0001-8750-8330>
 M. Drago  <https://orcid.org/0000-0002-3738-2431>
 J. C. Driggers  <https://orcid.org/0000-0002-6134-7628>
 L. Dunn  <https://orcid.org/0000-0002-1769-6097>
 P.-A. Duverne  <https://orcid.org/0000-0002-3906-0997>
 D. D'Urso  <https://orcid.org/0000-0002-8215-4542>
 P. Dutta Roy  <https://orcid.org/0000-0001-8874-4888>
 H. Duval  <https://orcid.org/0000-0002-2475-1728>
 M. Ebersold  <https://orcid.org/0000-0003-4631-1771>
 T. Eckhardt  <https://orcid.org/0000-0002-1224-4681>
 G. Eddolls  <https://orcid.org/0000-0002-5895-4523>
 A. Effler  <https://orcid.org/0000-0001-8242-3944>
 J. Eichholz  <https://orcid.org/0000-0002-2643-163X>
 M. Emma  <https://orcid.org/0000-0001-7943-0262>
 R. Enficiaud  <https://orcid.org/0000-0003-3908-1912>
 L. Errico  <https://orcid.org/0000-0003-2112-0653>
 M. Esposito  <https://orcid.org/0009-0009-8482-9417>
 R. C. Essick  <https://orcid.org/0000-0001-8196-9267>
 H. Estellés  <https://orcid.org/0000-0001-6143-5532>
 M. Evans  <https://orcid.org/0000-0001-8459-4499>
 J. M. Ezquiaga  <https://orcid.org/0000-0002-7213-3211>
 F. Fabrizi  <https://orcid.org/0000-0002-3809-065X>
 V. Fafone  <https://orcid.org/0000-0003-1314-1622>
 S. Fairhurst  <https://orcid.org/0000-0001-8480-1961>
 A. M. Farah  <https://orcid.org/0000-0002-6121-0285>
 B. Farr  <https://orcid.org/0000-0002-2916-9200>
 W. M. Farr  <https://orcid.org/0000-0003-1540-8562>
 G. Favaro  <https://orcid.org/0000-0002-0351-6833>
 M. Favata  <https://orcid.org/0000-0001-8270-9512>
 M. Fays  <https://orcid.org/0000-0002-4390-9746>
 M. Fazio  <https://orcid.org/0000-0002-9057-9663>
 R. Felicetti  <https://orcid.org/0009-0005-6263-5604>
 E. Fenyvesi  <https://orcid.org/0000-0003-2777-3719>
 T. Fernandes  <https://orcid.org/0009-0006-6820-2065>
 S. Ferraiuolo  <https://orcid.org/0009-0005-5582-2989>
 F. Fidecaro  <https://orcid.org/0000-0002-6189-3311>
 P. Figura  <https://orcid.org/0000-0002-8925-0393>
 A. Fiori  <https://orcid.org/0000-0003-3174-0688>
 I. Fiori  <https://orcid.org/0000-0002-0210-516X>
 M. Fishbach  <https://orcid.org/0000-0002-1980-5293>
 R. Fittipaldi  <https://orcid.org/0000-0003-2096-7983>
 V. Fiumara  <https://orcid.org/0000-0003-3644-217X>
 S. M. Fleischer  <https://orcid.org/0000-0001-7884-9993>
 J. A. Font  <https://orcid.org/0000-0001-6650-2634>
 B. Fornal  <https://orcid.org/0000-0003-3271-2080>
 F. Frascioni  <https://orcid.org/0000-0003-4204-6587>

- Z. Frei  <https://orcid.org/0000-0002-0181-8491>
A. Freise  <https://orcid.org/0000-0001-6586-9901>
O. Freitas  <https://orcid.org/0000-0002-2898-1256>
R. Frey  <https://orcid.org/0000-0003-0341-2636>
G. G. Fronzé  <https://orcid.org/0000-0003-0966-4279>
M. Fuentes-Garcia  <https://orcid.org/0000-0003-3390-8712>
B. Gadre  <https://orcid.org/0000-0002-1534-9761>
J. R. Gair  <https://orcid.org/0000-0002-1671-3668>
S. Galaudage  <https://orcid.org/0000-0002-1819-0215>
A. Gamboa  <https://orcid.org/0000-0001-8391-5596>
D. Ganapathy  <https://orcid.org/0000-0003-3028-4174>
A. Ganguly  <https://orcid.org/0000-0001-7394-0755>
B. Garaventa  <https://orcid.org/0000-0003-2490-404X>
J. García-Bellido  <https://orcid.org/0000-0002-9370-8360>
C. García-Quirós  <https://orcid.org/0000-0002-8059-2477>
J. W. Gardner  <https://orcid.org/0000-0002-8592-1452>
J. Gargiulo  <https://orcid.org/0000-0002-3507-6924>
X. Garrido  <https://orcid.org/0000-0002-7088-5831>
A. Garron  <https://orcid.org/0000-0002-1601-797X>
F. Garufi  <https://orcid.org/0000-0003-1391-6168>
C. Gasbarra  <https://orcid.org/0000-0001-8335-9614>
F. Gautier  <https://orcid.org/0000-0001-8006-9590>
V. Gayathri  <https://orcid.org/0000-0002-7167-9888>
G. Gemme  <https://orcid.org/0000-0002-1127-7406>
A. Gennai  <https://orcid.org/0000-0003-0149-2089>
V. Gennari  <https://orcid.org/0000-0002-0190-9262>
R. George  <https://orcid.org/0000-0002-7797-7683>
O. Gerberding  <https://orcid.org/0000-0001-7740-2698>
L. Gergely  <https://orcid.org/0000-0003-3146-6201>
Archisman Ghosh  <https://orcid.org/0000-0003-0423-3533>
Shaon Ghosh  <https://orcid.org/0000-0001-9901-6253>
Suprovo Ghosh  <https://orcid.org/0000-0002-1656-9870>
Tathagata Ghosh  <https://orcid.org/0000-0001-9848-9905>
J. A. Giaime  <https://orcid.org/0000-0002-3531-817X>
C. Gier  <https://orcid.org/0000-0003-0897-7943>
S. Gkaitatzis  <https://orcid.org/0000-0001-9420-7499>
J. Glanzer  <https://orcid.org/0009-0000-0808-0795>
F. Glotin  <https://orcid.org/0000-0003-2637-1187>
P. Godwin  <https://orcid.org/0000-0002-7489-4751>
A. S. Goettel  <https://orcid.org/0000-0002-6215-4641>
E. Goetz  <https://orcid.org/0000-0003-2666-721X>
S. Gomez Lopez  <https://orcid.org/0000-0002-9557-4706>
B. Goncharov  <https://orcid.org/0000-0003-3189-5807>
G. González  <https://orcid.org/0000-0003-0199-3158>
P. Goodarzi  <https://orcid.org/0009-0008-1093-6706>
A. W. Goodwin-Jones  <https://orcid.org/0000-0002-0395-0680>
R. Gouaty  <https://orcid.org/0000-0001-5372-7084>
A. Grado  <https://orcid.org/0000-0002-0501-8256>
V. Graham  <https://orcid.org/0000-0003-3633-0135>
A. E. Granados  <https://orcid.org/0000-0003-2099-9096>
M. Granata  <https://orcid.org/0000-0003-3275-1186>
V. Granata  <https://orcid.org/0000-0003-2246-6963>
R. Gray  <https://orcid.org/0000-0002-5556-9873>
A. C. Green  <https://orcid.org/0000-0002-6287-8746>
S. R. Green  <https://orcid.org/0000-0002-6987-6313>
H. L. Griggs  <https://orcid.org/0000-0001-5018-7908>
C. Grimaud  <https://orcid.org/0000-0001-7736-7730>
H. Grote  <https://orcid.org/0000-0002-0797-3943>
S. Grunewald  <https://orcid.org/0000-0003-4641-2791>
D. Guerra  <https://orcid.org/0000-0003-0029-5390>
D. Guetta  <https://orcid.org/0000-0002-7349-1109>
G. M. Guidi  <https://orcid.org/0000-0002-3061-9870>
F. Gulminelli  <https://orcid.org/0000-0003-4354-2849>
H. Guo  <https://orcid.org/0000-0002-3777-3117>
W. Guo  <https://orcid.org/0000-0002-4320-4420>
Y. Guo  <https://orcid.org/0000-0002-6959-9870>
Anuradha Gupta  <https://orcid.org/0000-0002-5441-9013>
I. Gupta  <https://orcid.org/0000-0001-6932-8715>
V. Gupta  <https://orcid.org/0000-0002-7672-0480>
F. Guzman  <https://orcid.org/0000-0001-9136-929X>
M. Haberland  <https://orcid.org/0000-0001-9816-5660>
E. D. Hall  <https://orcid.org/0000-0001-9018-666X>
E. Z. Hamilton  <https://orcid.org/0000-0003-0098-9114>
G. Hammond  <https://orcid.org/0000-0002-1414-3622>
C. Hanna  <https://orcid.org/0000-0002-0965-7493>
O. A. Hannuksela  <https://orcid.org/0000-0002-3887-7137>
A. G. Hanselman  <https://orcid.org/0000-0002-8304-0109>
S. Harikumar  <https://orcid.org/0000-0002-2653-7282>
T. Harmark  <https://orcid.org/0000-0002-2795-7035>
J. Harms  <https://orcid.org/0000-0002-7332-9806>
G. M. Harry  <https://orcid.org/0000-0002-8905-7622>
I. W. Harry  <https://orcid.org/0000-0002-5304-9372>
C.-J. Haster  <https://orcid.org/0000-0001-8040-9807>
K. Haughian  <https://orcid.org/0000-0002-1223-7342>
J. Heinze  <https://orcid.org/0000-0001-8692-2724>
H. Heitmann  <https://orcid.org/0000-0003-0625-5461>
F. Hellman  <https://orcid.org/0000-0002-9135-6330>
A. F. Helmling-Cornell  <https://orcid.org/0000-0002-7709-8638>
G. Hemming  <https://orcid.org/0000-0001-5268-4465>
O. Henderson-Sapir  <https://orcid.org/0000-0002-1613-9985>
M. Hendry  <https://orcid.org/0000-0001-8322-5405>
M. H. Hennig  <https://orcid.org/0000-0003-1531-8460>
C. Henshaw  <https://orcid.org/0000-0002-4206-3128>
M. Heurs  <https://orcid.org/0000-0002-5577-2273>
A. L. Hewitt  <https://orcid.org/0000-0002-1255-3492>
Y. Himemoto  <https://orcid.org/0000-0002-6856-3809>
I. J. Hollows  <https://orcid.org/0000-0002-3404-6459>
D. E. Holz  <https://orcid.org/0000-0002-0175-5064>
J. Hough  <https://orcid.org/0000-0003-3242-3123>
S. Hourihane  <https://orcid.org/0000-0002-9152-0719>
E. J. Howell  <https://orcid.org/0000-0001-7891-2817>
C. G. Hoy  <https://orcid.org/0000-0002-8843-6719>
H.-F. Hsieh  <https://orcid.org/0000-0002-8947-723X>
W.-F. Hsu  <https://orcid.org/0000-0001-5234-3804>
Q. Hu  <https://orcid.org/0000-0002-3033-6491>
H. Y. Huang  <https://orcid.org/0000-0002-1665-2383>
Y. Huang  <https://orcid.org/0000-0002-2952-8429>
V. Hui  <https://orcid.org/0000-0002-0233-2346>
S. Husa  <https://orcid.org/0000-0002-0445-1971>
L. Iampieri  <https://orcid.org/0009-0004-1161-2990>
G. A. Iandolo  <https://orcid.org/0000-0003-1155-4327>
G. Iannone  <https://orcid.org/0000-0001-8347-7549>
G. Iorio  <https://orcid.org/0000-0003-0293-503X>
P. Iosif  <https://orcid.org/0000-0003-1621-7709>
J. Irwin  <https://orcid.org/0000-0002-2364-2191>
M. Isi  <https://orcid.org/0000-0001-8830-8672>
K. S. Isleif  <https://orcid.org/0000-0001-7032-9440>
Y. Itoh  <https://orcid.org/0000-0003-2694-8935>
B. R. Iyer  <https://orcid.org/0000-0002-4141-5179>
P.-E. Jacquet  <https://orcid.org/0000-0001-9552-0057>
S. P. Jadhav  <https://orcid.org/0000-0003-0554-0084>

- A. L. James  <https://orcid.org/0000-0001-9165-0807>
K. Jani  <https://orcid.org/0000-0003-1007-8912>
J. Janquart  <https://orcid.org/0000-0003-2888-7152>
S. Jaraba  <https://orcid.org/0000-0002-4759-143X>
P. Jaranowski  <https://orcid.org/0000-0001-8085-3414>
R. Jaume  <https://orcid.org/0000-0001-8691-3166>
J. Jiang  <https://orcid.org/0000-0002-0154-3854>
H.-B. Jin  <https://orcid.org/0000-0002-6217-2428>
M. C. Johnston  <https://orcid.org/0000-0002-0663-9193>
D. H. Jones  <https://orcid.org/0000-0003-3987-068X>
P. Joshi  <https://orcid.org/0000-0002-4148-4932>
L. Ju  <https://orcid.org/0000-0002-7951-4295>
K. Jung  <https://orcid.org/0000-0003-4789-8893>
J. Junker  <https://orcid.org/0000-0002-3051-4374>
H. B. Kabagoz  <https://orcid.org/0000-0002-0900-8557>
T. Kajita  <https://orcid.org/0000-0003-1207-6638>
V. Kalogera  <https://orcid.org/0000-0001-9236-5469>
M. Kalomenopoulos  <https://orcid.org/0000-0001-6677-949X>
M. Kamiizumi  <https://orcid.org/0000-0001-7216-1784>
N. Kanda  <https://orcid.org/0000-0001-6291-0227>
S. Kandhasamy  <https://orcid.org/0000-0002-4825-6764>
G. Kang  <https://orcid.org/0000-0002-6072-8189>
S. J. Kapadia  <https://orcid.org/0000-0001-5318-1253>
D. P. Kapasi  <https://orcid.org/0000-0001-8189-4920>
M. Kasprzak  <https://orcid.org/0000-0003-4618-5939>
R. Kaushik  <https://orcid.org/0000-0003-4888-5154>
D. Keitel  <https://orcid.org/0000-0002-2824-626X>
L. J. Kemperman  <https://orcid.org/0009-0009-5254-8397>
J. Kennington  <https://orcid.org/0000-0002-6899-3833>
R. Kesharwani  <https://orcid.org/0009-0002-2528-5738>
J. S. Key  <https://orcid.org/0000-0003-0123-7600>
F. Y. Khalili  <https://orcid.org/0000-0001-7068-2332>
F. Khan  <https://orcid.org/0000-0001-6176-853X>
N. M. Khusid  <https://orcid.org/0000-0001-9304-7075>
W. Kiendrebeogo  <https://orcid.org/0000-0002-9108-5059>
N. Kijbunchoo  <https://orcid.org/0000-0002-2874-1228>
K. Kim  <https://orcid.org/0000-0003-1653-3795>
M. H. Kim  <https://orcid.org/0009-0009-9894-3640>
S. Kim  <https://orcid.org/0000-0003-1437-4647>
Y.-M. Kim  <https://orcid.org/0000-0001-8720-6113>
C. Kimball  <https://orcid.org/0000-0001-9879-6884>
J. S. Kissel  <https://orcid.org/0000-0002-1702-9577>
A. M. Knee  <https://orcid.org/0000-0003-0703-947X>
N. Knust  <https://orcid.org/0000-0002-5984-5353>
S. M. Koehlenbeck  <https://orcid.org/0000-0002-3842-9051>
K. Kohri  <https://orcid.org/0000-0003-3764-8612>
K. Kokeyama  <https://orcid.org/0000-0002-2896-1992>
S. Koley  <https://orcid.org/0000-0002-5793-6665>
P. Kolitsidou  <https://orcid.org/0000-0002-6719-8686>
A. E. Koloniari  <https://orcid.org/0000-0002-0546-5638>
K. Komori  <https://orcid.org/0000-0002-4092-9602>
A. K. H. Kong  <https://orcid.org/0000-0002-5105-344X>
A. Kontos  <https://orcid.org/0000-0002-1347-0680>
M. Korobko  <https://orcid.org/0000-0002-3839-3909>
A. Koushik  <https://orcid.org/0000-0002-7638-4544>
N. Kouvatso  <https://orcid.org/0000-0002-5497-3401>
N. V. Krishnendu  <https://orcid.org/0000-0002-3483-7517>
A. Królak  <https://orcid.org/0000-0003-4514-7690>
J. Kubisz  <https://orcid.org/0000-0001-7258-8673>
S. Kulkarni  <https://orcid.org/0000-0001-8057-0203>
A. Kulur Ramamohan  <https://orcid.org/0000-0003-3681-1887>
Praveen Kumar  <https://orcid.org/0000-0002-2288-4252>
Prayush Kumar  <https://orcid.org/0000-0001-5523-4603>
J. Kume  <https://orcid.org/0000-0003-3126-5100>
K. Kuns  <https://orcid.org/0000-0003-0630-3902>
S. Kuroyanagi  <https://orcid.org/0000-0001-6538-1447>
S. Kuwahara  <https://orcid.org/0009-0009-2249-8798>
K. Kwak  <https://orcid.org/0000-0002-2304-7798>
S. Kwon  <https://orcid.org/0009-0006-3770-7044>
D. Laghi  <https://orcid.org/0000-0001-7462-3794>
M. Lalleman  <https://orcid.org/0000-0002-2254-010X>
R. N. Lang  <https://orcid.org/0000-0002-4804-5537>
R. Langgin  <https://orcid.org/0000-0002-5116-6217>
B. Lantz  <https://orcid.org/0000-0002-7404-4845>
I. L. Rosa  <https://orcid.org/0000-0003-0107-1540>
A. Lartaux-Vollard  <https://orcid.org/0000-0003-1714-365X>
P. D. Lasky  <https://orcid.org/0000-0003-3763-1386>
J. Lawrence  <https://orcid.org/0000-0003-1222-0433>
M. Laxen  <https://orcid.org/0000-0001-7515-9639>
C. Lazarte  <https://orcid.org/0000-0002-6964-9321>
A. Lazzarini  <https://orcid.org/0000-0002-5993-8808>
P. Leaci  <https://orcid.org/0000-0002-3997-5046>
Y. K. Lecoche  <https://orcid.org/0000-0002-9186-7034>
H. M. Lee  <https://orcid.org/0000-0003-4412-7161>
H. W. Lee  <https://orcid.org/0000-0002-1998-3209>
K. Lee  <https://orcid.org/0000-0003-0470-3718>
R.-K. Lee  <https://orcid.org/0000-0002-7171-7274>
Sungho Lee  <https://orcid.org/0000-0001-6034-2238>
M. Le Jean  <https://orcid.org/0009-0003-8047-3958>
A. Lemaitre  <https://orcid.org/0000-0002-6865-9245>
M. Lenti  <https://orcid.org/0000-0002-2765-3955>
M. Leonardi  <https://orcid.org/0000-0002-7641-0060>
N. Leroy  <https://orcid.org/0000-0002-2321-1017>
M. Lethuillier  <https://orcid.org/0000-0001-6185-2045>
K. L. Li  <https://orcid.org/0000-0001-8229-2024>
X. Li  <https://orcid.org/0000-0002-3780-7735>
E. T. Lin  <https://orcid.org/0000-0002-0030-8051>
L. C.-C. Lin  <https://orcid.org/0000-0003-4083-9567>
Y.-C. Lin  <https://orcid.org/0000-0003-4939-1404>
A. Liu  <https://orcid.org/0000-0003-1081-8722>
G. C. Liu  <https://orcid.org/0000-0001-5663-3016>
Jian Liu  <https://orcid.org/0000-0001-6726-3268>
J. Llobera-Querol  <https://orcid.org/0000-0003-3322-6850>
R. K. L. Lo  <https://orcid.org/0000-0003-1561-6716>
A. Longo  <https://orcid.org/0000-0003-4254-8579>
D. Lopez  <https://orcid.org/0000-0003-3342-9906>
M. Lorenzini  <https://orcid.org/0000-0002-2765-7905>
A. Lorenzo-Medina  <https://orcid.org/0009-0006-0860-5700>
G. Losurdo  <https://orcid.org/0000-0003-0452-746X>
T. P. Lott IV  <https://orcid.org/0009-0002-2864-162X>
J. D. Lough  <https://orcid.org/0000-0002-5160-0239>
C. O. Lousto  <https://orcid.org/0000-0002-6400-9640>
N. Lu  <https://orcid.org/0000-0002-8861-9902>
L. Lucchesi  <https://orcid.org/0000-0002-5916-8014>
D. Lumaca  <https://orcid.org/0000-0002-3628-1591>
A. P. Lundgren  <https://orcid.org/0000-0002-0363-4469>
A. W. Lussier  <https://orcid.org/0000-0002-4507-1123>
R. Macas  <https://orcid.org/0000-0002-6096-8297>
D. M. Macleod  <https://orcid.org/0000-0002-1395-8694>

- I. A. O. MacMillan  <https://orcid.org/0000-0002-6927-1031>
A. Macquet  <https://orcid.org/0000-0001-5955-6415>
S. Maenaut  <https://orcid.org/0000-0003-1464-2605>
R. M. Magee  <https://orcid.org/0000-0001-9769-531X>
E. Maggio  <https://orcid.org/0000-0002-1960-8185>
M. Magnozzi  <https://orcid.org/0000-0003-4512-8430>
D. Malakar  <https://orcid.org/0000-0003-4234-4023>
U. Mali  <https://orcid.org/0009-0003-1285-2788>
L. Mallick  <https://orcid.org/0000-0001-8624-9162>
A.-K. Malz  <https://orcid.org/0009-0004-7196-4170>
M. Mancarella  <https://orcid.org/0000-0002-0675-508X>
V. Mandic  <https://orcid.org/0000-0001-6333-8621>
V. Mangano  <https://orcid.org/0000-0001-7902-8505>
G. L. Mansell  <https://orcid.org/0000-0003-4736-6678>
M. Manske  <https://orcid.org/0000-0002-7778-1189>
M. Mantovani  <https://orcid.org/0000-0002-4424-5726>
M. Mapelli  <https://orcid.org/0000-0001-8799-2548>
C. Marinelli  <https://orcid.org/0000-0002-3596-4307>
F. Marion  <https://orcid.org/0000-0002-8184-1017>
S. Marsat  <https://orcid.org/0000-0001-9449-1071>
F. Martelli  <https://orcid.org/0000-0003-3761-8616>
I. W. Martin  <https://orcid.org/0000-0001-7300-9151>
R. M. Martin  <https://orcid.org/0000-0001-9664-2216>
V. Martinez  <https://orcid.org/0000-0001-5852-2301>
J. C. Martins  <https://orcid.org/0000-0002-6099-4831>
M. Masso-Reid  <https://orcid.org/0000-0001-6177-8105>
S. Mastrogiovanni  <https://orcid.org/0000-0003-1606-4183>
T. Matcovich  <https://orcid.org/0009-0004-1209-008X>
M. Matushechkina  <https://orcid.org/0000-0002-9957-8720>
N. Mavalvala  <https://orcid.org/0000-0003-0219-9706>
D. E. McClelland  <https://orcid.org/0000-0001-6210-5842>
L. McCuller  <https://orcid.org/0000-0003-0851-0593>
G. I. McGhee  <https://orcid.org/0000-0001-5038-2658>
J. McIver  <https://orcid.org/0000-0003-0316-1355>
A. McLeod  <https://orcid.org/0000-0001-5424-8368>
I. McMahon  <https://orcid.org/0000-0002-4529-1505>
R. McTeague  <https://orcid.org/0009-0004-3329-6079>
D. Meacher  <https://orcid.org/0000-0001-5882-0368>
C. S. Menoni  <https://orcid.org/0000-0001-9185-2572>
R. A. Mercer  <https://orcid.org/0000-0001-8372-3914>
J. R. Mérou  <https://orcid.org/0000-0002-5776-6643>
C. Messick  <https://orcid.org/0000-0002-8230-3309>
M. Meyer-Conde  <https://orcid.org/0000-0003-2230-6310>
F. Meylahn  <https://orcid.org/0000-0002-9556-142X>
A. Miani  <https://orcid.org/0000-0001-7737-3129>
C. Michel  <https://orcid.org/0000-0003-0606-725X>
Y. Michimura  <https://orcid.org/0000-0002-2218-4002>
H. Middleton  <https://orcid.org/0000-0001-5532-3622>
D. P. Mihaylov  <https://orcid.org/0000-0002-8820-407X>
S. J. Miller  <https://orcid.org/0000-0001-5670-7046>
M. Millhouse  <https://orcid.org/0000-0002-8659-5898>
E. Milotti  <https://orcid.org/0000-0001-7348-9765>
V. Milotti  <https://orcid.org/0000-0003-4732-1226>
Ll. M. Mir  <https://orcid.org/0000-0002-4276-715X>
L. Mirasola  <https://orcid.org/0009-0004-0174-1377>
M. Miravet-Tenés  <https://orcid.org/0000-0002-8766-1156>
C.-A. Miritescu  <https://orcid.org/0000-0002-7716-0569>
C. Mishra  <https://orcid.org/0000-0002-8115-8728>
T. Mishra  <https://orcid.org/0000-0002-7881-1677>
S. Mitra  <https://orcid.org/0000-0002-0800-4626>
V. P. Mitrofanov  <https://orcid.org/0000-0002-6983-4981>
O. Miyakawa  <https://orcid.org/0000-0002-9085-7600>
S. Miyoki  <https://orcid.org/0000-0002-1213-8416>
G. Mo  <https://orcid.org/0000-0001-6331-112X>
L. Mobilia  <https://orcid.org/0009-0000-3022-2358>
S. R. Mohite  <https://orcid.org/0000-0003-1356-7156>
M. Molina- Ruiz  <https://orcid.org/0000-0003-4892-3042>
A. More  <https://orcid.org/0000-0001-7714-7076>
S. More  <https://orcid.org/0000-0002-2986-2371>
C. Moreno  <https://orcid.org/0000-0002-0496-032X>
E. A. Moreno  <https://orcid.org/0000-0001-5666-3637>
S. Morisaki  <https://orcid.org/0000-0002-8445-6747>
Y. Moriwaki  <https://orcid.org/0000-0002-4497-6908>
G. Morras  <https://orcid.org/0000-0002-9977-8546>
A. Moscatello  <https://orcid.org/0000-0001-5480-7406>
M. Mould  <https://orcid.org/0000-0001-5460-2910>
B. Mours  <https://orcid.org/0000-0002-6444-6402>
C. M. Mow-Lowry  <https://orcid.org/0000-0002-0351-4555>
L. Muccillo  <https://orcid.org/0009-0000-6237-0590>
F. Muciaccia  <https://orcid.org/0000-0003-0850-2649>
D. Mukherjee  <https://orcid.org/0000-0001-7335-9418>
Suvodip Mukherjee  <https://orcid.org/0000-0002-3373-5236>
N. Mukund  <https://orcid.org/0000-0002-8666-9156>
P. G. Murray  <https://orcid.org/0000-0002-8218-2404>
D. Nabari  <https://orcid.org/0009-0006-8500-7624>
N. Nagarajan  <https://orcid.org/0000-0003-3695-0078>
K. Nakamura  <https://orcid.org/0000-0001-6148-4289>
H. Nakano  <https://orcid.org/0000-0001-7665-0796>
D. Nanadoumgar-Lacroze  <https://orcid.org/0009-0009-7255-8111>
P. Narayan  <https://orcid.org/0009-0009-0599-532X>
I. Nardecchia  <https://orcid.org/0000-0001-5558-2595>
L. Naticchioni  <https://orcid.org/0000-0003-2918-0730>
R. K. Nayak  <https://orcid.org/0000-0002-6814-7792>
A. Nelson  <https://orcid.org/0000-0002-5909-4692>
A. Neunzert  <https://orcid.org/0000-0002-0323-0111>
L. Nguyen Quynh  <https://orcid.org/0000-0002-1828-3702>
A. B. Nielsen  <https://orcid.org/0000-0001-8694-4026>
A. Nishizawa  <https://orcid.org/0000-0003-3562-0990>
W. Niu  <https://orcid.org/0000-0003-1470-532X>
J. Novak  <https://orcid.org/0000-0002-6029-4712>
R. Nowicki  <https://orcid.org/0009-0008-6626-0725>
J. F. Nuño Siles  <https://orcid.org/0000-0001-8304-8066>
L. K. Nuttall  <https://orcid.org/0000-0002-8599-8791>
J. Oberling  <https://orcid.org/0009-0001-4174-3973>
E. Oelker  <https://orcid.org/0000-0002-3916-1595>
M. Oertel  <https://orcid.org/0000-0002-1884-8654>
M. Ohashi  <https://orcid.org/0000-0001-8072-0304>
F. Ohme  <https://orcid.org/0000-0003-0493-5607>
R. Oliveri  <https://orcid.org/0000-0002-7497-871X>
K. Oohara  <https://orcid.org/0000-0002-7518-6677>
B. O'Reilly  <https://orcid.org/0000-0002-3874-8335>
M. Orsell  <https://orcid.org/0000-0003-3563-8576>
R. O'Shaughnessy  <https://orcid.org/0000-0001-5832-8517>
S. Oshino  <https://orcid.org/0000-0002-2794-6029>
I. Ota  <https://orcid.org/0000-0001-5045-2484>
D. J. Ottaway  <https://orcid.org/0000-0001-6794-1591>
B. J. Owen  <https://orcid.org/0000-0003-3919-0780>
A. E. Pace  <https://orcid.org/0009-0003-4044-0334>
R. Pagano  <https://orcid.org/0000-0001-8362-0130>
M. A. Page  <https://orcid.org/0000-0002-5298-7914>
A. Pai  <https://orcid.org/0000-0003-3476-4589>
S. Pal  <https://orcid.org/0000-0003-2172-8589>

- M. A. Palaia  <https://orcid.org/0009-0007-3296-8648>
 C. Palomba  <https://orcid.org/0000-0002-4450-9883>
 P. Palud  <https://orcid.org/0000-0002-5850-6325>
 K. C. Pan  <https://orcid.org/0000-0002-1473-9880>
 F. Pannarale  <https://orcid.org/0000-0002-7537-3210>
 F. Paoletti  <https://orcid.org/0000-0001-8898-1963>
 A. Paolone  <https://orcid.org/0000-0002-4839-7815>
 A. Papadopoulos  <https://orcid.org/0009-0006-1882-996X>
 L. Papalini  <https://orcid.org/0000-0002-5219-0454>
 G. Papigkiotis  <https://orcid.org/0009-0008-2205-7426>
 A. Parisi  <https://orcid.org/0000-0003-0251-8914>
 J. Park  <https://orcid.org/0000-0002-7510-0079>
 W. Parker  <https://orcid.org/0000-0002-7711-4423>
 D. Pascucci  <https://orcid.org/0000-0003-1907-0175>
 A. Pasqualetti  <https://orcid.org/0000-0003-0620-5990>
 R. Passaquieti  <https://orcid.org/0000-0003-4753-9428>
 O. Patane  <https://orcid.org/0000-0002-4850-2355>
 A. V. Patel  <https://orcid.org/0000-0001-6872-9197>
 B. Patricelli  <https://orcid.org/0000-0001-6709-0969>
 K. Paul  <https://orcid.org/0000-0002-8406-6503>
 S. Paul  <https://orcid.org/0000-0002-4449-1732>
 E. Payne  <https://orcid.org/0000-0003-4507-8373>
 A. Pele  <https://orcid.org/0000-0002-1873-3769>
 F. E. Peña Arellano  <https://orcid.org/0000-0002-8516-5159>
 S. Penn  <https://orcid.org/0000-0003-4956-0853>
 A. Perego  <https://orcid.org/0000-0002-0936-8237>
 C. Périgois  <https://orcid.org/0000-0002-9779-2838>
 G. Perna  <https://orcid.org/0000-0002-7364-1904>
 A. Perreca  <https://orcid.org/0000-0002-6269-2490>
 J. Perret  <https://orcid.org/0009-0006-4975-1536>
 S. Perriès  <https://orcid.org/0000-0003-2213-3579>
 H. P. Pfeiffer  <https://orcid.org/0000-0001-9288-519X>
 K. A. Pham  <https://orcid.org/0000-0002-7650-1034>
 K. S. Phukon  <https://orcid.org/0000-0003-1561-0760>
 L. Piccari  <https://orcid.org/0009-0000-0247-4339>
 O. J. Piccinni  <https://orcid.org/0000-0001-5478-3950>
 M. Pichot  <https://orcid.org/0000-0002-4439-8968>
 M. Piendibene  <https://orcid.org/0000-0003-2434-488X>
 F. Piergiovanni  <https://orcid.org/0000-0001-8063-828X>
 L. Pierini  <https://orcid.org/0000-0003-0945-2196>
 G. Pierra  <https://orcid.org/0000-0003-3970-7970>
 V. Pierro  <https://orcid.org/0000-0002-6020-5521>
 M. Pillas  <https://orcid.org/0000-0003-3224-2146>
 F. Pilo  <https://orcid.org/0000-0003-4967-7090>
 L. Pinard  <https://orcid.org/0000-0002-8842-1867>
 I. M. Pinto  <https://orcid.org/0000-0002-2679-4457>
 M. Pinto  <https://orcid.org/0009-0003-4339-9971>
 B. J. Piotrkowski  <https://orcid.org/0000-0001-8919-0899>
 M. D. Pitkin  <https://orcid.org/0000-0003-4548-526X>
 A. Placidi  <https://orcid.org/0000-0001-8032-4416>
 E. Placidi  <https://orcid.org/0000-0002-3820-8451>
 M. L. Planas  <https://orcid.org/0000-0001-8278-7406>
 W. Plastino  <https://orcid.org/0000-0002-5737-6346>
 C. Plunkett  <https://orcid.org/0000-0002-1144-6708>
 R. Poggiani  <https://orcid.org/0000-0002-9968-2464>
 L. Pompili  <https://orcid.org/0000-0002-0710-6778>
 C. Posnansky  <https://orcid.org/0009-0009-7137-9795>
 R. Poulton  <https://orcid.org/0000-0003-2049-520X>
 J. Powell  <https://orcid.org/0000-0002-1357-4164>
 M. Pracchia  <https://orcid.org/0009-0001-8343-719X>
 B. K. Pradhan  <https://orcid.org/0000-0002-2526-1421>
 T. Pradier  <https://orcid.org/0000-0001-5501-0060>
 K. Prasai  <https://orcid.org/0000-0001-6552-097X>
 G. Pratten  <https://orcid.org/0000-0003-4984-0775>
 G. Principe  <https://orcid.org/0000-0003-0406-7387>
 G. A. Prodi  <https://orcid.org/0000-0001-5256-915X>
 A. Puecher  <https://orcid.org/0000-0003-1357-4348>
 J. Pullin  <https://orcid.org/0000-0001-8248-603X>
 M. Pürner  <https://orcid.org/0000-0002-3329-9788>
 H. Qi  <https://orcid.org/0000-0001-6339-1537>
 J. Qin  <https://orcid.org/0000-0002-7120-9026>
 G. Quémeré  <https://orcid.org/0000-0001-6703-6655>
 B. Rajbhandari  <https://orcid.org/0000-0001-7568-1611>
 K. E. Ramirez  <https://orcid.org/0000-0003-2194-7669>
 F. A. Ramis Vidal  <https://orcid.org/0000-0001-6143-2104>
 M. Ramos Arevalo  <https://orcid.org/0009-0003-1528-8326>
 A. Ramos-Buades  <https://orcid.org/0000-0002-6874-7421>
 S. Ranjan  <https://orcid.org/0000-0001-7480-9329>
 P. Rapagnani  <https://orcid.org/0000-0002-1865-6126>
 A. Ray  <https://orcid.org/0000-0002-7322-4748>
 V. Raymond  <https://orcid.org/0000-0003-0066-0095>
 M. Razzano  <https://orcid.org/0000-0003-4825-1629>
 D. H. Reitze  <https://orcid.org/0000-0002-5756-1111>
 A. I. Renzini  <https://orcid.org/0000-0002-4589-3987>
 B. Revenu  <https://orcid.org/0000-0002-7629-4805>
 L. Ricca  <https://orcid.org/0009-0002-1638-0610>
 F. Ricci  <https://orcid.org/0000-0001-5475-4447>
 M. Ricci  <https://orcid.org/0009-0008-7421-4331>
 A. Ricciardone  <https://orcid.org/0000-0002-5688-455X>
 J. W. Richardson  <https://orcid.org/0000-0002-1472-4806>
 K. Riles  <https://orcid.org/0000-0002-6418-5812>
 S. Rinaldi  <https://orcid.org/0000-0001-5799-4155>
 A. Rocchi  <https://orcid.org/0000-0002-1382-9016>
 L. Rolland  <https://orcid.org/0000-0003-0589-9687>
 J. G. Rollins  <https://orcid.org/0000-0002-9388-2799>
 A. E. Romano  <https://orcid.org/0000-0002-0314-8698>
 R. Romano  <https://orcid.org/0000-0002-0485-6936>
 A. Romero  <https://orcid.org/0000-0003-2275-4164>
 S. Ronchini  <https://orcid.org/0000-0003-0020-687X>
 T. J. Rooke  <https://orcid.org/0000-0003-2640-9683>
 D. Rosińska  <https://orcid.org/0000-0002-3681-9304>
 M. P. Ross  <https://orcid.org/0000-0002-8955-5269>
 M. Rossello-Sastre  <https://orcid.org/0000-0002-3341-3480>
 S. Rowan  <https://orcid.org/0000-0002-0666-9907>
 S. K. Roy  <https://orcid.org/0000-0001-9295-5119>
 S. Roy  <https://orcid.org/0000-0003-2147-5411>
 D. Rozza  <https://orcid.org/0000-0002-7378-6353>
 E. Ruiz Morales  <https://orcid.org/0000-0002-0995-595X>
 S. Sachdev  <https://orcid.org/0000-0002-0525-2317>
 P. Saffarieh  <https://orcid.org/0009-0000-7504-3660>
 S. Safi-Harb  <https://orcid.org/0000-0001-6189-7665>
 M. R. Sah  <https://orcid.org/0009-0005-9881-1788>
 S. Saha  <https://orcid.org/0000-0002-3333-8070>
 T. Sainrat  <https://orcid.org/0009-0003-0169-266X>
 S. Sajith Menon  <https://orcid.org/0009-0008-4985-1320>
 Y. Sakai  <https://orcid.org/0000-0001-8810-4813>
 M. Sakellariadou  <https://orcid.org/0000-0002-2715-1517>
 S. Sakon  <https://orcid.org/0000-0002-5861-3024>
 O. S. Salafia  <https://orcid.org/0000-0003-4924-7322>
 F. Salces-Carcoba  <https://orcid.org/0000-0001-7049-4438>
 M. Saleem  <https://orcid.org/0000-0002-3836-7751>
 F. Salemi  <https://orcid.org/0000-0002-9511-3846>
 M. Sallé  <https://orcid.org/0000-0002-6620-6672>

- S. Salvador  <https://orcid.org/0000-0003-3444-7807>
A. Samajdar  <https://orcid.org/0000-0002-0857-6018>
N. Sanchis-Gual  <https://orcid.org/0000-0001-5375-7494>
E. M. Sanger  <https://orcid.org/0009-0003-6642-8974>
F. Santoliquido  <https://orcid.org/0000-0003-3752-1400>
A. Sasli  <https://orcid.org/0000-0001-7357-0889>
P. Sassi  <https://orcid.org/0000-0002-4920-2784>
B. Sassolas  <https://orcid.org/0000-0002-3077-8951>
O. Sauter  <https://orcid.org/0000-0003-2293-1554>
R. L. Savage  <https://orcid.org/0000-0003-3317-1036>
T. Sawada  <https://orcid.org/0000-0001-5726-7150>
M. G. Schiowski  <https://orcid.org/0000-0001-9298-004X>
P. Schmidt  <https://orcid.org/0000-0003-1542-1791>
S. Schmidt  <https://orcid.org/0000-0002-8206-8089>
R. Schnabel  <https://orcid.org/0000-0003-2896-4218>
K. Schouteden  <https://orcid.org/0000-0002-5975-585X>
E. Schwartz  <https://orcid.org/0000-0001-8922-7794>
M. Scialpi  <https://orcid.org/0009-0007-6434-1460>
J. Scott  <https://orcid.org/0000-0001-6701-6515>
S. M. Scott  <https://orcid.org/0000-0002-9875-7700>
R. M. Sedas  <https://orcid.org/0000-0001-8961-3855>
M. Seglar-Arroyo  <https://orcid.org/0000-0001-8654-409X>
Y. Sekiguchi  <https://orcid.org/0000-0002-2648-3835>
A. S. Sengupta  <https://orcid.org/0000-0002-3212-0475>
E. G. Seo  <https://orcid.org/0000-0002-8588-4794>
J. W. Seo  <https://orcid.org/0000-0003-4937-0769>
M. Serra  <https://orcid.org/0000-0002-6093-8063>
U. S. Shah  <https://orcid.org/0000-0001-8249-7425>
M. A. Shaikh  <https://orcid.org/0000-0003-0826-6164>
L. Shao  <https://orcid.org/0000-0002-1334-8853>
A. K. Sharma  <https://orcid.org/0000-0003-0067-346X>
P. Shawhan  <https://orcid.org/0000-0002-8249-8070>
N. S. Shcheblanov  <https://orcid.org/0000-0001-8696-2435>
H. Shinkai  <https://orcid.org/0000-0003-1082-2844>
D. H. Shoemaker  <https://orcid.org/0000-0002-4147-2560>
D. M. Shoemaker  <https://orcid.org/0000-0002-9899-6357>
H. Siegel  <https://orcid.org/0000-0001-5161-4617>
D. Sigg  <https://orcid.org/0000-0003-4606-6526>
L. Silenzi  <https://orcid.org/0000-0001-7316-3239>
L. Silvestri  <https://orcid.org/0009-0008-5207-661X>
L. P. Singer  <https://orcid.org/0000-0001-9898-5597>
D. Singh  <https://orcid.org/0000-0001-9675-4584>
N. Singh  <https://orcid.org/0000-0002-1135-3456>
A. M. Sintesi  <https://orcid.org/0000-0001-9050-7515>
V. Skliris  <https://orcid.org/0000-0003-0902-9216>
B. J. J. Slagmolen  <https://orcid.org/0000-0002-2471-3828>
J. R. Smith  <https://orcid.org/0000-0003-0638-9670>
L. Smith  <https://orcid.org/0000-0002-3035-0947>
R. J. E. Smith  <https://orcid.org/0000-0001-8516-3324>
W. J. Smith  <https://orcid.org/0009-0003-7949-4911>
K. Somiya  <https://orcid.org/0000-0003-2601-2264>
I. Song  <https://orcid.org/0000-0002-4301-8281>
S. Soni  <https://orcid.org/0000-0003-3856-8534>
V. Sordini  <https://orcid.org/0000-0003-0885-824X>
H. Sotani  <https://orcid.org/0000-0002-3239-2921>
F. Spada  <https://orcid.org/0000-0001-5664-1657>
V. Spagnuolo  <https://orcid.org/0000-0002-0098-4260>
A. P. Spencer  <https://orcid.org/0000-0003-4418-3366>
P. Spinicelli  <https://orcid.org/0000-0001-8078-6047>
F. Stachurski  <https://orcid.org/0000-0002-8658-5753>
D. A. Steer  <https://orcid.org/0000-0002-8781-1273>
N. Steinle  <https://orcid.org/0000-0003-0658-402X>
S. Steinlechner  <https://orcid.org/0000-0003-4710-8548>
N. Stergioulas  <https://orcid.org/0000-0002-5490-5302>
S. Sudhagar  <https://orcid.org/0000-0001-8578-4665>
L. Suleiman  <https://orcid.org/0000-0003-3783-7448>
J. Sun  <https://orcid.org/0009-0008-8278-0077>
L. Sun  <https://orcid.org/0000-0001-7959-892X>
J. Suresh  <https://orcid.org/0000-0003-2389-6666>
P. J. Sutton  <https://orcid.org/0000-0003-1614-3922>
B. L. Swinkels  <https://orcid.org/0000-0002-3066-3601>
A. Syx  <https://orcid.org/0009-0000-6424-6411>
M. J. Szczepańczyk  <https://orcid.org/0000-0002-6167-6149>
P. Szewczyk  <https://orcid.org/0000-0002-1339-9167>
M. Tacca  <https://orcid.org/0000-0003-1353-0441>
H. Tagoshi  <https://orcid.org/0000-0001-8530-9178>
S. C. Tait  <https://orcid.org/0000-0003-0327-953X>
H. Takahashi  <https://orcid.org/0000-0003-0596-4397>
R. Takahashi  <https://orcid.org/0000-0003-1367-5149>
A. Takamori  <https://orcid.org/0000-0001-6032-1330>
S. Takano  <https://orcid.org/0000-0002-1266-4555>
H. Takeda  <https://orcid.org/0000-0001-9937-2557>
N. Tamanini  <https://orcid.org/0000-0001-8760-5421>
S. J. Tanaka  <https://orcid.org/0000-0002-8796-1992>
S. Tanioka  <https://orcid.org/0000-0003-3321-1018>
L. Tao  <https://orcid.org/0000-0003-4382-5507>
E. N. Tapia San Martn  <https://orcid.org/0000-0002-4817-5606>
A. Taruya  <https://orcid.org/0000-0002-4016-1955>
J. D. Tasson  <https://orcid.org/0000-0002-4777-5087>
J. G. Tau  <https://orcid.org/0009-0004-7428-762X>
R. Tenorio  <https://orcid.org/0000-0002-3582-2587>
A. Theodoropoulos  <https://orcid.org/0000-0003-4486-7135>
L. M. Thomas  <https://orcid.org/0000-0003-3271-6436>
J. E. Thompson  <https://orcid.org/0000-0002-0419-5517>
E. Thrane  <https://orcid.org/0000-0002-4418-3895>
J. Tissino  <https://orcid.org/0000-0003-2483-6710>
S. Tiwari  <https://orcid.org/0000-0003-1611-6625>
V. Tiwari  <https://orcid.org/0000-0002-1602-4176>
A. M. Toivonen  <https://orcid.org/0009-0008-9546-2035>
K. Toland  <https://orcid.org/0000-0001-9537-9698>
A. E. Tolley  <https://orcid.org/0000-0001-9841-943X>
T. Tomaru  <https://orcid.org/0000-0002-8927-9014>
T. Tomura  <https://orcid.org/0000-0002-7504-8258>
H. Tong  <https://orcid.org/0000-0002-4534-0485>
A. Torres-Forne  <https://orcid.org/0000-0001-8709-5118>
I. Tosta e Melo  <https://orcid.org/0000-0001-5833-4052>
E. Tournefier  <https://orcid.org/0000-0002-5465-9607>
A. Trapananti  <https://orcid.org/0000-0001-7763-5758>
R. Travaglini  <https://orcid.org/0000-0002-5288-1407>
F. Travasso  <https://orcid.org/0000-0002-4653-6156>
M. C. Tringali  <https://orcid.org/0000-0001-5087-189X>
A. Tripathi  <https://orcid.org/0000-0002-6976-5576>
G. Troian  <https://orcid.org/0000-0001-6837-607X>
A. Trovato  <https://orcid.org/0000-0002-9714-1904>
T. Tsang  <https://orcid.org/0000-0003-3666-686X>
S. Tsuchida  <https://orcid.org/0000-0001-8217-0764>
L. Tsukada  <https://orcid.org/0000-0003-0596-5648>
K. Turbang  <https://orcid.org/0000-0002-9296-8603>
M. Turconi  <https://orcid.org/0000-0001-9999-2027>
H. Ubach  <https://orcid.org/0000-0002-0679-9074>
T. Uchiyama  <https://orcid.org/0000-0003-2148-1694>
R. P. Udall  <https://orcid.org/0000-0001-6877-3278>

- T. Uehara  <https://orcid.org/0000-0003-4375-098X>
 K. Ueno  <https://orcid.org/0000-0003-3227-6055>
 V. Undheim  <https://orcid.org/0000-0003-4028-0054>
 T. Ushiba  <https://orcid.org/0000-0002-5059-4033>
 M. Vacatello  <https://orcid.org/0009-0006-0934-1014>
 H. Vahlbruch  <https://orcid.org/0000-0003-2357-2338>
 N. Vaidya  <https://orcid.org/0000-0003-1843-7545>
 G. Vajente  <https://orcid.org/0000-0002-7656-6882>
 J. Valencia  <https://orcid.org/0000-0003-2648-9759>
 M. Valentini  <https://orcid.org/0000-0003-1215-4552>
 S. A. Vallejo-Peña  <https://orcid.org/0000-0002-6827-9509>
 V. Valsan  <https://orcid.org/0000-0003-0315-4091>
 M. van Dael  <https://orcid.org/0000-0002-6061-8131>
 E. Van den Bossche  <https://orcid.org/0009-0009-2070-0964>
 J. F. J. van den Brand  <https://orcid.org/0000-0003-4434-5353>
 M. van der Sluys  <https://orcid.org/0000-0003-1231-0762>
 J. van Dongen  <https://orcid.org/0000-0003-0964-2483>
 H. van Haeveermaet  <https://orcid.org/0000-0003-2386-957X>
 J. V. van Heijningen  <https://orcid.org/0000-0002-8391-7513>
 P. Van Hove  <https://orcid.org/0000-0002-2431-3381>
 N. van Remortel  <https://orcid.org/0000-0003-4180-8199>
 A. F. Vargas  <https://orcid.org/0000-0001-8396-5227>
 V. Varma  <https://orcid.org/0000-0002-9994-1761>
 A. Vecchio  <https://orcid.org/0000-0002-6254-1617>
 J. Veitch  <https://orcid.org/0000-0002-6508-0713>
 P. J. Veitch  <https://orcid.org/0000-0002-2597-435X>
 R. C. Venterea  <https://orcid.org/0000-0003-3299-3804>
 P. Verdier  <https://orcid.org/0000-0003-3090-2948>
 D. Verkindt  <https://orcid.org/0000-0003-4344-7227>
 Y. Verma  <https://orcid.org/0000-0003-4147-3173>
 S. M. Vermeulen  <https://orcid.org/0000-0003-4227-8214>
 A. Vestro  <https://orcid.org/0009-0002-9160-5808>
 A. Viceré  <https://orcid.org/0000-0003-0624-6231>
 A. D. Viets  <https://orcid.org/0000-0002-4241-1428>
 A. Vijaykumar  <https://orcid.org/0000-0002-4103-0666>
 V. Villa-Ortega  <https://orcid.org/0000-0001-7983-1963>
 E. T. Vincent  <https://orcid.org/0000-0002-0442-1916>
 S. Vitale  <https://orcid.org/0000-0003-2700-0767>
 H. Vocca  <https://orcid.org/0000-0002-1200-3917>
 D. Voigt  <https://orcid.org/0000-0001-9075-6503>
 L. Vujeva  <https://orcid.org/0000-0001-7697-8361>
 S. P. Vyatchanin  <https://orcid.org/0000-0002-6823-911X>
 M. Wade  <https://orcid.org/0000-0002-5703-4469>
 K. J. Wagner  <https://orcid.org/0000-0002-7255-4251>
 H. Wang  <https://orcid.org/0000-0002-6589-2738>
 Y. F. Wang  <https://orcid.org/0000-0002-2928-2916>
 G. Waratkar  <https://orcid.org/0000-0003-3630-9440>
 M. Was  <https://orcid.org/0000-0002-1890-1128>
 T. Washimi  <https://orcid.org/0000-0001-5792-4907>
 N. L. Weickhardt  <https://orcid.org/0000-0002-3923-5806>
 A. J. Weinstein  <https://orcid.org/0000-0002-0928-6784>
 L. Wen  <https://orcid.org/0000-0001-7987-295X>
 K. Wette  <https://orcid.org/0000-0002-4394-7179>
 J. T. Whelan  <https://orcid.org/0000-0001-5710-6576>
 B. F. Whiting  <https://orcid.org/0000-0002-8501-8669>
 C. Whittle  <https://orcid.org/0000-0002-8833-7438>
 D. Wilken  <https://orcid.org/0000-0002-7290-9411>
 D. Williams  <https://orcid.org/0000-0003-3772-198X>
 M. J. Williams  <https://orcid.org/0000-0003-2198-2974>
 N. S. Williams  <https://orcid.org/0000-0002-5656-8119>
 J. L. Willis  <https://orcid.org/0000-0002-9929-0225>
 B. Willke  <https://orcid.org/0000-0003-0524-2925>
 M. Wils  <https://orcid.org/0000-0002-1544-7193>
 G. Woan  <https://orcid.org/0000-0003-0381-0394>
 H. T. Wong  <https://orcid.org/0000-0003-4145-4394>
 I. C. F. Wong  <https://orcid.org/0000-0003-2166-0027>
 M. Wright  <https://orcid.org/0000-0003-1829-7482>
 C. Wu  <https://orcid.org/0000-0003-3191-8845>
 D. S. Wu  <https://orcid.org/0000-0003-2849-3751>
 H. Wu  <https://orcid.org/0000-0003-4813-3833>
 Z. Wu  <https://orcid.org/0000-0002-0032-5257>
 D. M. Wysocki  <https://orcid.org/0000-0001-9138-4078>
 V. A. Xu  <https://orcid.org/0000-0002-3020-3293>
 Y. Xu  <https://orcid.org/0000-0001-8697-3505>
 N. Yadav  <https://orcid.org/0009-0009-5010-1065>
 H. Yamamoto  <https://orcid.org/0000-0001-6919-9570>
 K. Yamamoto  <https://orcid.org/0000-0002-3033-2845>
 T. S. Yamamoto  <https://orcid.org/0000-0002-8181-924X>
 T. Yamamoto  <https://orcid.org/0000-0002-0808-4822>
 R. Yamazaki  <https://orcid.org/0000-0002-1251-7889>
 K. Z. Yang  <https://orcid.org/0000-0001-8083-4037>
 Y. Yang  <https://orcid.org/0000-0002-3780-1413>
 Z. Yarbrough  <https://orcid.org/0000-0002-9825-1136>
 A. B. Yeliker  <https://orcid.org/0000-0002-8065-1174>
 J. Yokoyama  <https://orcid.org/0000-0001-7127-4808>
 H. Yuzurihara  <https://orcid.org/0000-0002-3710-6613>
 M. Zeeshan  <https://orcid.org/0000-0002-6494-7303>
 M. Zeoli  <https://orcid.org/0009-0007-1898-4844>
 M. Zevin  <https://orcid.org/0000-0002-0147-0835>
 R. Zhang  <https://orcid.org/0000-0001-8095-483X>
 C. Zhao  <https://orcid.org/0000-0001-5825-2401>
 Z.-C. Zhao  <https://orcid.org/0000-0001-5180-4496>
 Y. Zheng  <https://orcid.org/0000-0002-5432-1331>
 H. Zhong  <https://orcid.org/0000-0001-8324-5158>
 Z.-H. Zhu  <https://orcid.org/0000-0002-3567-6743>
 A. B. Zimmerman  <https://orcid.org/0000-0002-7453-6372>
 M. E. Zucker  <https://orcid.org/0000-0002-2544-1596>
 J. Zweizig  <https://orcid.org/0000-0002-1521-3397>

References

- Aasi, J., Abbott, B., Abbott, R., et al. 2015a, *ApJ*, 813, 39
 Aasi, J., Abbott, B. P., Abbott, R., et al. 2015b, *CQGra*, 32, 074001
 Abac, A. G., Abouelfettouh, I., Acernese, F., et al. 2025a, arXiv:2508.18079
 Abac, A. G., Abouelfettouh, I., Acernese, F., et al. 2025b, arXiv:2508.20721
 Abac, A. G., Abouelfettouh, I., Acernese, F., et al. 2025c, arXiv:2510.17487
 Abadie, J., Abbott, B. P., Abbott, R., et al. 2010, *ApJ*, 722, 1504
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017, *PhRvL*, 118, 121102
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2019a, *PhRvX*, 9, 031040
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2019b, *ApJ*, 875, 122
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2019c, *PhRvD*, 100, 062001
 Abbott, R., Abbott, T. D., Abraham, S., et al. 2021a, *PhRvX*, 11, 021053
 Abbott, R., Abbott, T. D., Abraham, S., et al. 2021b, *ApJ*, 921, 80
 Abbott, R., Abbott, T. D., Abraham, S., et al. 2021c, *PhRvD*, 104, 022005
 Abbott, R., Abbott, T. D., Abraham, S., et al. 2021d, *PhRvD*, 104, 082004
 Abbott, R., Abbott, T. D., Acernese, F., et al. 2022a, *PhRvD*, 105, 082005
 Abbott, R., Abe, H., Acernese, F., et al. 2022b, *PhRvD*, 106, 042003
 Abbott, R., Abbott, T. D., Acernese, F., et al. 2023, *PhRvX*, 13, 041039
 Abbott, R., Abbott, T. D., Acernese, F., et al. 2024, *PhRvD*, 109, 022001
 Abbott, R., Abouelfettouh, I., Acernese, F., et al. 2025, arXiv:2508.18082
 Acernese, F., Alshourbagy, M., Antonucci, F., et al. 2009, *CQGra*, 26, 204002
 Ain, A., Suresh, J., & Mitra, S. 2018, *PhRvD*, 98, 024001
 Amicucci, F., Leaci, P., Astone, P., et al. 2025, *CQGra*, 42, 145008
 Andersson, N. 1998, *ApJ*, 502, 708
 Antonucci, F., Astone, P., D'Antonio, S., Frasca, S., & Palomba, C. 2008, *CQGra*, 25, 184015
 Astone, P., Colla, A., D'Antonio, S., Frasca, S., & Palomba, C. 2014a, *PhRvD*, 90, 042002
 Astone, P., Colla, A., D'Antonio, S., et al. 2014b, *PhRvD*, 89, 062008

- Astone, P., D’Antonio, S., Frasca, S., & Palomba, C. 2010, *CQGra*, **27**, 194016
- Astone, P., Frasca, S., & Palomba, C. 2005, *CQGra*, **22**, 1197
- Behnke, B., Papa, M. A., & Prix, R. 2015, *PhRvD*, **91**, 064007
- Benival, D., Clearwater, P., Dunn, L., et al. 2022, *PhRvD*, **106**, 103018
- Biwer, C., Barker, D., Batch, J. C., et al. 2017, *PhRvD*, **95**, 062002
- Bondarescu, R., Teukolsky, S. A., & Wasserman, I. 2009, *PhRvD*, **79**, 104003
- Borkowski, K. J., Reynolds, S. P., Hwang, U., et al. 2013, *ApJL*, **771**, L9
- Braun, C., Safi-Harb, S., Fryer, C. L., & Zhou, P. 2023, *MNRAS*, **525**, 6257
- Bu, Y., & Melatos, A. 2025, Details of Parameter Choices for the O4a Supernova Remnant Search Using the Single-Harmonic Viterbi Pipeline LIGO-T2500368, LIGO-Virgo-KAGRA Collaboration <https://dcc.ligo.org/LIGO-T2500368/public>
- Capote, E., Jia, W., Aritomi, N., et al. 2025, *PhRvD*, **111**, 062002
- Caride, S., Inta, R., Owen, B. J., & Rajbhandari, B. 2019, *PhRvD*, **100**, 064013
- Cheunichitra, T., Melatos, A., Carlin, J. B., & Howitt, G. 2024, *MNRAS*, **528**, 1360
- Cheung, D. 2025, Outlier parameters from the Weave pipeline in the study “Searches for Continuous Gravitational Waves from Supernova Remnants in the first part of the LIGO-Virgo-KAGRA Fourth Observing run” v1, Zenodo, doi:10.5281/zenodo.14232423
- Cheung, D. H. T., Riles, K., Amengual, R., et al. 2026, arXiv:2603.04792
- Cordes, J. M., & Greenstein, G. 1981, *ApJ*, **245**, 1060
- Cutler, C. 2002, *PhRvD*, **66**, 084025
- Davis, D., Neunzert, A., Goetz, E., et al. 2024, Self-gating of O4a h(t) for use in continuous-wave searches T2400003, LIGO-Virgo-KAGRA Collaboration <https://dcc.ligo.org/T2400003>
- De Luca, A. 2017, *JPhCS*, **932**, 012006
- Dreissigacker, C., Prix, R., & Wette, K. 2018, *PhRvD*, **98**, 084058
- Ferrand, G., & Safi-Harb, S. 2012, *AdSpR*, **49**, 1313
- Fransson, C., Barlow, M. J., Kavanagh, P. J., et al. 2024, *Sci*, **383**, 898
- Gaensler, B. M., & Slane, P. O. 2006, *ARA&A*, **44**, 17
- Ganapathy, D., Jia, W., Nakano, M., et al. 2023, *PhRvX*, **13**, 041021
- Gittins, F. 2024, *CQGra*, **41**, 043001
- Gittins, F., & Andersson, N. 2023, *MNRAS*, **521**, 3043
- Glampedakis, K., & Gualtieri, L. 2018, in *Gravitational Waves from Single Neutron Stars: An Advanced Detector Era Survey*, ed. L. Rezzolla et al. (Springer International Publishing), 673
- Gnedin, O. Y., Yakovlev, D. G., & Potekhin, A. Y. 2001, *MNRAS*, **324**, 725
- Goetz, E., & Riles, K. 2024, Segments used for creating standard SFTs in O4 data T2400058, LIGO-Virgo-KAGRA Collaboration <https://dcc.ligo.org/T2400058>
- Goetz, E., Neunzert, A., Knee, A., et al. 2024, O4a lines and combs in found in self-gated C00 cleaned data T2400204, LIGO-Virgo-KAGRA Collaboration <https://dcc.ligo.org/T2400204>
- Gottlieb, E. V., Halpern, J. P., & Alford, J. 2013, *ApJ*, **765**, 58
- Greco, E., Miceli, M., Orlando, S., et al. 2021, *ApJL*, **908**, L45
- Green, D. A. 2025, yCat, VII/297, Originally published in: 2025
- Haensel, P., Urpin, V. A., & Yakovlev, D. G. 1990, *A&A*, **229**, 133
- Hamil, O., Stone, J., Urbanec, M., & Urbancová, G. 2015, *PhRvD*, **91**, 063007
- Haskell, B., Antonelli, M., & Pizzochero, P. 2022, *Univ*, **8**, 619
- Ho, W. C. G., & Lai, D. 2000, *ApJ*, **543**, 386
- Hurley-Walker, N., Gaensler, B. M., Leahy, D. A., et al. 2019, *PASA*, **36**, e048
- Idrisy, A., Owen, B. J., & Jones, D. I. 2015, *PhRvD*, **91**, 024001
- Jaranowski, P., Królak, A., & Schutz, B. F. 1998, *PhRvD*, **58**, 063001
- Jia, W., Xu, V., Kuns, K., et al. 2024, *Sci*, **385**, 1318
- Johnson-McDaniel, N. K., & Owen, B. J. 2013, *PhRvD*, **88**, 044004
- Jones, D. I. 2010, *MNRAS*, **402**, 2503
- Knispel, B., & Allen, B. 2008, *PhRvD*, **78**, 044031
- Lasky, P. D., & Melatos, A. 2013, *PhRvD*, **88**, 103005
- LIGO Scientific Collaboration, Virgo Collaboration, and KAGRA Collaboration 2026, Upper limits for the O4a Supernova Remnant search, v1, Zenodo, doi:10.5281/zenodo.19567772
- LIGO-Virgo-KAGRA Collaboration, et al. 2025, LIGO Virgo KAGRA Calibration Uncertainty (O4) T2500288, LIGO-Virgo-KAGRA Collaboration <https://dcc.ligo.org/T2500288/public>
- Lindblom, L., & Owen, B. J. 2020, *PhRvD*, **101**, 083023
- Lower, M. E., Bailes, M., Shannon, R. M., et al. 2020, *MNRAS*, **494**, 228
- Melatos, A., Douglass, J. A., & Simula, T. P. 2015, *ApJ*, **807**, 132
- Melatos, A., & Link, B. 2014, *MNRAS*, **437**, 21
- Millhouse, M., Strang, L., & Melatos, A. 2020, *PhRvD*, **102**, 083025
- Ming, J., Papa, M. A., Eggenstein, H.-B., et al. 2022, *ApJ*, **925**, 8
- Ming, J., Papa, M. A., Eggenstein, H.-B., et al. 2024, *ApJ*, **977**, 154
- Ming, J., Papa, M. A., Eggenstein, H.-B., et al. 2025, arXiv:2512.15672
- Ming, J., Papa, M. A., Singh, A., et al. 2019, *PhRvD*, **100**, 024063
- Morales, J., Ming, J., Papa, M. A., Eggenstein, H.-B., & Machenschalk, B. 2025, *ApJ*, **986**, 202
- Namkham, N., Jaroenjittichai, P., & Johnston, S. 2019, *MNRAS*, **487**, 5854
- Nitz, A. H., Kumar, S., Wang, Y.-F., et al. 2023, *ApJ*, **946**, 59
- Olsen, S., Venumadhav, T., Mushkin, J., et al. 2022, *PhRvD*, **106**, 043009
- Owen, B. J. 2010, *PhRvD*, **82**, 104002
- Owen, B. J. 2026, *RvMP*, **98**, 011002
- Owen, B. J., Lindblom, L., Cutler, C., et al. 1998, *PhRvD*, **58**, 084020
- Owen, B. J., Lindblom, L., & Pinheiro, L. S. 2022, *ApJL*, **935**, L7
- Owen, B. J., Lindblom, L., Soares Pinheiro, L., & Rajbhandari, B. 2024, *The ApJL*, **962**, L23
- Palomba, C. 2005, *MNRAS*, **359**, 1150
- Palomba, C. 2025, On the Sensitivity of Peakmap-based methods for the Search of Continuous Gravitational Wave Signals VIR-0724A-25, Virgo Collaboration <https://tds.virgo-gw.eu/?r=25026>
- Papa, M. A., Ming, J., Gottlieb, E. V., et al. 2020, *ApJ*, **897**, 22
- Parthasarathy, A., Shannon, R. M., Johnston, S., et al. 2019, *MNRAS*, **489**, 3810
- Pavlov, G. G., Zavlin, V. E., Aschenbach, B., Trümper, J., & Sanwal, D. 2000, *ApJL*, **531**, L53
- Piccinni, O. J. 2022, *Galax*, **10**, 72
- Piccinni, O. J., Astone, P., D’Antonio, S., et al. 2018, *CQGra*, **36**, 015008
- Piccinni, O. J., Astone, P., D’Antonio, S., et al. 2020, *PhRvD*, **101**, 082004
- Potekhin, A. Y., Pons, J. A., & Page, D. 2015, *SSRv*, **191**, 239
- Ravi, A. P., Park, S., Zhekov, S. A., et al. 2024, *ApJ*, **966**, 147
- Renzini, A. I., Romero-Rodríguez, A., Talbot, C., et al. 2023, *ApJ*, **952**, 25
- Reynolds, S. P., Borkowski, K. J., Green, D. A., et al. 2008, *ApJL*, **680**, L41
- Riles, K. 2023, *LRR*, **26**, 3
- Salvadore, C., L. Rosa, I., Leaci, P., et al. 2025, *PhRvD*, **112**, 083051
- Sieniawska, M., & Beijer, M. 2019, *Univ*, **5**, 217
- Singhal, A., Leaci, P., Astone, P., et al. 2019, *CQGra*, **36**, 205015
- Soni, S., Berger, B. K., Davis, D., et al. 2025, *CQGra*, **42**, 085016
- Suherli, J., Seitzenzahl, I. R., Safi-Harb, S., et al. 2026, *ApJL*, **997**, L36
- Sun, L., Goetz, E., Kissel, J. S., et al. 2020, *CQGra*, **37**, 225008
- Sun, L., Melatos, A., & Lasky, P. D. 2019, *PhRvD*, **99**, 123010
- Sun, L., Melatos, A., Lasky, P. D., Chung, C. T. Y., & Darman, N. S. 2016, *PhRvD*, **94**, 082004
- Sun, L., Melatos, A., Suvorova, S., Moran, W., & Evans, R. J. 2018, *PhRvD*, **97**, 043013
- Suvorova, S., Sun, L., Melatos, A., Moran, W., & Evans, R. 2016, *PhRvD*, **93**, 123009
- LIGO Scientific Collaboration 2022, SFT Data Format Version 2–3 Specification. LIGO-T040164, LIGO-Virgo-KAGRA Collaboration <https://dcc.ligo.org/LIGO-T040164/public>
- Treyturić, C., Braun, C., Safi-Harb, S., Fryer, C. L., & Ferrand, G. 2026, *ApJ*, **997**, 211
- Ushomirsky, G., Cutler, C., & Bildsten, L. 2000, *MNRAS*, **319**, 902
- Vajente, G., Huang, Y., Isi, M., et al. 2020, *PhRvD*, **101**, 042003
- Van Den Broeck, C. 2005, *CQGra*, **22**, 1825
- Viets, A. D. 2019, PhD thesis, Univ. Wisconsin-Milwaukee, Milwaukee, WI <http://digital.library.wisc.edu/1793/92045>
- Viets, A. D., Wade, M., Urban, A. L., et al. 2018, *CQGra*, **35**, 095015
- Wade, M., Betzwieser, J., Bhattacharjee, D., et al. 2025, *CQGra*, **42**, 215016
- Wang, J., & Riles, K. 2024, *PhRvD*, **110**, 042006
- Wette, K. 2014, *PhRvD*, **90**, 122010
- Wette, K. 2015, *PhRvD*, **92**, 082003
- Wette, K. 2023, *APH*, **153**, 102880
- Wette, K., & Prix, R. 2013, *PhRvD*, **88**, 123005
- Wette, K., Owen, B. J., Allen, B., et al. 2008, *CQGra*, **25**, 235011
- Wette, K., Walsh, S., Prix, R., & Papa, M. A. 2018, *PhRvD*, **97**, 123016
- Zimmermann, M., & Szedenits, E., Jr. 1979, *PhRvD*, **20**, 351