



Origins of a new economic union (7th-12th centuries)

Preliminary results of the nEU-Med project:
October 2015-March 2017

edited by

Giovanna Bianchi, Richard Hodges



BIBLIOTECA DI
**Archeologia
Medievale**

25

ARCHEOLOGIA MEDIEVALE

Cultura materiale. Insediamenti. Territorio.

Rivista fondata da Riccardo Francovich

Comitato di Direzione

SAURO GELICHI (responsabile) (Dipartimento di Studi Umanistici – Università Ca' Foscari di Venezia)

GIAN PIETRO BROGIOLO (già Università degli Studi di Padova)

Comitato Scientifico

LANFREDO CASTELLETTI (già Direttore dei Musei Civici di Como)

ANTONIO MALPICA CUELLO (Departamento de Historia – Universidad de Granada)

CARLO VARALDO (Dipartimento di antichità, filosofia, storia, geografia – Università degli Studi di Genova)

RINALDO COMBA (già Università degli Studi di Milano)

GHISLAINE NOYÉ (École nationale des chartes)

CHRIS WICKHAM (già Faculty of History – University of Oxford)

PAOLO DELOGU (Professore emerito, Sapienza Università di Roma)

PAOLO PEDUTO (già Università degli Studi di Salerno)

JUAN ANTONIO QUIRÓS CASTILLO (Departamento de Geografía, Prehistoria y Arqueología de la Universidad del País Vasco)

RICHARD HODGES (President of the American University of Rome)

Redazione

ANDREA AUGENTI (Dipartimento di Storia Culture Civiltà – Università degli Studi di Bologna)

CRISTINA LA ROCCA (Dipartimento di Scienze storiche, geografiche e dell'antichità – Università degli Studi di Padova)

SERGIO NEPOTI (responsabile sezione scavi in Italia) (Archeologo libero professionista)

GIOVANNA BIANCHI (Dipartimento di Scienze Storiche e dei Beni Culturali – Università degli Studi di Siena)

MARCO MILANESE (Dipartimento di Storia, Scienze dell'uomo e della Formazione – Università degli Studi di Sassari)

ALDO A. SETTIA (già Università degli Studi di Pavia)

ENRICO GIANNICCHEDDA (Istituto per la Storia della Cultura Materiale di Genova [ISCuM])

ALESSANDRA MOLINARI (Dipartimento di Storia – Università degli Studi di Roma Tor Vergata)

MARCO VALENTI (Dipartimento di Scienze Storiche e dei Beni Culturali – Università degli Studi di Siena)

GUIDO VANNINI (Dipartimento di Storia, Archeologia, Geografia, Arte e Spettacolo – Università degli Studi di Firenze)

Corrispondenti

PAUL ARTHUR (Dipartimento di Beni Culturali – Università degli Studi di Lecce)

CATERINA GIOSTRA (Dipartimento di Storia, archeologia e storia dell'arte – Università Cattolica del Sacro Cuore)

PHILIPPE PERGOLA (LAM3 – Laboratoire d'Archéologie Médiévale et Moderne en Méditerranée – Université d'Aix-Marseille CNRS/Pontificio istituto di archeologia cristiana)

VOLKER BIERBRAUER (Professore emerito, Ludwig-Maximilians-Universität München)

FEDERICO MARAZZI (Dipartimento di Scienze Storiche e dei Beni Culturali – Università degli Studi Suor Orsola Benincasa)

RENATO PERINETTI (già Soprintendenza per i Beni e le Attività Culturali della Regione Autonoma Valle d'Aosta)

HUGO BLAKE (già Royal Holloway – University of London)

ROBERTO MENEHINI (Sovrintendenza Capitolina ai Beni Culturali)

GIULIANO PINTO (già Università degli Studi di Firenze)

MAURIZIO BUORA (Società friulana di archeologia)

EGLÉ MICHELETTI (direttore della Soprintendenza Archeologia, Belle Arti e Paesaggio per le Province di Alessandria, Asti e Cuneo)

MARCELLO ROTILI (Seconda Università degli Studi di Napoli)

FEDERICO CANTINI (Dipartimento di Civiltà e Forme del Sapere – Università degli Studi di Pisa)

MASSIMO MONTANARI (Dipartimento di Storia Culture Civiltà – Università degli Studi di Bologna)

DANIELA ROVINA (Soprintendenza Archeologia, Belle Arti e Paesaggio per le Province di Sassari, Olbia-Tempio e Nuoro)

GISELLA CANTINO WATAGHIN (già Università del Piemonte Orientale)

GIOVANNI MURIALDO (Museo Archeologico del Finale – Finale Ligure Borgo SV)

LUCIA SAGUÌ (già Sapienza Università di Roma)

ENRICO CAVADA (Soprintendenza per i beni librari, archivistici e archeologici – Trento)

CLAUDIO NEGRELLI (Dipartimento di Studi Umanistici – Università Ca' Foscari di Venezia)

PIERGIORGIO SPANU (Dipartimento di Storia, Scienze dell'uomo e della Formazione – Università degli Studi di Sassari)

NEIL CHRISTIE (School of Archaeology and Ancient History – University of Leicester)

MICHELE NUCCIOTTI (Dipartimento di Storia, Archeologia, Geografia, Arte e Spettacolo – Università degli Studi di Firenze)

ANDREA R. STAFFA (Soprintendenza Archeologia, Belle Arti e Paesaggio dell'Abruzzo)

MAURO CORTELAZZO (Archeologo libero professionista)

GABRIELLA PANTÒ (Musei Reali di Torino – Museo di Antichità)

DANIELA STIAFFINI (Archeologa libera professionista)

FRANCESCO CUTERI (AISB, Associazione Italiana Studi Bizantini)

HELEN PATTERSON (già British School at Rome)

STANISŁAW TABACZYŃSKI (Polskiej Akademii Nauk)

LORENZO DAL RI (già Direttore ufficio Beni archeologici – Provincia autonoma di Bolzano Alto Adige)

LUISELLA PEJRANI BARICCO (già Soprintendenza Archeologia del Piemonte e del Museo Antichità Egizie)

BRYAN WARD PERKINS (History Faculty – Trinity College University of Oxford)

FRANCO D'ANGELO (già Direttore del Settore Cultura e della Tutela dell'Ambiente della Provincia di Palermo)

ALESSANDRA FRONDONI (già Soprintendenza Archeologia della Liguria)

UNIVERSITÀ DEGLI STUDI DI SIENA



nEU-Med

Origins of a new economic union (7th-12th centuries)

Preliminary results of the nEU-Med project:
October 2015-March 2017

edited by

Giovanna Bianchi, Richard Hodges

with contributions by

Andrea Bardi, Marco Benvenuti, Giovanna Bianchi, Andrea Biondi,
Arianna Briano, Mauro Paolo Buonincontri, Mirko Buono, Laura Chiarantini,
Cristina Cicali, Simone M. Collavini, Luisa Dallai, Gaetano Di Pasquale,
Alessandro Donati, Stefania Fineschi, Cristina Fornacelli, Simon Greenslade,
Richard Hodges, Sarah Leppard, Carmine Lubritto, Lorenzo Marasco,
Pierluigi Pieruccini, Giulio Poggi, Elisabetta Ponta, Paola Ricci,
Marta Rossi, Alessia Rovelli, Luisa Russo, Davide Susini, Paolo Tomei,
Emanuele Vaccaro, Igor Villa, Vanessa Volpi

Cover: Aerial view of excavations at Vetricella (Scarlino, GR) (photo ATS s.r.l.).

Unless otherwise specified, all photos are by the authors of the individual articles.

Italian to English translation: Gavin Williams

English to Italian translation for 'Towards a new Mediterranean narrative for Early Medieval Tuscany' by Richard Hodges:
Alexander Agostini

This volume has been subjected to double-blind peer review.

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement n. 670792)



ISSN 2035-5319
ISBN 978-88-7814-879-6
e-ISBN 978-88-7814-880-2
© 2018 All'Insegna del Giglio s.a.s. – Firenze

Edizioni All'Insegna del Giglio s.a.s
via del Termine, 36; 50019 Sesto Fiorentino (FI)
tel. +39 055 8450 216; fax +39 055 8453 188
e-mail redazione@insegnadelgiglio.it; ordini@insegnadelgiglio.it
sito web www.insegnadelgiglio.it

Printed in Florence, October 2018
Tecnografica Rossi

CONTENTS

Giovanna Bianchi, Richard Hodges <i>Introduction</i>	7
Richard Hodges <i>Towards a new Mediterranean narrative for Early Medieval Tuscany</i>	11
Pierluigi Pieruccini, Mauro Paolo Buonincontri, Davide Susini, Carmine Lubritto, Gaetano Di Pasquale <i>Changing landscapes in the Colline Metallifere (Southern Tuscany, Italy): early medieval palaeohydrology and land management along the Pecora river valley</i>	19
Luisa Dallai with contributions by Andrea Bardi, Arianna Briano, Mauro Paolo Buonincontri, Mirko Buono, Luisa Dallai, Gaetano Di Pasquale, Stefania Fineschi, Giulio Poggi, Elisabetta Ponta, Marta Rossi, Luisa Russo, Vanessa Volpi <i>Investigations at Carlappiano: new archaeological findings in anthropic and natural landscapes</i>	29
Lorenzo Marasco with contributions by Arianna Briano, Simon Greenslade, Sarah Leppard, Carmine Lubritto, Paola Ricci <i>Investigations at Vetricella: new archaeological findings in anthropic and natural landscapes</i>	57
Emanuele Vaccaro <i>Long-distance ceramic connections: Portus Scabris (Portiglioni-GR), coastal Tuscany and the Tyrrhenian Sea</i>	81
Arianna Briano, Cristina Fornacelli, Elisabetta Ponta, Luisa Russo <i>Pottery circulation and wares in the rural world: the Colline Metallifere and south-eastern Tuscany in the Early Medieval period</i> .	101
Paolo Tomei <i>The power of the gift. Early Medieval Lucca and its court</i>	123
Marco Benvenuti, Laura Chiarantini, Cristina Cicali, Alessandro Donati, Alessia Rovelli, Igor Villa, Vanessa Volpi <i>Metals and coinage in Medieval Tuscany: the Colline Metallifere</i>	135
Giovanna Bianchi, Simone M. Collavini <i>Public estates and economic strategies in Early Medieval Tuscany: towards a new interpretation</i>	147

ITALIAN ABSTRACTS

Giovanna Bianchi, Richard Hodges <i>Introduzione</i>	163
Richard Hodges <i>Verso una nuova narrativa mediterranea per l'alto Medioevo toscano</i>	165
Pierluigi Pieruccini, Mauro Paolo Buonincontri, Davide Susini, Carmine Lubritto, Gaetano Di Pasquale <i>Alterazioni di paesaggio nelle Colline Metallifere nell'alto Medioevo: paleoidrologia e pianificazione territoriale nella valle del fiume Pecora</i>	167
Luisa Dallai con contributi di Andrea Bardi, Arianna Briano, Mauro Paolo Buonincontri, Mirko Buono, Luisa Dallai, Gaetano Di Pasquale, Stefania Fineschi, Giulio Poggi, Elisabetta Ponta, Marta Rossi, Luisa Russo, Vanessa Volpi <i>Le ricerche a Carlappiano: nuove evidenze archeologiche tra paesaggi antropici e naturali</i>	169
Lorenzo Marasco con contributi di Arianna Briano, Simon Greenslade, Sarah Leppard, Carmine Lubritto, Paola Ricci <i>Le ricerche a Vetricella: nuove evidenze archeologiche tra paesaggi antropici e naturali</i>	183
Emanuele Vaccaro <i>Ceramica e commerci di lunga distanza: Portus Scabris (Portiglioni-GR), la Toscana costiera ed il Mar Tirreno</i>	197
Arianna Briano, Cristina Fornacelli, Elisabetta Ponta, Luisa Russo <i>Circolazione e produzioni ceramiche nelle campagne: le Colline Metallifere e l'area grossetana nell'alto Medioevo</i>	199
Paolo Tomei <i>Il potere del dono. Lucca e la sua corte nell'alto Medioevo</i>	209
Marco Benvenuti, Laura Chiarantini, Cristina Cicali, Alessandro Donati, Alessia Rovelli, Igor Villa, Vanessa Volpi <i>Metalli e monete nella Toscana medievale: le Colline Metallifere</i>	217
Giovanna Bianchi, Simone M. Collavini <i>Beni fiscali e strategie economiche nell'alto Medioevo toscano: verso una nuova lettura</i>	223

Pierluigi Pieruccini*, Mauro Paolo Buonincontri**/***,
Davide Susini*, Carmine Lubritto****, Gaetano Di Pasquale**

CHANGING LANDSCAPES IN THE COLLINE METALLIFERE (SOUTHERN TUSCANY, ITALY): EARLY MEDIEVAL PALAEOHYDROLOGY AND LAND MANAGEMENT ALONG THE PECORA RIVER VALLEY

1. INTRODUCTION

One of the most important aspects of geoarchaeological research is the modelling of the landscape evolution in the past (VAN ANDEL 1994; FRENCH 2003). Landscape changes are investigated coupling the paleoenvironmental signal coming from geological records such as sediments and soils, and the biological signal coming from the proxies found within the geological record itself. Finally, stratigraphy and geochronology provide the sequence of events which occurred both before, during and/or after the settlements.

Therefore, sedimentology, stratigraphy, geomorphology and palaeopedology are the main branches of earth science that provide sets of information about palaeoenvironmental settings and their evolution through time and space.

This kind of analysis area is a useful archaeological tool as well, providing the information needed for a more extensive geoarchaeological approach at the regional and site level. In fact, archaeological observations are usually limited to a site environmental approach with little regard to meso- and macro-scale landscape dynamics (FRENCH 2003; RAPP, HILL 2006; GOLDBERG, MACPHAIL 2006).

In this context, river valleys have strongly influenced the strategies of the populations for exploitation of the inland slopes and valley floors, the latter being a natural migration, commercial and communication paths or due to factors such as the availability of water, favourable topography, presence of “soft” soils and suitable areas for settlements and productive activities.

The Pecora river offers a unique chance to reconstruct the fluvial landscape before and during the Early Middle Ages in this Mediterranean area and to evaluate the influence of human impact on the environment of the river basin due to its proximity to the medieval archaeological site of Vetricella (MARASCO *et al.* this volume). The importance of this area is also marked by the presence of fiscal and public estates from the mid-9th c AD network of earlier settlements (HODGES, BIANCHI; COLLAVINI in this volume). A new environment

became suitable for productive large kernelled crops as well as fruit tree growing, serving the landlords’ will to achieve the highest production of any possible lands and to optimize surpluses from their estates (BUONINCONTRI *et al.* 2015, 2017).

Thus, the main objective of this work is the reconstruction of the changes which occurred to the alluvial environments and dynamics, including channel patterns, at the scale of the river basin. As explained further on in this work, the investigations carried out in the distal reach provide strong constraints for the evaluation of the changing dynamics in the mid- and upper reaches of the Pecora valley. Therefore, unravelling the changes in the landscape and river dynamics at such macro-scale, that is the whole valley system, provides information about the effects of and relationships between human behaviour (land use) and environment.

2. STUDY AREA AND ENVIRONMENTAL BACKGROUND

2.1 GEOLOGICAL AND GEOMORPHOLOGICAL BACKGROUND

The Pecora river basin extends between the towns of Follonica to the SW and Massa Marittima to the NE with a catchment of about 250 km² (fig. 1). The bedrock of the basin is mainly made of Cretaceous marine shales and limestones belonging to the Ligurian lithostructural Units (“Argille a Palombini Fm.”, ISPRA 2002) that rest uncomfortably on the massive Triassic limestones belonging to the Tuscan lithostructural Units (“Calcere Cavernoso Fm.”, ISPRA 2002). The unconformity is related to the eastward migration of compressional tectonic movements until the Early Miocene. At the end of the Miocene the area underwent a vertical uplift and the onset of continental conditions are testified in the area by the presence of thick alluvial fan conglomerates of the Messinian age (“Conglomerati di Monte Bamboli Fm”, ISPRA 2002). Marine transgression occurred during the Pliocene, although in the Pecora basin the associated sediments (“Argille Azzurre Fm.”, ISPRA 2002) are not present due to erosion.

The area is known also as part of “Colline Metallifere”, a mining district exploited in the last three thousand years (LATTANZI *et al.* 1994 and ref. therein) for ore deposits of pyrite, Fe, Cu-Pb-Zn, Ag, Sb, Hg, Sn and Au. Their occurrence is related to hydrothermal fluids circulating along extensional faults that affect the bedrock following the emplacement of Plio-Quaternary magmatic bodies (BENVENUTI *et al.* 2009).

The definitive emersion of the area is recorded during the Quaternary period with the deposition of a complex of continental Unconformity Bounded Stratigraphic Units including

* Dipartimento di Scienze Fisiche, della Terra e dell’Ambiente, DSFTA – Università di Siena, Italy (pierluigi.pieruccini@unisi.it; susini.davide@gmail.com).

** Dipartimento di Agraria – Università di Napoli “Federico II”, Italy (mauropaulo.buonincontri@unina.it; gaetano.dipasquale@unina.it).

*** Dipartimento di Scienze Storiche e dei Beni Culturali, DSSBC – Università di Siena, Italy (mauro.buonincontri@unisi.it).

**** Dipartimento di Scienze e Tecnologie Ambientali, Biologiche e Farmaceutiche, Università della Campania “Luigi Vanvitelli”, Italy (carmine.lubritto@unicampania.it).

alluvial fan, fluvial and Calcareous Tufaterraces and barrages (BENVENUTI *et al.* 2009). The present-day hilly landscape is mainly controlled by selective erosion and fluvial dynamics with formations of steeper slopes on the harder pre-Miocene and Miocene bedrock and wide, flat alluvial and Calcareous Tufa terraces along the valleys. Moreover, karst processes deeply influenced landscape evolution as indicated by the presence of wide coalescent doline depressions, karst springs (i.e. “Le Venelle” and “Aronne”, *fig. 1*) and related depositional environments. The evolution of the present-day drainage network is controlled by tectonic uplift coupled with Quaternary climatic changes through alternating phases of fluvial incision and aggradation, river captures and terrigenous and carbonate deposition (BENVENUTI *et al.* 2009). The historical and present-day slope and river dynamics are mainly controlled by land use and hydraulic regulations (*fig. 1*). In this paper we focus on the early medieval fluvial landscape evolution including continental carbonatic depositional environments. The latter is characterised by Calcareous Tufa (CT from now on) sediments characterized by physio-chemical and microbiological carbonate precipitation by fluvial and spring-fed fresh and cool water carbonates, generally restricted to localized karstic areas (FORD, PEDLEY 1996; PEDLEY 2009; CAPEZZUOLI *et al.* 2014). The typical landscape associated with the deposition of CT is formed by terraced flat palustrine environments bound by barrages and waterfalls. Such landscapes were widely present along the Pecora and the Le Venelle-Ferriere tributary (*fig. 1*). CT deposition is usually associated with the presence of carbonate-rich waters and humid and warm phases since CO₂, necessary for CaCO₃ dissolution and precipitation, is normally enriched under forest cover and the associated soils. However, human activity, namely deforestation, agricultural practices and drainage regulations, is the main factor that triggers their depositional decline in proto-historical and historical times (GOUDIE *et al.* 1993).

2.2 CLIMATE AND VEGETATION

According to the weather station of Follonica (4.34 m a.s.l., UTM 643775 E, 4753770 N, data source <http://www.sir.toscana.it/>), the area is characterised by a Mediterranean climate, with an average minimum temperature of 3.1°C during the coldest month (25 years observations) and an average annual precipitation of 592 mm (observed over 80 years). Arable crops, vineyards and olive groves are present in the flat valley floors and/or on the gentler slopes. The evergreen sclerophyllous forest, dominated by *Quercus ilex* L. with *Arbutus unedo* L., prevails on the steeper slopes of the Pecora river basin. Small stands of deciduous broadleaved species, such as *Q. cerris* L., *Q. pubescens* Willd., and *Fraxinus ornus* L. are sparse, whereas the deciduous oak forest, dominated by *Q. cerris* L., is located only on the cooler north-western slopes of the basin.

3. MATERIALS AND TOOLS

3.1 SEDIMENTOLOGY AND GEOMORPHOLOGY

The investigations have been mainly carried out in a retention basin excavated ca. 500 m to the NW of the Vetricella site (*fig. 1*) on the hydrographic left of the Pecora river. The




fig. 1 – Geological and geomorphological sketch of the studied area. Legend: 1) Localities cited in the text (1-Pian del Padule; 2-Padule del Moreta; 3-La Cascata; 4-Le Venelle Spring; 5-Aronne Spring); 2) Artificial levees; 3) Pecora river palaeochannels; 4) CT barrage; 5) Artificial channels; 6) Fluvial scarps; 7) Pecora and Le Venelle-Ferriere rivers and major tributaries; 8) Alluvia fan; 9) Lagoonal deposit reclaimed after 1950; 10) Beach deposit (Holocene); 11) CT (Holocene); 12) Unactive alluvial deposit (Late Pleistocene-Holocene); 13) Active alluvial deposit (Late Pleistocene-Holocene); 14) CT (Late Pleistocene); 15) Polygenic conglomerates (Messinian); 16) Macigno Fm (Upper Oligocene – Lower Miocene); 17) Argille a Palombini Fm (Upper Cretaceous – Paleocene); 18) Calcare Cavernoso Fm (Upper Triassic).

retention basin is c. 400 m long parallel and c. 100 m wide perpendicular to the Pecora river. The works started in 2015 and ended in 2016. During the excavations our research group was granted special access to these sections thanks to the kind collaboration of the “Consorzio di Bonifica 5 Toscana Coste” and the “Soprintendenza Archeologia, Belle Arti e Paesaggio per le Province di Siena, Grosseto e Arezzo”. The excavations allowed the observation of a sedimentary section, around 8 m thick, cutting the alluvial terrace on top of which is located the Vetricella site (Section E, *fig. 1*) and c. 3 m thick sequence perpendicular to the river flow direction (Sections




fig. 2 – The NW section studied within the retention basin (for location see fig. 1). The US numbering follows the description in the text whereas the asterisks indicates the radiocarbon dates on charcoal samples.

NW, figg. 1 and 2). The outcropping sequence was analysed in the field, drawing the sedimentological characteristics in terms of facies analysis following the method proposed by Miall (1996). Samples of sediments were taken for routine analysis whereas charcoals were sampled both for radiometric dating and anthracological analysis. Facies analysis allows the reconstruction of the depositional environment starting from the sedimentological characteristics (grain size, composition, shape of clasts, internal geometry and fabric) or lithofacies that indicate the flow dynamics; lithofacies are in turn grouped into associations or architectural elements that correspond to the characters of the internal depositional environments. Finally, the associations of architectural elements and the presence of bounding surfaces allow the definition of fluvial models or styles that correspond to the river dynamics and associated landscape.

To establish the relationships of the observed sedimentary characteristics within the valley, further investigations have been carried out along the thalweg. In the studied area the River Pecora is confined downflow within artificial levees and the river is hanging on the floodplain; this prevents the observation of the sediments. Upflow, the thalweg is downcut within the floodplain sediments allowing the description of further sedimentary sections. The analysis was also coupled with geomorphological investigations along the Pecora basin; these have been carried out in the field and in the laboratory by remote analysis of colour and black and white multitemporal aerial photos, spanning from 1954 to 2016. Moreover, the imagery was integrated with a highly detailed Digital Terrain Model derived by LiDAR and 10 m DTM. All the remote images and topographic models are available from the Regione Toscana Environmental Agency (<http://www.regione.toscana.it/web/geoblog/-/open-geodata>).

The geomorphological mapping was carried out following the guidelines of the Italian National Geological Survey for the 1:50,000 scale geomorphological maps of Italy (ISPRA-APAT, 1994). Punctual, linear, polygonal landforms and related deposits are classified according to the prevalent geomorphic process (i.e. superficial running water, gravity, karst, anthropic etc.). The field and remote maps have now been digitalized in the ArcGIS© environment and the features classified within a geodatabase structured following the adopted legend.

3.2 CHARCOAL

The abundance of charcoal within the sediments is the result of the incomplete combustion of plant vegetation (FORBES *et al.* 2006). Usually, it is difficult to establish the

source and the formation processes leading to the creation of the charcoal assemblage. The main source is generally forest fires naturally ignited or fired by humans in relation to anthropogenic activities (PYNE, GOLDAMMER 1997; MOORE 2000). The possible provenance of charcoals from nearby human-related contexts, such as archaeological sites or charcoal kilns, must also be taken into account when interpreting the charcoal record. Furthermore, in cases of forest fires, topographic conditions can influence the dispersion and deposition of charcoal during and after a fire (CLARK 1988; THINON 1992; SCOTT *et al.* 2000).

In the case of Pecora river, bedforms were characterised by the abundance of very fine to very coarse charcoal. In the Section NW, 23 sediment samples were collected at different levels, ranging from 500 to 2390 ml of volume. The samples were firstly air-dried and weighed, and successively wet-sieved through two sieves with 1 and 0.4 mm mesh-size. Charcoal concentration and taxonomical identification were preliminarily performed for charcoal remains larger than 1 mm. Smaller charcoal fragments were excluded since they are usually not suitable for taxonomical identification and cannot provide information on past forest composition (ROBIN *et al.* 2014). Charcoal concentration is expressed as specific anthracomass (SA) in milligrams of charcoal per kilogram of dried sediment (TALON 2010). Specific anthracomass is calculated per sample layer (SAL).

Taxonomical identification was carried out with an incident light microscope at a 100×, 200× and 500× magnification and supported with wood anatomy atlases (ABBATE EDLMANN, DE LUCA, LAZZERI 1994; SCHWEINGRUBER 1990; VERNET *et al.* 2001) and the reference collection at the Dipartimento di Agraria, Università di Napoli “Federico II”. Charcoal fragments were identified at the species or genus level for the most part, thanks to their good preservation, whilst for only a few poorly preserved fragments (bad preservation or vitrification) the identification was possible at a family level, or no identification was possible at all.

4. DATA

4.1 SEDIMENTOLOGY AND FACIES ANALYSIS

The retention basins allowed the observation of 4 sections and the presence of 4 well distinguishable Stratigraphic Units (US) separated by important sedimentary unconformities (fig. 3).

Here, we mainly describe and discuss the palaeochannel, c. 50 m wide and 3 m deep, observed along a section (NW

section) perpendicular to the Pecora flow direction excavated within a retention basin (*figg.* 1-2). The section is located in the distal reach of the Pecora river where the thalweg is regulated by artificial levees and the landscape opens into a wide flat floodplain. The current geomorphological setting is the result of the reclamation of the wide lagoonal and swampy system that characterised the area at least until the beginning of the 19thc, as is also indicated by the Maps of the Catasto Leopoldino available for the area (<http://www502.regione.toscana.it/castoreapp/>). Up valley the watercourse is not regulated and becomes suddenly incised within the floodplain. The incision increases up to 10 m in the upper reach where the CT characterises the landscape with wide flat terraces alternated with barrages.

4.1.1 US1

US1 is made of massive to laminated, bioturbated, horizontally bedded greyish clays (Fm, Fl; MIALI 1995), with rare sandy-silty lenses and blankets (Sm, Sh; MIALI 1995). Rare vegetal fragments (woods, leaves) and charcoals are present. The overall sedimentary characteristics are typical of a very low-energy lagoonal environment, perennially flooded as indicated by the lack of any pedogenetical evidence.

A clear erosional unconformity cuts US1 on top. The unconformity is undulated and corresponds to a channel system more than 8 m deep that shows a planar geometry parallel to the river flow. This unconformity is buried under US2 (*figg.* 2-3).

4.1.2 US2

US2 is separated by a minor unconformity and characterised by different facies associations (US2.1 and US2.2). US2.1 is made of trough and planar cross-bedded, or massive sands and silts (Sp, St, Sh, Sm) with very rare polygenic fine gravels and very thin muddy lenses (Fm). The gravels represent the lithologies of the Ligurian and Tuscan Units outcropping in the catchment area. The facies association, mainly made of downstream accretion and sandy bedforms with minor floodplain architectural elements, are typical of a deep sand-bed braided river.

A minor unconformity is in turn buried under US2.2 sediments. They are packed, fine to coarse-grained, subrounded to rounded trough cross-bedded or massive, poorly sorted gravels (Gt, Gh, Gm) with abundant sandy matrix. The gravels are polygenic and represent all the Ligurian and Tuscan Units present in the drainage basin. The architectural elements made of gravelly bedforms indicate massive sedimentation within a multi-storey channel typical of a shallow gravel-bed braided river (MIALI 1995). US2.2 constitutes the uppermost part of the alluvial terrace whose top surface is at a maximum elevation of about 5 m above the present-day thalweg and floodplain and gently sloping to the south, where the site of Vetricella is located. US2.2 is also weathered on top by a polycyclic, reddish, strongly leached argillic soil (ALFISOL, USDA 2015, *fig.* 4).

4.1.3 US3

On the N section a ca. 4 m deep channel cuts both US2 and US1. This channel is filled by US3 sediments and, in turn, is cut by a further weakly undulated and shallower channel filled by US 4 sediments (*figg.* 2-3).




fig. 3 – The US observed within the retention basin and described in the text. Bold lines are the main unconformities, dotted lines represent the minor unconformities.

US3 is made of loose planar cross-bedded, poorly sorted, rounded to subrounded fine to coarse-grained gravels (Gp), variable amount of sandy matrix. Minor planar or low angle cross-bedded sands (Sp) and lenses and blankets of massive to finely laminated silts and clays (Fm, Fl) are also present. The composition of the gravels also include rare clasts of CT and rare charcoal fragments. The facies association is typical of a gravel-sand wandering to meandering river (MIALI 1995) with a westward lateral accretion of gravelly and sandy bars and filling of abandoned channels.

4.1.4 US4

US4 is composed of loose, unsorted, trough crossbedded to massive, fine to medium-grained, rounded to subrounded gravels (Gt, Gh), locally matrix supported (Gsm). Minor trough cross-bedded or massive sands (St, Sh) and lenses and blankets of massive to finely laminated silts and clays (Fm, Fl) are also present. The gravels are made of very abundant clasts of CT, locally, filling entirely smaller channels. The bedforms are also characterised by the presence of very abundant fine to very coarse charcoals, both scattered within the sediments or concentrated along the base of the beds. The facies association, made of fine gravelly and sandy downstream accretion and mass flows bedforms with minor floodplain sediments are typical of a distal shallow gravel/sand-bed braided river environment. The same sediments filling smaller channels have also been observed a few tens of metres downflow alternated to laminated clays and silts, weakly weathered by palaeosols with root tracks (*fig.* 4) typical of shallow swampy or lagoonal environment.

Although the sedimentary characteristics of US4 are homogeneous, a further subdivision has been made on the base of the presence of minor unconformities: US4.1 is observed in the western part of the section that buries a shallow and almost flat unconformity that cuts both the US1 and the US3 sediments. US4.2, towards the east, fills a deeper channel cutting entirely into the US3 sediments. US4.2 also represents the thicker sub-unit suggesting that this is the most important depositional phase. Finally, both US4.1 and US4.2 are cut by a further very shallow and slightly undulated unconformity buried under US4.3 sediments.

4.2 CHRONOLOGY

The whole palaeochannel filled with US3 and US4 sediments is the last sedimentary event recorded in the succession investigated in the retention basin. For this reason,




fig. 4 – A) The unconformity affecting the basal US1 sediments filled by US2 sediments subdivided into two minor sub-units. US2.2 is weathered by a reddish palaeosol on top. B) Angular, unsorted, polygenic gravels of the US2.2 unit. C) The almost monogenic composition of the US4 gravels and sands made of clasts coming from the erosion of the CT. D) Root tracks and weakly developed palaeosols affecting the shallow swampy or lagoonal deposits entered by the US4 sediments.

it has been investigated also to assess its chronology and the sedimentary changes observed. This was made through radiocarbon dating of the abundant charcoals found mixed with the sediments. Preliminary results (fig. 2; tab. 1) show that the US3 sediments were left at least until 787-471 CAL age BC (Sample Fi3497) whereas the filling of the upper palaeochannels (US4) can be dated between 660-779 CAL age AD (US4.1, Sample Fi3554) and 1150-1290 AD (US4.3, Sample Fi3451).

4.3 SEDIMENT CHARCOAL ANALYSIS

Sediment charcoal analysis was preliminarily carried on 14.5 litres of sediment from 13 samples, with ca. 19.4 g of extracted charcoal. The samples were collected from US4, characterised by braided shallow channel (sub-units US4.3 and US4.2) and braided sheet channel (sub-units US4.1; Table 1). The sample 1.3 collected in US4.1 was found to be sterile.

To assess the taxonomical information, 191 charcoal fragments were preliminarily analysed allowing the identification of 17 taxa. The results of the taxonomical identification,

together with the SAL and the radiocarbon dating, are presented for each sample in the diagram (fig. 5). The main identified taxa useful to reconstruct the forest types, such as *Alnus*, *Erica*, *F. cf. oxycarpa*, *F. cf. ornus*, *Q. cf. cerris*, *Q. cf. ilex*, *Salix*, and *Ulmus* are presented in detail, whilst other identified broadleaved deciduous taxa (*Fraxinus*, Maloideae, Monocotyledon, *Prunus*, and *Quercus*) are grouped under the definition “Other broadleaves”. Calculations and percentages are based on the charcoal frequencies (number of identified charcoal remains per sample). To better highlight the vegetation changes in the record, key taxa have been grouped based on their ecological significance.

Among the identified taxa, deciduous broadleaved trees prevail on the total of the charcoal remains. *Ulmus* is the most attested (23%), followed by *Alnus* (7.3%), *F. cf. oxycarpa* (6.8%), *Q. cf. cerris* (5.8%), *Salix* and *F. cf. ornus* (3.7%). Among sclerophyllous evergreen taxa, *Erica* (4.7%) and *Q. cf. ilex* (2.1%) are attested. Other broadleaves constitute 25.7% of the total. Unidentifiable charcoals (because of bad preservation status or vitrification) constitute 13.1% of the total.

5. DISCUSSION

Paleoenvironmental reconstruction of fluvial landscapes along the Pecora river valley was made possible by an integrated sedimentological, geomorphological, stratigraphical, geochronological and charcoal analysis. In particular, the sections opened in the retention basin along the distal reach of the Pecora river allowed a detailed modelling of the sedimentary history along the valley and its relationships with the changing environment in the upper reach of the drainage basin.

The older depositional environment (US1) is characterised by clayey sediments deposited within a stable palustrine/lagoon environment with no evidence of palaeosols or arrival of coarse-grained sediments. Such an environment can be related to a marine high-stand during periods of warm and wet climatic conditions, possibly associated with the Last Interglacial (COLTORTI, PIERUCCINI 2006). US1 is abruptly cut by a major unconformity buried under US2 sediments. They are coarse-grained gravels and sands deposited in a high-energy fluvial environment typical of deep and shallow gravel and sand-bed braided rivers (MIALL 1995). These sediments form an alluvial fan whose top surface is gently sloping to the south. US2 indicates an environmental change with an increase in gravelly bedload and the formation of a braidplain. This environment is associated with a wide alluvial fan filling the earlier lagoon. This phase can be linked to the onset of cool and arid climatic conditions typical of the cooler stages of the Last Glaciation that enhanced the production of debris from denudated slopes driving the sedimentary aggradation of the valley floors. The c. 120 m sea level fall due to the cool climatic conditions shifted the coastline to the south. The climatic amelioration at the beginning of the Holocene, due to the abrupt bedload decrease together with the rapid sea-level rise, enhanced the capacity of the river system to downcut the valley floors (COLTORTI, PIERUCCINI 2006). Therefore, the previous alluvial fan forms an alluvial terrace where the Vetricella site is located, that in its apical part is c. 10 m high on the present-day thalweg.

A single channel, up to 3 m deep, cuts both the US1 and US2 and is filled by US3 sediments. These are gravels and sands with facies associations typical of a gravel-sand wandering to meandering river with the formation of point-bars and swamps along the valley floor following meander-cuts. The gravels, finer-grained than US2, are made of lithologies belonging to the bedrock of the basin and by scarce clasts deriving from the erosion of CT. This composition suggests that the CT systems, present in the mid- and upper reaches of the Pecora and Le Venelle-Ferriere rivers (*fig. 1*), were already in formation although not strongly eroded by fluvial dynamics. US3 sediments also commonly contain charcoals that allowed the dating of this sedimentary phase at least up to 787-471 BC Cal Age (Sample FI 3497) and possibly up to the Roman age. The presence of burnt vegetation also suggests the onset of land use practices related to fuel production or land clearing and vegetation opening.

US3 is cut by a further unconformity represented by a composite channel buried under c. 2.5 m of US4 sediments that can be subdivided into 3 main subunits bounded by minor unconformities (*figg. 2-3*). US4 sediments are characterised by unsorted fine-grained gravels, coarse-grained sands

and silts deposited within shallow and small channels with cut-and-fill geometries that indicate an abrupt change from a sinuous wandering river to a shallow braided river testifying to an increase of bedload and fast deposition in a less confined channel system. The most striking feature of US4 is the composition of the sediments that shows an abrupt change becoming predominantly made of unsorted clasts coming from the erosion of the CT. This is the very first evidence of the erosion of the up-valley CT environments (*fig. 1*), although the amount of sediments and the geometry of the sedimentary body suggest that only minor changes occurred upstream. These environments are typically made of wide flat swampy areas alternated with barrages or waterfalls. Both environments enhance the deposition of calcium carbonate due to the biological activity of algal and microbial floras and faunas within the swamps and to turbulence in the correspondence of the waterfalls (CAPEZZUOLI *et al.* 2014). The sedimentary characteristics and the chronology of US 4 indicates that the process of inactivation and erosion of the CT systems occurred in a very short time span that starts in the 8th c AD (US4.1) but is mostly concentrated in the 10th-11th c AD (US4.2), with sedimentation rates up to 0,7 cm/yr. The response of the fluvial system to the changes occurring in the upper reach of the catchment lasted until the 13th c AD (US4.3).

The presence of swampy areas along the Pecora river valley is also testified by the place names reported in the topographic maps such as “Padule Moreta” and “Pian del Padule” (“Padule” is a Tuscan term for swamp, *fig. 1*). The drainage of these wet areas occurred by means of drainage regulations still observable today. Moreover, the main barrage of the CT system (named “La Cascata” meaning Waterfall, *fig. 1*) is bypassed by a deep artificial cut that diverted the natural course of the river to the west, draining the whole wetland up-valley (*fig. 1*). All these artificial water regulations are responsible for the drainage of the CT environments and for their erosion due to the subsequent river down-cut that led to the transport and deposition of the amazing amount of CT clasts found in the NW section down valley. Furthermore, the NW section is located very close to the mouth of the Pecora river entering the early medieval shallow swampy or lagoonal environment as testified by the interlayering of these deposits with the US4 sediments (*fig. 4*).

US4 sediments also contain very abundant charcoals, from very coarse to fine, concentrated within the troughs and beds. Charcoals are within sedimentary record and their abundance coupled with the absence of human artefacts allows us to exclude an archaeological-related contamination (i.e. there are not derived from a nearby archaeological site, see chapter 3.2.). Moreover, the dispersion and deposition of charcoal during and after the fire (CLARK 1988; THINON 1992; SCOTT *et al.* 2000) is strictly dependent on run-off processes and the capability of the river to re-distribute sediments and charcoals down-valley. Such fast and widespread vegetation fires linked to the erosion of the CT environments and their drainage cannot be attributed to climate changes also because the climatic signals in the Mediterranean are not strong enough (LUTERBACHER *et al.* 2012) to be able to generate such a rapid response of the fluvial system for the time interval under consideration.

The preliminary radiocarbon dating, pointed to the overlapping alluvial phases, shows a coherent and linear




fig. 5 – Section NW charcoal analysis. From left to right: charcoal samples listed according to the depth; light grey bars: SAL (sample layer) values; black bars: percentage of most important taxa with indication of the AMS radiocarbon dates; dark grey bars: related ecological groups; stratigraphic units according to facies analysis.

chronological sequence. Assuming that the charcoal record is a result of forest fires, we can argue that the period between the end of the 8th c AD and the 13th c AD recorded the presence of fire activity within the Pecora river basin, with a peak of fire activity in the 9th-11th c AD period.

US4.1 is preserved on the right side of the palaeochannel and indicates a first phase of deposition occurring in a very shallow, flat and unconfined channel system dated to 660-779 AD Cal Age (Sample FI3554). This age postdates the beginning of the erosion of the CT systems and the onset of a braidplain in the distal reach of the Pecora river. However, the limited thickness of the sediments and the unconfined geometry of the depositional unit indicates only minor changes occurring in the upper catchment.

US4.1 taxonomical identifications document the presence of *Alnus* and, at a lesser extent, *F. cf. oxycarpa*, indicating that in this period mainly riparian and mixed floodplain vegetation is affected by fire, coupled with the onset of fluvial erosional events along the swampy areas within the CT environments. This suggests that the drainage and reclamation of these wet environments was also due to vegetation clearance along the floodplain, which could be evidence of woodland management carried out by 8th c AD settlements capable of seeking high-quality crop production in a period of involution of the medieval state (BUONINCONTRI *et al.* 2017). However, fire activity and sedimentary processes increase in the following centuries.

In fact, US4.1 is in turn cut by a minor unconformity that indicates the formation of a deeper channel whose filling, US4.2, shows the same facies model, with the predominance of cut-and-fill bedforms made again of CT fine-grained gravels. Nevertheless, the increasing thickness of the sedimentary succession and its deposition within a deeper and more confined channel indicates that this is the main depositional phase, possibly associated with the major drainage and environmental changes occurring upstream. Also, the frequency and volume increase of charcoal fragments lead to the older dating of this depositional stage to 880-980 AD Cal Age (Sample FI3452).

US4.2 anthracomass shows that fire activity included more established forest vegetation made of *Ulmus*, *F. cf. oxycarpa*, *Alnus*, *Populus* and *Salix*. This indicates that fire events affected a wider environment involving the riparian and mixed flood-plain forest occurring along the riverbed, wetlands, and distal alluvial plain. The presence of *Erica*, *Q. cf. ilex* and *F. cf. ornus* suggests that broadleaved evergreen forest of the foothill areas was also affected by fire activity.

Thus, the observed evidence argues in favour of the clearing and reclaiming of woodland in the whole Pecora basin suitable for the cultivation of large kernelled crops as well as fruit tree cultivation (BUONINCONTRI *et al.* 2015, 2017), probably due to the establishment of a new economic system (HODGES, BIANCHI; COLLAVINI *et al.* this volume). This hypothesis is also supported by the comparison of charcoal data with pollen analysis in the region, which show a noteworthy match of fire signals and forest clearance. At the Accesa lake the pollen sequence attests a decrease of wild arboreal pollen and an increase of cultivated tree pollen (olive and chestnut) from c. 850-950 AD (MAGNY *et al.* 2007).

US	Sub-Units	CT abundance	Facies model	¹⁴ C Age	Ch samples	Ch abundance	Fired vegetation
US4	US4.3		Distal sheetflood unconfined gravel/sand-bed braided river	1150-1290 AD	1.1	***	Riparian, proximal, distal floodplain and foothill
					2.1		
					3.1		
					1.2		
					2.3		
	US4.2	*****	Distal shallow gravel/sand-bed braided river	820-980 AD	2	****	Riparian, proximal and distal floodplain
					3		
					1.3		
					4		
					1		
US4.1		Distal sheetflood unconfined gravel/sand-bed braided river	660-779 AD	2.4	***	Riparian and proximal floodplain	
				1.4			
				7			
US 3		*	Gravel-sand wandering to meandering river	787-471 BC		*	

tab. 1 – Section NW sedimentary facies and related chronology and environments CT (Calcareous Tufa) and Ch (Charcoal) abundance: * rare; **scarce; ***common **** abundant *****very abundant.

A further minor unconformity is buried under US4.3 sediments that show the same sedimentary and compositional characteristics in the period of 1150-1290 AD Cal Age (Sample FI3451). However, the thickness of the sedimentary record is very limited and, as with US4.1, deposition occurred in an unconfined shallow channel, indicating a decreasing in sedimentary and associated environmental dynamics in the surrounding landscape.

In this unit a strong presence of *Q. cf. cerris* and, at a lesser extent, *Erica*, *Ulmus*, *F. cf. oxycarpa*, *Populus* and/or *Salix*, *Q. cf. ilex* and *F. cf. ornus* is recorded. Overall, the data suggest that in this phase fire events occurred intensely on the foothill areas, affecting the thermophilous deciduous forest dominated by *Q. cerris*.

The reclamation of the CT environments seems to come to an end in the 12th and 13th c AD, when exchange, productive specialization and semi-industrial activity became very complex and affected agriculture. In the same period, the farming system reached the highest levels and its expansion reached the foothills (DI PASQUALE *et al.* 2014; BUONINCONTRI *et al.* 2015, 2017).

This is the last sedimentary stage recorded within the palaeochannel, although we cannot exclude the erosion of younger sediments. However, the geomorphological investigations indicate that after this moment the river dynamics were not confined within channels until the definitive reclamation occurred after the 19th c AD.

6. CONCLUSIONS

The analysis of the NW section within the retention basin allowed a detailed reconstruction of the physical and biological environments of the fluvial landscape in the distal reach of the Pecora river, which represents an exceptional case of study for the Early Medieval times for this Mediterranean area. Sedimentological investigations revealed the erosion of the CT systems in the mid-to proximal reach of the river indicated by the occurrence of CT clasts. The erosion must be related to the disappearance of these environments that occurred in a short time span comprised between 8th and 13th c AD together with an abrupt change from sinuous-

meandering to braided river model. These changes are also characterised by the appearance within the sedimentary record of abundant charcoal remnants. The processes related to the CT swamps and systems of barrages/waterfalls inactivation must be attributed to fast drainage of the swampy areas and bypass of the waterfalls that work as barrages for the wet environments. Such processes cannot be attributed to natural events (i.e. climate changes) due to the very short time duration necessary for the erosion, transport and deposition downvalley of such an abundant sedimentary record and charcoals associated with vegetation fires. Moreover, for the time span under consideration there is no global or local climatic signal able to trigger such strong environmental changes. On the other hand, artificial run-off regulations for the improvement of wet environments drainage and bypassing of waterfalls can serve as the determinant factor for the recorded changes. Furthermore, these processes occurred at the same time with widespread vegetation fires that can be easily attributed to land clearance for agricultural purposes as suggested by the selective fire activities along the ecological vegetation groups typical of environments proximal to the valley floors such as riparian, floodplains and foothills. Within the medieval palaeochannel we recognised 3 main depositional events that embrace respectively the 8th c, 9th-11th c and 12th-13th c AD. The older event is characterised by a limited thickness of sediments and the minor amount of associated charcoals, suggesting as in this chronological interval that the drainage and vegetational changes in the upper catchment were at an embryonic stage. The most important event in terms of sedimentary response and charcoal amount is associated with the 9th-11th c AD when the CT environments become strongly eroded and mostly drained, as suggested also by the broader ecological groups affected by fires. Finally, the last evidence of this sedimentary history spans the 12th-13th c centuries AD, when the depositional processes decreased and the fires also affected the hilly landscape.

This analysis revealed that the fluvial dynamics and associated landscapes changed in the Early Medieval times, with major changes occurring at the Vetricella settlement providing information about site formation processes and landscape exploitation strategies.

BIBLIOGRAPHY

- ABBATE EDMANN M.L., DE LUCA L., LAZZERI S., 1994, *Atlante anatomico degli alberi e arbusti della macchia mediterranea*. Relazioni e monografie agrarie subtropicali e tropicali n.s. 114, Firenze.
- BENVENUTI M., COSTAGLIOLA P., TANELLI G., 2009, *Quaternary evolution of the Pecora River (southern Tuscany, Italy): paleohydrography and sediments provenance*, «Bollettino Della Società Geologica Italiana», 128, pp. 61-72.
- BENVENUTI *et al.* 2009 = BENVENUTI M.G., COSTAGLIOLA P., BENVENUTI M., BUCCIANI A., DI BENEDETTO F., PAOLIERI M., LATTANZI P., *Anthropogenic vs. natural sources of toxic elements in mining districts: The case of Val di Pecora (Tuscany)*, «Rendiconti Online Della Società Geologica Italiana», 6, pp. 49-50.
- BUONINCONTRI *et al.* 2017 = BUONINCONTRI M.P., PECCI A., DI PASQUALE G., RICCI P., LUBRITTO C., *Multiproxy approach to the study of Medieval food habits in Tuscany (central Italy)*, «Archaeological and Anthropological Sciences», 9, pp. 653-671.
- BUONINCONTRI M.P., SARACINO A., DI PASQUALE G., 2015, *The transition of chestnut (Castanea sativa Miller) from timber to fruit tree: Cultural and economic inferences in the Italian peninsula, «The Holocene»*, 25, pp. 1111-1123.
- BUTZER K.W., 1982, *Archaeology as human ecology: Method and theory for a contextual approach*, Cambridge.
- CAPEZZUOLI E., GANDIN A., PEDLEY M., 2014, *Decoding tufa and travertine (fresh water carbonates) in the sedimentary record: The state of the art*, «Sedimentology», 61, pp. 1-21.
- CLARK J.S., 1988, *Particle motion and the theory of charcoal analysis: source area, transport, deposition, and sampling*, «Quaternary Research», 30, pp. 67-80.
- COLTORTI M., PIERUCCINI P., 2006, *The Last Interglacial Pedocomplexes in the litho- and morpho-stratigraphical framework of the Central-Northern Apennines (Central Italy)*, «Quaternary International», 156-157, pp. 118-132.
- DI PASQUALE *et al.* 2014 = DI PASQUALE G., BUONINCONTRI M.P., ALLEVATO E., SARACINO A., *Human-derived landscape changes on the northern Etruria coast (western Italy) between Roman times and the late Middle Ages, «The Holocene»*, 24, pp. 1491-1502.
- FEDELE F.G., 1976, *Sediments as paleo-land segments: the excavation side of study*, in D.A. DAVIDSON, M.L. SHACKLEY, *Geoarchaeology*, London, pp. 23-48.
- FORBES M.S., RAISON R.J., SKJEMSTAD J.O., 2006, *Formation, transformation and transport of black carbon (charcoal) in terrestrial and aquatic ecosystems*, «Science of the Total Environment», 370, pp. 190-206.
- FORD T.D., PEDLEY H.M., 1996, *A review of tufa and travertine deposits of the world*, «Earth-Science Reviews», 41, pp. 117-175.
- FRENCH C., 2003, *Geoarchaeology in action*, London.
- GLADFELTER B.G., 1977, *Geoarchaeology: the geomorphologist and archaeology*, «American Antiquity», 42 (4), pp. 519-538.
- GOLDBERG P., MACPHAIL R.I., 2006, *Practical and Theoretical Geoarchaeology*, Oxford.
- GOUDIE A.S., VILES H.A., PENTECOST A., 1993, *The late-Holocene tufa decline in Europe*, «The Holocene», 3 (2), pp. 181-186.
- ISPRA (Istituto Superiore per la Protezione dell'Ambiente), 2002, *Carta Geologica d'Italia in scala 1:50.000 e Note Illustrative, Foglio 306 Massa Marittima*. http://www.isprambiente.gov.it/Media/carg/306_MASSA_MARITTIMA/Foglio.html
- ISPRA-APAT, 1994, *Carta Geomorfologica d'Italia 1:50,000, Guida al Rilevamento, Quaderno 4*, Roma.
- LATTANZI *et al.* 1994 = LATTANZI P.F., BENVENUTI M., COSTAGLIOLA P., TANELLI G., *An overview on recent research on the metallogeny of Tuscany, with special reference to the Apuane Alps*, «Memorie della Società Geologica Italiana», 48, pp. 613-625.
- LUTERBACHER *et al.* 2012 = LUTERBACHER J., GARCÍA-HERRERA R., AKÇER-ÖN S., ALLANE R., ALVAREZ-CASTROF M.-C., BENITO G., BOOTH J., BÜNTGEN U., ÇAĞATAY N., COLOMBAROLI D., DAVIS B., ESPER J., FELIS T., FLEITMANN D., FRANK D., GALLEGO D., GARCIA-BUSTAMANTE E., GLASER R., GONZÁLEZ-ROUCO F.J., GOOSSE H., KIEFER T., MACKLIN M.G., MANNING S.W., MONTAGNA P., NEWMAN L., POWER M.J., RATH V., RIBERA P., RIEMANN D., ROBERTS N., SICRE M.-A., SILENZI S., TINNER W., TZEDAKIS P.C., VALERO-GARCÉS B., VAN DER SCHRIERA G., VANNIÈRE B., VOGT S., WANNER H., WERNER J.P., WILLETT G., WILLIAMS M.H., XOPLAKI E., ZEREFOS C.S., ZORITA E., *A review of 2000 years of Paleoclimatic evidence in the Mediterranean*, in P. LIONELLO, *The Climate of the Mediterranean Region: From the Past to the Future*, New York, pp. 87-185.
- MAGNY *et al.* 2007 = MAGNY M., DE BEAULIEU J.-L., DRESCHER-SCHNEIDER R., VANNIÈRE B., WALTER-SIMONNET A.-V., MIRAS Y., MIRET L., BOSSUET G., PEYRON O., BRUGIAPAGLIA E., LEROUX A., *Holocene climate changes in the central Mediterranean as recorded by lake-level fluctuations at Lake Accesa (Tuscany, Italy)*, «Quaternary Science Reviews», 26 (13-14), pp. 1736-1758.
- MIALL A.D., 1996, *The Geology of Fluvial Deposits*, Berlin.
- MOORE J., 2000, *Forest fire and human interaction in the early Holocene woodlands of Britain*, «Palaeogeography, Palaeoclimatology, Palaeoecology», 164, pp. 125-137.
- PEDLEY M., 2009, *Tufas and travertines of the Mediterranean region: a testing ground for freshwater carbonate concepts and developments*, «Sedimentology», 56, pp. 221-246.
- PYNE S.J., GOLDAMMER J.G., 1997, *The culture of fire: an introduction to anthropogenic fire history*, in J.S. CLARK, H. CACHIER, J.G. GOLDAMMER, B. STOCKS, *Sediment Records of Biomass Burning and Global Change*, Proceedings of the NATO Advanced Study Institute "Biomass burning and global change", Praia de Alvor, Portugal, (October 1994), Berlin-Heidelberg, pp. 71-114.
- RAPP G., HILL C.L., 2006, *Geoarchaeology. The Earth Science approach to archaeological interpretation*, New Haven.
- ROBIN *et al.* 2014 = ROBIN V., BORK H.-R., NADEAU M.-J., NELLE O., *Fire and forest history of central European low mountain forest sites based on soil charcoal analysis: the case of the eastern Harz, «The Holocene»*, 24, pp. 35-47.
- SCHWEINGRUBER F.H., 1990, *European Wood Anatomy*, Bern.
- SCOTT *et al.* 2000 = SCOTT A.C., CRIPPS J.A., COLLINSON M.E., NICHOLS G.J., *The taphonomy of charcoal following a recent heath land fire and some implications for the interpretation of fossil charcoal deposits*, «Palaeogeography, Palaeoclimatology, Palaeoecology», 164, pp. 1-31.
- TALON B., 2010, *Reconstruction of Holocene high-altitude vegetation cover in the French southern Alps: evidence from soil charcoal, «The Holocene»*, 20, pp. 35-44.
- THINON M., 1992, *L'analyse pédoanthracologique e Aspect méthodologique et applications*, Thèse, Faculté sciences et techniques de St. Jérôme.
- USDA, 2015, *Keys to Soil Taxonomy*, U.S. Department of Agriculture.
- VAN ANDEL T.H., 1994, *New views on an old planet: a history of global change*, Cambridge.
- VERNET *et al.* 2001 = VERNET J.L., OGÉREAU P., FIGUEIRAL I., MACHADO YANES C., UZQUIANO P., *Guide d'identification des charbons de bois préhistoriques et récents*, Paris.

Abstract

Sedimentary filling within fluvial valleys are excellent archives for landscape changes along the river basins since the valleys work as traps for sediments coming from the slopes and later re-worked, distributed and deposited by fluvial processes. In terms of physical and associated biological landscapes, fluvial sediments reveal if and how the riverine dynamics changed, providing information about different fluvial styles. At La Vetricella we had the chance to observe thick sedimentary records in the distal reach of the Pecora river basin. At c. 1 km to the NW from the site, engineering works for the excavation of a retention basin on the left bank of the river revealed the presence of a wide palaeochannel filled with up to 3 m of sediments. The combined sedimentological, geomorphological, anthracological and geochronological analyses revealed an abrupt change of the river dynamics and associated landscape that mainly occurred between the 8th and 13th c., testifying to strong human impact on the surface hydrology and vegetation in the mid- and upper reach of the Pecora river basin.

ITALIAN ABSTRACTS

Pierluigi Pieruccini, Mauro Paolo Buonincontri,
Davide Susini, Carmine Lubritto,
Gaetano Di Pasquale

ALTERAZIONI DI PAESAGGIO
NELLE COLLINE METALLIFERE
NELL'ALTO MEDIOEVO:
PALEOIDROLOGIA
E PIANIFICAZIONE TERRITORIALE
NELLA VALLE DEL FIUME PECORA

I riempimenti sedimentari delle valli fluviali costituiscono degli eccellenti archivi per stabilire variazioni le ambientali e quindi di paesaggio occorse all'interno dei bacini idrografici. Infatti le valli funzionano come trappole sedimentarie per i depositi provenienti dall'erosione dei versanti e dei suoli che successivamente vengono trasportati e ridistribuiti all'interno del sistema vallivo. L'analisi delle facies sedimentarie e delle successioni stratigrafiche consente di stabilire se gli stili fluviali e, di conseguenza, l'ambiente a scala di bacino ha subito nel tempo variazioni legate a dinamiche superficiali sia naturali che antropiche. Inoltre, i sedimenti all'interno delle valli possono contenere altri tipi di *proxies*, anche di origine biologica, che possono fornire dati e informazioni circa l'evoluzione dell'ambiente e del paesaggio biologico alla scala del bacino.

Circa 1 km a NO del sito di La Vetricella, a partire dal 2015, i lavori di realizzazione di una cassa di espansione in sinistra idrografica del Fiume Pecora hanno consentito l'osservazione e l'analisi della sequenza sedimentaria fluviale relativa all'apporto fluviale e, di conseguenza, le relazioni con l'insediamento della Vetricella in termini di ambiente fisico e biologico. Lo spessore totale di sedimenti osservati all'interno dell'opera idraulica è di circa 8 m nelle sezioni parallele al corso attuale del Fiume Pecora e di circa 3 m nelle sezioni perpendicolari al corso stesso. L'analisi geomorfologica e sedimentologica ha consentito di definire la successione stratigrafica e di eventi che ha condotto alla formazione della superficie su cui si sviluppa l'insediamento della Vetricella. Questa superficie costituisce la sommità di una conoide alluvionale re-incisa dal corso del Fiume Pecora, la cui parte apicale si trova sospesa di circa 5 m sul fondovalle diminuendo progressivamente di quota verso sud. I sedimenti osservati indicano la presenza di una sequenza argillosa di origine lagunare o palustre antica a cui si sovrappone in discontinuità erosiva una sequenza tipica di conoide alluvionale prevalentemente ghiaiosa. L'assenza di materiali databili non ne consente una precisa attribuzione cronologica ma è probabile che si tratti di una sequenza legata a condizioni climatiche caldo umide e di alto stazionamento marino (Ultimo Interglaciale) a cui si sovrappone il *record* grossolano legato alle fasi fredde ed aride e di basso stazionamento marino dell'Ultima Glaciazione.

Di maggiore interesse, per la cronologia relativa al Progetto nEU-Med, si è rivelata l'analisi delle sezioni perpendicolari al

corso d'acqua e che mostravano il riempimento sedimentario del fondovalle attuale. Infatti qui gli scavi hanno mostrato la presenza di un paleoalveo ampio circa 50 m e profondo circa 3 m, al cui interno si riconoscono due distinte facies sedimentarie che indicano in origine la presenza di un corso d'acqua molto sinuoso o meandriforme a cui succedono in discontinuità facies associabili ad un corso d'acqua a canali intrecciati. L'aspetto più caratteristico delle due facies è costituito dalla composizione dei sedimenti che nel paleoalveo più antico contengono scarsi clasti provenienti dall'erosione di Tufi Calcarei mentre in quello più recente tali clasti diventano predominanti. Complessi sedimentari costituiti da Tufi Calcarei sono diffusamente presenti nel tratto a monte sia del Fiume Pecora sia del tributario di sinistra Le Venelle-Le Ferriere. Gli ambienti deposizionali tipici dei Tufi Calcarei sono costituiti da alternanze di paludi e cascate la cui morfologia è chiaramente distinguibile ancora oggi e la cui presenza fino a tempi recenti è anche suggerita da alcuni toponimi. La cronologia delle variazioni occorse nella deposizione del paleoalveo è stata resa possibile dalla datazione al radiocarbonio dei rari frammenti di carbone presenti all'interno del primo riempimento del paleoalveo e dell'eccezionale quantità di carboni associati ai sedimenti provenienti dall'erosione dei Tufi Calcarei che costituiscono il riempimento più recente. I carboni, provenienti da incendi della vegetazione, inoltre consentono di definire il tipo di vegetazione interessata dagli incendi e la variazione nel tempo delle porzioni di paesaggio interessate. Nel complesso l'analisi integrata ci ha consentito di stabilire che il Fiume Pecora aveva un corso molto sinuoso e con scarso apporto di sedimenti provenienti dall'erosione dei Tufi Calcarei almeno fino al VI secolo AC mentre il paleoalveo più recente ha restituito una cronologia compresa tra l'VIII e il XIII secolo DC. All'interno di questo paleoalveo infine si possono riconoscere tre diverse fasi deposizionali attribuibili rispettivamente al VIII, IX-XI e XII-XIII secolo. La fase deposizionale più antica (VIII secolo) è caratterizzata da un sottile record sedimentologico a cui sono comunemente associati carboni mentre la seconda fase (IX-XI secolo) è più spessa e con resti di carboni molto abbondanti. Infine la terza fase (XII-XIII) mostra una diminuzione in termini di sedimenti e di carboni associati. Il drenaggio e la successiva erosione dei complessi di Tufi Calcarei sono quindi associabili a interventi antropici nel tratto a monte della Valle del Fiume Pecora i cui effetti deposizionali si osservano nel tratto a valle, in prossimità della pianura costiera. La variazione di sedimentazione indica come tali processi abbiano avuto gli effetti maggiori nell'intervallo IX-XI secolo impattando sia sui sistemi idrici e quindi sul paesaggio fisico sia sul paesaggio vegetale attraverso un graduale ampliamento delle aree interessate da incendi di vegetazione. Infatti nella fase più antica i resti paleoantropologici ci rivelano che la vegetazione interessata da incendi era quella palustre e ripariale mentre nelle fasi successive, la più importante, gli incendi si estendevano a tutta la pianura alluvionale andando ad interessare i versanti nell'ultima e più recente fase.

The nEU-Med project is part of the Horizon 2020 programme, in the ERC Advanced project category. It began in October 2015 and will be concluded in October 2020. The University of Siena is the host institution of the project.

The project is focussed upon two Tuscan riverine corridors leading from the Gulf of Follonica in the Tyrrhenian Sea to the Colline Metallifere. It aims to document and analyze the form and timeframe of economic growth in this part of the Mediterranean, which took place between the 7th and the 12thc. Central to this is an understanding of the processes of change in human settlements, in the natural and farming landscapes in relation to the exploitation of resources, and in the implementation of differing political strategies.

This volume brings together the research presented at the first nEU-Med workshop, held in Siena on 11-12 April, 2017. The aim of the workshop was to draw up an initial survey of research and related work on the project, one and a half years after its inception. The project is composed of several research units. Each unit covers an aspect of the interdisciplinary research underpinning the nEU-Med project, each with their own methodology. For this first volume of results, it was decided not to give an account of all the work carried out within all the units, but to select those lines of investigation which, at the end of the first year and a half, have made it possible to articulate and develop an interdisciplinary research strategy.

edited by
Giovanna Bianchi, Richard Hodges

Origins of a new economic union (7th-12th centuries)
Preliminary results of the nEU-Med project: October 2015-March 2017

€ 39,00

ISSN 2035-5319

ISBN 978-88-7814-879-6

e-ISBN 978-88-7814-880-2

BAM-25

