Geodynamics of the central-western Mediterranean region: plausible and non-plausible driving forces

Journal Pre-proof



Geodynamics of the central-western Mediterranean region: Plausible and non-plausible driving forces

E. Mantovani, M. Viti, D. Babbucci, C. Tamburelli, N. Cenni

PII: S0264-8172(19)30570-7

DOI: https://doi.org/10.1016/j.marpetgeo.2019.104121

Reference: JMPG 104121

To appear in: *Marine and Petroleum Geology*

Received Date: 10 July 2019

Revised Date: 30 October 2019

Accepted Date: 31 October 2019

Please cite this article as: Mantovani, E., Viti, M., Babbucci, D., Tamburelli, C., Cenni, N., Geodynamics of the central-western Mediterranean region: Plausible and non-plausible driving forces, *Marine and Petroleum Geology* (2019), doi: https://doi.org/10.1016/j.marpetgeo.2019.104121.

Geodynamics of the central-western Mediterranean region: plausible and non-plausible driving forces

Mantovani E.[1*], Viti M.[1], Babbucci D.[1], Tamburelli C.[1], Cenni N.[2]

[1]*Dipartimento di Scienze Fisiche, della Terra e dell'Ambiente, Università di Siena, Italy*

[2]*Dipartimento di GeoScienze, Università di Padova, Italy*

[*] *Corresponding author (enzo.mantovani@unisi.it)*

Corresponding author*:*

Enzo Mantovani

Dipartimento di Scienze Fisiche, della Terra e dell'Ambiente., Università di Siena

Via Laterina, 8 - 53100 Siena (I)

e-mail: *enzo.mantovani@unisi.it*

Abstract. The observed deformation pattern in the central-western Mediterranean area, in particular the development of the Northern, Central and Southern Tyrrhenian basins in three well distinct phases, can hardly be explained as an effect of the gravitational sinking of subducted lithosphere, an hypothesis often advanced in literature. A more plausible and coherent explanation of the spatio-temporal distribution of major tectonic events in the study area can instead be achieved by supposing that tectonic activity has mainly been driven by the convergence of Africa and Eurasia and the roughly westward displacement of the Anatolian-Aegean-Pelagonian belt. The development of Arc-Trench-Back Arc systems is interpreted as an effect of extrusion processes, that in some constricted contexts have represented the most convenient shortening process for accommodating plate convergence.

Keywords: Mediterranean, geodynamics, slab-pull, extrusion

1. Introduction

In the last 30-35 My the Mediterranean area has undergone a drastic change (Fig. 1), as suggested by several authors (e.g., Dercourt et al., 1986; Meulenkamp and Sissingh, 2003; Mantovani, 2005, Mantovani et al., 2006, 2009; Viti et al., 2009 and references therein). The pre-existing Alpine orogenic belt has undergone considerable distortions and long migrations of even several hundreds of Km. Some zones, mainly located in the internal side of the migrating arcs, have undergone crustal extension and consequent thinning, with the formation of large basins (Balearic, Tyrrhenian, Aegean and Pannonian). The oceanic and thinned continental zones which surrounded the African/Adriatic promontory have been almost completely consumed by subduction. Since extensional deformations are generally considered as scarcely compatible with the compressional context induced by plate convergence, some authors have tentatively advanced the hypothesis that other types of forces acted in the Mediterranean region. In particular, the most often cited interpretation suggests that the migration of arcs and the related back-arc extension have been driven by the gravitational sinking of subducted lithosphere, as originally proposed by Malinverno and Ryan (1986) and Royden (1993). In this work, we argue that the main implications of the above interpretation can hardly be reconciled with the observed features and that a more plausibile and coherent explanation of the evolutionary history can be achieved by supposing that Arc-Trench-Back Arc systems (ATBA) have developed in the framework of extrusion processes, driven by the convergence of the confining plates. With respect to previous attempts (e.g., Mantovani et al., 2009, 2014; Viti et al., 2009), this work reports new evidence and arguments about the compatibility of the interpretations here discussed with the observed deformations.

2. Balearic and Tyrrhenian basins interpreted as effects of the slab-pull genetic mechanism: main problems

Balearic basin

In the Oligocene, the western Alpine belt, tentatively reconstructed by the Al-Ka-Pe-Ca system, and the Corsica-Sardinia block (Fig.1a) detached from the European-Iberian foreland and underwent a fairly long eastward migration and strong bowing (Fig.2), while crustal extension occurred in the wake of the migrating arc, forming the Balearic basin (e.g., Maillard and Mauffret, 1999; Gaspar-Escribano et al., 2004; Gattacceca et al., 2007; Lustrino et al., 2009; Etheve et al., 2016)

Several authors (e.g., Royden and Burchfiel, 1989; Schmid et al., 1996, 2004; Stampfli et al., 1998; Finetti et al., 2005a; Handy et al., 2015) suggest that the western Alpine belt built up as an effect of the subduction of the Valais and Piemont-Ligurian oceanic domains and the European thinned continental margin, which generated a southward verging slab. In the Oligocene, such process underwent slowdown/cessation, when continental crust entered the trench zone. Then, along the Alpine sector lying aside the Iberian domain, another opposite verging (North to NW ward) subduction process started, involving the consumption of the Tethyan oceanic domain under the migrating Al-Ka-Pe-Ca belt (Gaspar-Escribano et al., 2004; Lustrino et al., 2009; Schettino and Turco, 2006; Handy et al., 2015; Etheve et al., 2016). This implies that when the ATBA system began to migrate, in the Oligocene, the northward dipping slab did not exist or was scarcely developed, being thus unable to induce a sufficient slab-pull force. To this regard, it should be considered that laboratory and numerical modelings suggest that a minimum slab length (150-300 km) is required in order to initiate back-arc extension driven by slab-pull (e.g., Hassani et al., 1997; Faccenna et al., 1999; Schellart, 2005).

Other authors instead, suggest that prior to the formation of the Liguro-Provencal basin (30-25 My) there was a well developed northward dipping slab at least 150 km long (Jolivet and Faccenna, 2000, Carminati et al., 2012; van Hinsbergen et al., 2014, Faccenna et al., 2014a) beneath the Alpine belt. However, no clear information is provided about the correspondence between the dimension and age of the present Alpine-Apennine belt and the orogenic material that the presumed subduction process would have accumulated in the trench zone.

The fact that during the formation of the Balearic basin the Northern Al-Ka-Pe-Ca arc underwent an evident sinistral NE-SW shift with respect to the Western Alps (Fig.2), leading to the formation of the Ligurian basin (e.g., Makris et al., 1999; Mosca et al., 2010), suggests that such arc was stressed by belt-parallel push. This kind of stress field is also suggested by the kinematics of the strike-slip faults and lateral escape of blocks in the Sardinia-Corsica microplate (e.g., Carmignani et al., 1994, 2004; Oggiano et al., 2009 and references therein). The Oligocene-Aquitanian age of this tectonic regime has been mainly constrained by the analysis of the infilling succession of transtensive basins. The above evidence is not compatible with the roughly eastward orientation of the trench suction force that would have been induced by a slab-pull mechanism.

Another major feature of the Balearic ATBA system that cannot easily be explained as an effect of slab-pull forces is the strong bowing that the Arc underwent during its migration (Fig. 2b). Numerical modelings of a slab-pull mechanism (e.g., Schellart et al., 2007) could justify a curvature of the migrating arc, but the resulting effect is much smaller than the one that really occurred in the western Mediterranean area (Mantovani et al., 2001a; 2002; Viti et al., 2009 and references therein**).** Moreover, the horizontal bowing predicted by modellings would need more than 16 My to develop (e.g., Schellart et al., 2007). Other laboratory experiments suggest instead a linear shape of the retreating subduction boundary, without significant arc bowing (Becker et al., 1999).

It is not demonstrated that the trench-suction force induced by gravitational sinking of a slab can break the upper plate. The results of some laboratory and numerical experiments (e.g., Shemenda, 1993; Hassani et al., 1997; Capitanio et al., 2010) indicate that slab-pull cannot produce

104 significant deformation in the upper plate unless it was not previously weakened by other tectonic
105 or magmatic processes. Even in this last case, the trench suction force would be unable to produce
106 extension in the overriding plate until the slab has reached a length of about 150-300 km.
107
108 *Northern Tyrrhenian basin (9-6 My)*
109 From about 9 to 6 My, the sector of the Alpine-Apennine belt that lay aside the Corsica-
110 Sardinia block (Fig. 3a) underwent roughly E-W crustal extension and consequent thinning (e.g.,
111 Mauffret et al., 1999; Sartori et al., 2001; Carmignani et al., 2004; Finetti et al., 2005a; Peccerillo,
112 2005; Sartori, 2005; Moeller et al., 2013; Cornamusini et al., 2014). This process formed the
113 Northern Tyrrhenian basin (Fig.3b), a roughly triangular zone confined to the South by the Selli
114 fault, which divided the extending Tyrrhenian area from an orogenic Alpine-Apennine body
115 (Sartori et al., 2001, 2004).
116
117 Explaining the main features of such extensional event as an effect of slab-pull forces (e.g.,
118 Malinverno and Ryan, 1986; Royden, 1993) involves serious difficulties.
119 Most authors suggest that the stop of the northern Al-Ka-Pe-Ca Arc and the Corsica-Sardinia
120 block occurred when the Adriatic continental domain entered the trench zone, around 15 My
121 (Speranza et al., 2002; Finetti et al., 2005a; Gattacceca et al., 2007; Molli 2008, Malusà et al.,
122 2016). In that context, any further development of slab roll-back would have encountered very
123 strong resistance from buoyancy forces (e.g., Brancolini et al., 2019). In fact, in the subsequent
124 period (from about 15 to 9 My) no possible effects of trench retreat (such as crustal extension) has
125 developed in the zone overlying the slab (Faccenna et al., 2001, 2014a; Finetti et al., 2005a). Thus,
126 the hypothesis that the subducted Adriatic margin has undergone gravitational sinking from about 9
127 to 6 My, forming the Northern Tyrrhenian basin, suggests some questions:
128 - Why slab-roll back did not occur for about 6 My and why it would have resumed at 9 My ?

- Why from 9 to 6 My gravitational sinking would have only affected the northern sector of a long slab lying beneath the whole Apennine belt ? in spite that the slab sector lying under the Central-Southern Apennines and Calabria was certainly more developed.

- Why was there no accretionary activity in the Northern Apennine belt during the formation of the Northern Tyrrhenian basin ? If crustal extension is supposed to be an effect of slab-pull forces, one could expect to observe as well the other major effect of that driving mechanism, i.e. the development of accretionary activity along the front of the migrating Arc, as indeed occurred in the previous period, during the formation of the Balearic basin.

- Why was the zone affected by crustal extension confined by the Selli fault, which is obliquely oriented with respect to the eastward orientation of slab roll-back (and of crustal extension) ?

- Why in the Messinian (6-5 My) gravitational sinking of the slab would have ceased beneath the Northern Apennines and started beneath the Southern Apennines ?

The attempts at interpreting the evolution of the Northern Tyrrhenian-Apennine system as an effect of slab-pull forces are encouraged by some tomographic studies (Lucente et al., 1999; Piromallo and Morelli, 2003; Spakman and Wortel, 2004) which suggest the presence of a well developed lithospheric body (hundreds of Km long) beneath that belt. However, these interpretations cannot easily be reconciled with the results of other tomographic investigations (Scafidi et al., 2009; Scafidi and Solarino, 2012) and CROP seismic soundings (e.g., Finetti et al., 2001, 2005a), which do not evidence any well developed lithospheric slab beneath the Apennine-Tyrrhenian system. The occurrence of some subcrustal earthquakes beneath the Northern Apennines cannot be used to confirm the results of tomography, since focal depths are mainly lower that 70 km (e.g., Chiarabba et al., 2005) and the magnitudes are very low (ISIDe Working Group, 2016). It is worth noting that the distribution of subcrustal earthquakes is compatible with the geometry of the crustal slivers suggested by CROP sections (Finetti et al., 2005a).

Some authors have tried to identify possible explanations for episodic effects of gravity on subducted lithosphere. For instance, Faccenna et al. (2001, 2014a) recognize the problem of

episodic slab retreat, stating that it could be explained by the interaction of the subducted lithosphere with the 670 km-depth mantle discontinuity. The inability of the slab to penetrate across such discontinuity would cause a sudden decrease of the subduction rate and trench retreat. Only after the slab has folded over the mantle discontinuity, subduction and trench migration may resume again, leading to a new phase of back-arc extension (Faccenna et al., 2004, 2007). However, it must be considered that the above results have been obtained by experiments where the upper plate is not present at all, a quite irrealistic condition. Other experiments have shown that the presence of a sufficiently strong upper plate can prevent slab roll-back (see figure 11 by Capitanio et al., 2010 and Shemenda, 1993; Hassani et al., 1997).

Central Tyrrhenian basin (about 5-2 My)

In the late Miocene-Early Pliocene crustal extension ceased in the Northern Tyrrhenian and began to develop in the Alpine-Apennine sector located south of the Selli fault, generating oceanic crust in the central Tyrrhenian area (Magnaghi and Vavilov basins, Fig.4, e.g., Finetti and Del Ben, 1986, 2005a; Mascle and Rehault, 1990; Sartori, 1990, 2005; Sartori et al., 2004; Guillaume et al., 2010).

Such extension was accompanied by other coeval major tectonic events in the central Mediterranean region, as listed in the following.

- Thrusting and folding underwent a significant acceleration in the Apennine belt (e.g., Patacca et al., 1990; Coward et al., 1999; Catalano et al., 2004; Cerrina Feroni et al., 2004; Parotto and Praturlon, 2004; Patacca and Scandone, 2007; Ghielmi et al., 2013).
- In the Northern Apennines, the foredeep geometry underwent a major change during the Late Pliocene, passing from a continuous (cylindrical) to highly fragmented (non-cylindrical) pattern (Amadori et al., 2019).

- Seismic soundings (Finetti et al. 2001, Finetti et al., 2005b) indicate that since the latest Miocene the Adriatic lithosphere has undergone strong shortening, accommodated by major thrust faults cutting the entire crust and that significant shortening (several tens of Km) also affected the migrating arc.

- Since the late Messinian, a major transtensional fault system (Sicily Channel, Medina and Victor Hensen, Fig. 4) developed in the Pelagian and Ionian zones (e.g., Finetti and Del Ben, 1986, 2005b; Boccaletti et al., 1987; Cello, 1987; Reuther et al., 1993; Hieke et al., 2003, 2006 and references therein).

- During the Lower Pliocene, the western border of the thinned Hyblean domain (Sciacca fault) was activated as a dextral shear zone (Civile et al., 2018), while compressional deformation is recognized in the Maghrebian sector lying north of the Adventure block (Pepe et al., 2005) and an extensional regime formed the Pantelleria trough (Civile et al., 2014, 2018, Finetti and Del Ben, 2005b).

- The Malta and Linosa troughs started developing (Catalano et al., 1994; Furlani et al., 2013).

- The Sicilian Apennines underwent a southward bending, attributed to E-W shortening (e.g., Ghisetti et al., 2009), forming the Gela nappe.

- Dextral transpressional deformation, associated with a system of NW-SE faults, is recognized in the Maghrebian belt lying north of the Hyblean-Adventure promontory (Fig. 4, e.g., Catalano et al., 1994, 1996; Sulli, 2000; Guarnieri, 2004; Finetti et al., 2005c).

- An old NW-SE discontinuity in the northern Adriatic foreland reactivated with a sinistral strike-slip regime (the Schio-Vicenza fault system, Fig. 4, e.g., Castellarin and Cantelli, 2000; Zampieri et al., 2003; Massironi et al., 2006; Pola et al., 2014).

- Since the late Miocene, NW-SE to N-S thrusting in the Southern Alps mostly affected the sector lying east of the Schio-Vicenza fault (e.g., Bressan et al., 1998; Galadini et al., 2005), while such activity almost ceased west of that discontinuity.

- In the Northern Dinarides several thrust faults reactivated as dextral strike-slip faults (Placer et al., 2010).

Any attempt at explaining the formation of the Central Tyrrhenian basin as an effect of slab-pull forces should provide plausible answers to some demanding questions:

- Why in the Pliocene would gravitational sinking have only affected the sector of the slab located beneath the Southern Apennines ? (although at that time the slab was well developed under the Calabrian Arc).
- Why would such sinking have only begun in the latest Miocene ? notwithstanding the slab under the Apennines was already well developed in the middle Miocene.
- Why would the effects of slab-pull forces beneath the southern Apennines have ceased in the late Pliocene-early Pleistocene ? i.e. when crustal stretching ended in the Central Tyrrhenian basin.

Other major problems for slab-pull supporters are discussed in the following:

The subsidence predicted by the slab-pull mechanism in the trench zone (Shemenda, 1993; Hassani et al., 1997; Buiter et al., 2001; Hampel and Pfiffner, 2006; Husson, 2006) is just opposite to the uplift that most of the Apennines belt and Calabrian Arc have undergone in the Pliocene and Quaternary (e.g., Westaway, 1993; Pizzi, 2003; Schiattarella et al., 2003; Rusciadelli, 2005, Ghielmi et al., 2013).

Belt-parallel shortening of the whole Apennine chain, evidenced by the formation of major and minor arcs and transversal thrust fronts, as the Olevano-Antrodoco and Sangro-Volturno (e.g., Pizzi and Galadini, 2009; Di Domenica et al., 2012; Ghielmi et al., 2013; Amadori et al., 2019), is not compatible with the trench suction forces implied by slab roll-back.

The Northern Adriatic foredeep does not show any evidence of tilting after the early Pliocene, suggesting that tectonic activity in the Apennine belt can hardly be imputed to slab-pull forces (Brancolini et al., 2019).

231 Since the activation of major fault systems in the Pelagian zone (Sicily Channel) and the
232 northern Adriatic foreland (Schio-Vicenza) can hardly be explained as effects of slab-pull forces
233 beneath the Southern Apennines, there remains the problem of identifying which other driving
234 mechanisms were active in the central Mediterranean zone during the Pliocene. In this regard,
235 Faccenna et al. (2004, 2007) suggest that transtensional deformation in the Sicily Channel could be
236 related to the breaking that would have affected the most curved sector of the wide slab formed
237 during the Balearic phase. Such deep break would have caused the Late Miocene alkaline
238 volcanism in northern Tunisia and Sardinia and deformation in the Sicily Channel zone. However,
239 slab breaking and related magmatism (e.g., Lustrino and Wilson, 2007) do not necessarily
240 presuppose passive lithosphere sinking. Moreover, one must consider that tectonic activity in the
241 Sicily Channel relates to transcurrent fault systems, with some sectors of pull-apart extension,
242 which cannot simply be generated by uprising of magma.
243
244 *Southernmost Tyrrhenian (Pleistocene)*
245 In the early Pleistocene, crustal stretching ended in the Central Tyrrhenian and started
246 developing in the southernmost Tyrrhenian area (Marsili basin, e.g., Finetti and Del Ben, 1986;
247 Sartori, 1990; Savelli, 2002). This extension (Fig. 5) was accompanied by other major tectonic
248 events in the surrounding regions:
249 - In the late Pliocene-Early Pleistocene thrusting underwent slowdown/cessation in the Southern
250 Apennines (e.g., Cello and Mazzoli, 1999; Catalano et al., 2004; Patacca and Scandone, 2004,
251 2007).
252 - Tectonic activity considerably strengthened in the Calabria-Peloritani wedge, with development of
253 major troughs, transversal discontinuities and relative block rotations, accompanied by fast uplift
254 (e.g., Westaway, 1993; Van Dijk and Scheepers, 1995; Zecchin et al., 2004, 2010, 2011, 2015;
255 Finetti, 2005a; Tansi et al., 2007; Del Ben et al., 2008; Spina et al., 2011; Roda-Boluda and
256 Whittaker, 2017; Tripodi et al., 2018).

257 - Thrusting and folding increased along the external front of the Calabrian wedge, with the
258 formation of the External Calabrian Arc complex (e.g., Patacca et al., 1990; Del Ben, 1993; Finetti,
259 2005a).
260 -Tectonic and volcanic activity increased at the Taormina and Palinuro fault systems (e.g., Finetti
261 and Del Ben, 1986; Finetti et al., 1996; Ventura et al., 1999; Savelli, 2002; Peccerillo, 2005).
262 - Thrusting at the front of the Adventure block and extension in the Pantelleria graben slowed down
263 with respect to the Pliocene (Pepe et al., 2005; Civile et al., 2010, 2018).
264 - The southern Adriatic plate underwent upward flexure, with the formation of the Apulian swell
265 (e.g., Finetti and Del Ben, 1986; Tropeano et al., 2002; Santangelo et al., 2012).
266 - In the Early Pleistocene, the northern Adriatic foreland and the surrounding foredeeps were
267 affected by intense subsidence (Ghielmi et al., 2013; Zecchin et al., 2017).
268
269 Explaining the formation of the southernmost Tyrrhenian basin and the coeval tectonic events
270 cited above (Fig. 5) as an effect of slab-pull forces involves some major problems, as discussed in
271 the following.
272 -The uplift that the Calabrian Arc underwent in the Pleistocene cannot be reconciled with the
273 subsidence predicted by gravitational sinking of the underlying slab (Shemenda, 1993; Hassani et
274 al., 1997: Buiter et al., 2001; Hampel and Pfiffner, 2006; Husson, 2006; Faccenna et al., 2014b).
275 - The strong deformation that the Calabria-Peloritani wedge has undergone since the late Pliocene,
276 with the formation of several longitudinal troughs and transversal sphenocasms and the bowing and
277 uplift of that arc, is compatible with belt-parallel compression, which can hardly be taken as an
278 effect of slab roll-back.
279 - The upward flexure of the southern Adriatic domain (Apulian swell) cannot easily be explained as
280 an effect of slab roll back.
281 - The very small width of the Marsili basin (about 50 Km, Fig.5) would imply the roll-back of a
282 corresponding narrow slab. However, such slab shape would require the presence of major

283 decoupling tear faults between the sinking lithosphere and the lateral (not sinking) sectors. When
284 and why did such tears develop ?
285
286 *Recent/present tectonic setting in the Apennine belt*
287 Around the middle Pleistocene (Fig.6), the deformation pattern in the Apennines underwent an
288 important change, recognized by most authors (e.g., Hippolyte et al., 1994; Galadini, 1999;
289 Bartolini, 2003; Piccardi et al., 2006):
290 - Strike-slip tectonics has developed in the southernmost (Lucanian) sector of the Southern
291 Apennines, with the formation of a system of NW-SE sinistral strike-slip faults, accompanied by
292 compressional and tensional features at restraining and releasing stepovers respectively (e.g. Cello
293 et al., 2003; Catalano et al., 2004; Maschio et al., 2005; Ferranti et al., 2009).
294 - In the northern sector of the Southern Apennines, from the Irpinia to Matese zones, a system of
295 normal faults roughly trending NW-SE has developed in the axial part of the belt (e.g., Ascione et
296 al., 2003, 2007; Brozzetti, 2011).
297 - In the Central Apennines, sinistral transtension is recognized in the system of NW-SE normal-
298 oblique faults located in the axial belt (e.g. Galadini, 1999; Piccardi et al., 1999; Galadini and
299 Messina, 2004; Elter et al., 2012).
300 - In the Northern Apennines, sinistral transtensional faulting in the axial belt (e.g., Cello et al.,
301 1998, 2000; Boncio and Lavecchia, 2000; Tondi and Cello, 2003; Piccardi et al., 2006) and
302 thrusting at the outer border (e.g., Boncio and Bracone, 2009; Scisciani and Calamita, 2009;
303 Boccaletti et al., 2011) have occurred. The middle-upper Pleistocene evolution of this belt sector
304 indicates a predominance of vertical motion, with uplift and widening of the Northern Apennine
305 range (Ghielmi et al., 2013).
306 - Uplift has affected the axial and outer sectors of the whole Apennine belt (e.g., Argnani et al.,
307 2003; Bartolini, 2003; Schiattarella et al., 2003, Ascione et al., 2008).

- Compressional features have developed in the inner part of Adria, such as the Middle Adriatic Ridge (Fig. 6, e.g., Scisciani and Calamita, 2009).
- Two major volcanic episodes (Roman and Campanian provinces, Fig.6) occurred in the western side of the Apennine belt (Peccerillo, 2005; Alagna et al., 2010). The emplacement of such volcanism has been related to transtensional faulting (Milia and Torrente, 2003; Acocella and Funiciello, 2006).
- Around the middle-late Pleistocene, the northern part of the Calabrian wedge reaches the continental Adriatic domain and a new major fault system (Sibari), almost parallel to the Adria border develops (Fig. 6, e.g., Guarnieri, 2006; Del Ben et al., 2008; Ferranti et al., 2014; Zecchin et al., 2015, Volpi et al., 2017).
- The Crati and Mesima longitudinal troughs and the Catanzaro and CapoVaticano transversal troughs/faults develope in the Calabrian Arc (e.g., Spina et al., 2011; Zecchin et al., 2015; Tripodi et al., 2018).
- The Vulcano-Syracuse fault system activates (e.g., Finetti and del Ben, 1986; Del Ben et al., 2008; Sulli et al., 2013).
- The Sciacca fault system becomes a sinistral shear zone and is affected by magmatic activity (e.g., Lodolo et al., 2012; Civile et al., 2018, Fedorik et al., 2018).

The present kinematic pattern in the Italian region is fairly well defined by the analyisis of geodetic data observed in more than 700 permanent GPS stations in the period running from January 1, 2001 to December 31, 2018. The horizontal velocity field derived by such data (Fig. 7) with respect to a fixed Eurasian frame (Euler pole at 54.23˚N, 98.83˚W, ω = 0.257˚/Myr, Altamimi et al., 2016) shows that the outer sector of the Apennine belt, including the buried thrusts and folds under the Po Plain, moves considerably faster (4-5 mm/y) and with a greater eastward component with respect to the inner belt (1-2 mm/y). In the Padanian zone that lies west of the Giudicarie fault system velocity values show a significant decrease and a different trend, with respect to the outer

Apennine belt. In southern Italy, the geodetic field confirms the long-term roughly NE ward motion trend of Calabria and the NNW ward motion trend of the Hyblean zone.

To interpret the drastic change of tectonic style that occurred in the Apennine belt around the middle Pleistocene as an effect of slab-pull forces, one should explain why the effect of such deep driving mechanism on shallow structures has considerably changed at that time. An attempt in this sense has been made by some authors (e.g., Faccenna et al., 2014b), who suggest that the Pleistocene deformation pattern in the Apennine belt has been controlled by changes of deep seated dynamic processes (mantle convection) connected with the development of windows in the western subducted margin of Adria located beneath the Central and Southern Apennines (e.g., Wortel and Spakman, 2000; Piromallo and Morelli, 2003; Faccenna et al., 2007)**.** However, this interpretation cannot easily be reconciled with some major features of the observed deformation pattern, as discussed in the following:

- Pleistocene uplift is recognized in the whole belt, from the Northern Apennines to Calabria, although this effect would only be expected in the belt sectors (Central and Southern Apennines) located above the supposed slab windows.
- The velocity field derived from GPS measurements in the Italian region (Fig. 8) indicates that the outer sector of the Apennine belt is moving faster (4-5 mm/y NEward) than the inner sector (1-2 mm/y N to NWward). This almost homogeneous kinematic pattern all along the whole belt can hardly be imputed to mantle upwelling above slab windows in the Central and Southern Apennines.
- The Hyblean wedge, confined by tectonic zones (Sicily channel, Syracuse fault and Maghrebian belt) is moving roughly North to NW ward, while the Calabrian wedge is moving roughly ENE ward (Fig. 8). Explaining such microplate kinematics as an effect of mantle upwelling in the Southern and Central Apennines appears to be a rather difficult task.

3. Proposed geodynamic interpretation

Main concepts

The observed deformation pattern in the Mediterranean region has been driven by the convergence of the Africa/Adriatic and Eurasia plates and by the roughly westward motion of the Anatolian-Aegean-Pelagonian belt, induced by the indentation of Arabia (e.g., Mantovani et al., 2006, 2009, 2014; Viti et al., 2009). The shortening of the interposed Mediterranean structures, constituted by orogenic belts, oceanic zones and the Adriatic continental and thinned continental domains, has been accommodated by a variety of tectonic processes, whose spatio-temporal distribution has been controlled by the well know least-action principle, aimed at minimizing the resistance of gravity, fault friction and mantle viscosity.

It is known that in a buoyancy-controlled context the most efficient way to gain space in a constricted area is the subduction of lithosphere and that such process is mainly favored when oceanic domains are involved (e.g., Cloos, 1993; Stern, 2004). However, one must take into account that the consumption of such domains cannot simply be determined by horizontal compression exerted by plate convergence. This consideration is mainly suggested by the fact that during the long collisional phase between the Africa-Adriatic plate and Eurasia, the Ionian oceanic domain did not undergo any subduction beneath the African and Adriatic continental domains. The fact that during such phase the push of Africa was very efficiently transmitted by the Ionian zone to the continental Adriatic domain is testified by the compressional deformations that took place along the northern boundary of that promontory (e.g., Castellarin and Cantelli, 2000; Ceriani and Schmid, 2004; Rosemberg and Kissling, 2013; Handy et al., 2015). The physical plausibility of the above evidence is confirmed by the computation of rheological profiles (e.g., Viti et al., 1997), which indicates that in long-term (geological) time intervals the horizontal compressional strength of the oceanic lithosphere is not lower than the one of a continental domain. That old and cold oceanic lithosphere can effectively transmit horizontal compression is also suggested by the persistence of continental collision at the Himalayan-Tibet boundary, without any new subduction zone in the

386 central Indian Ocean (e.g. Stern, 2004; Copley et al., 2010). Another example is the Black Sea
387 oceanic lithosphere, which has transmitted compression from northern Anatolia to central-southern
388 Eurasia since Eocene (e.g., Hippolyte et al., 2018).

389 To understand why the consumption of oceanic domains has often occurred in the
390 Mediterranean region since the Oligocene, it is necessary to take into account that the lateral escape
391 of orogenic wedges may have created the conditions which can make that process feasible. When
392 the margin of an oceanic domain is overthrust by extruding orogenic material, it undergoes
393 downward flexure, due to isostasy. This deformation perturbs the previous equilibrium between
394 horizontal forces, triggering the sink of the denser lithosphere, under the action of plate
395 convergence (see e.g. the mechanism of subduction initiation given by Hall, 2018).

396 Laboratory experiments (Driehaus et al., 2013) show that the extrusion of a continental belt over
397 an adjacent oceanic domain (driven by belt-parallel compression) may induce subduction when the
398 density ratio between the oceanic and continental plates is larger than 1.4.

399 The above considerations can explain why the consumption of most Tethyan zones has often
400 occurred since the Oligocene, when most extrusion processes have developed in the study area (e.g.,
401 Mantovani et al., 2006, 2009, 2014; Viti et al., 2009).

402 Another type of tectonic reorganization that may allow the activation of more convenient
403 shortening processes in a constricted context, involves a change of plate mosaic, through the
404 activation of major decoupling faults. In the new plate configuration, the kinematic pattern of
405 buoyant blocks (microplates and orogenic wedges) is reorganized in order to address most intense
406 stresses towards the less buoyant domains. Two major examples of this kind of tectonic
407 reorganization have occurred in the middle Miocene and late Miocene in the central Mediterranean
408 area, as discussed in the next sections.

409 Laboratory (e.g., Ratschbacher et al., 1991; Davy et al., 1995; Faccenna et al., 1996; Driehaus
410 et al., 2013; Boutelier et al., 2018) and numerical (e.g., Mantovani et al., 2000, 2001b, 2007b)
411 experiments suggest that in constricted buoyant structures the lateral escape of wedges (arcs)

412 is the most convenient shortening process and that in the internal side of an extruding/migrating
413 arc crustal extension may develop. Extension occurs when the divergence between the migrating arc
414 and the stable foreland is faster than the shortening induced by plate convergence.
415
416 *Balearic basin*
417 The development of the Balearic ATBA system (Fig.2) was triggered by the collision between
418 the northern margin of the African continent and the southernmost edge of the Al-Ka-Pe-Ca belt
419 (the Atlas compressional phase, e.g. Benaouali-Mebarek et al., 2006; Tesòn and Teixell, 2008;
420 Frizon de Lamotte et al., 2009). After the remarkable increase of resistance induced by that
421 collision, the prosecution of the Africa-Eurasia convergence was allowed by a peculiar shortening
422 process, given by the eastward bowing/extrusion of the Al-Ka-Pe-Ca-Corsica-Sardinia arc, at the
423 expense of the adjacent Tethyan domain (Fig. 2). Crustal extension developed in the wake of the
424 migrating arc, leading to the formation of the Balearic and Algerian basins, while thrustings
425 developed in the trench zones (Apennines and Maghrebides). A detailed description of the proposed
426 geodynamic interpretation and of how it can account for the observed deformation pattern in the
427 Western Mediterranean region is given by Viti et al. (2009).
428
429 *Northern Tyrrhenian basin (9-6 My)*
430 The conditions that led to the formation of the Northern Tyrrhenian basin were created by the
431 reactivation of a major old discontinuity in the Northwestern part of the Adriatic promontory (the
432 Giudicarie fault system, Fig. 9a,b). This decoupling allowed to overcome the critical situation that
433 had gradually developed along the collision zone between the Adriatic promontory and the
434 European plate. In the Oligocene-Lower Miocene, such collision caused the accumulation of a huge
435 amount of light crustal material in the trench zone (e.g., Schmid et al., 2004; Finetti, 2005b). Thus,
436 the resistance of gravity against any further underthrusting was getting higher and higher.

The possibility of mitigating such critical situation was offered to the Adriatic promontory by the development, in the early-middle Miocene of a weak lateral boundary zone, constituted by the Carpatho-Pannonian ATBA system (e.g., Royden et al. 1983; Tari and Pamic, 1998; Horvath et al, 2015, Rèka et al., 2018; Mantovani et al., 2006, 2009). The presence of that tectonized structure allowed the eastward extrusion of buoyant crustal wedges from the Eastern Alps (a process also revealed by the formation of the Tauern window in the wake of the extruding wedges, Fig.4b, e.g., Ratschbacher et al., 1991; Robl and Stuwe, 2005; Wolfler et al., 2011), which favoured the NNEward displacement of the Adriatic domain (Peresson and Decker, 1997; Frisch et al., 2000). However, this displacement could only occur after the decoupling of the main Adriatic domain from its northwestern protuberance, which at that time was deeply stacked beneath the Western Alps. Such decoupling was allowed by the reactivation, around the Tortonian (Martin et al. 1998; Castellarin and Cantelli, 2000; Viola et al., 2001; Fellin et al., 2002), of an old discontinuity in the northern Adriatic foreland, the Giudicarie fault system (Fig. 9b), where sinistral transpressional activity is recognized until the early Messinian (e.g., Favaro et al., 2017). The NNE ward displacement, with clockwise rotation, of the northern Adriatic domain that followed such decoupling allowed that promontory to release the internal elastic deformation that such indenter had accumulated during its oblique collision with the European domain.

The sinistral motion between the decoupled Adriatic domain and its northwestern Padanian protuberance (by then closely connected with the Western Alps) is testified by the fact that since then thrusting activity in the Alps mostly occurred in the sector lying east of the Giudicarie fault (e.g., Frisch et al., 2000, Viola et al., 2001; Zampieri et al., 2003; Castellarin et al., 2004).

The consequent divergence between the mobile Adriatic domain and the Corsica-Sardinia block (stable since the middle Miocene, Gattacceca et al., 2007 and references therein) caused crustal extension in the interposed zone, constituted by a sector of the Alpine-Apennine orogenic belt lying north of the Selli fault, leading to the formation of the Northern Tyrrhenian basin (Fig.9b). This hypothesis can explain why crustal extension affected the zone comprised between

the Adriatic promontory and the Corsica-Sardinia block, why such activity started around the middle Tortonian (about 9 My), just after the reactivation of the Giudicarie fault system, and why the extended area was confined to the South by the Selli fault.

Central Tyrrhenian basin and other major coeval tectonic events

Since the late Miocene, tectonic activity in the Mediterranean region was significantly influenced by the westward displacement of the Anatolian-Aegean-Pelagonian belt (Fig.1). In the previous evolution, this last kinematic boundary condition did not have significant effects in the Central Mediterranean area, because the convergence between the Africa-Adriatic plate and the above belt was mainly accommodated by the consumption of the interposed thinned domain (the Ionian zone in figure 9a, e.g., Robertson and Shallo, 2000). The resistance against this convergence considerably increased in the upper Miocene, when more buoyant domains reached that consuming boundary (Mercier et al., 1987; Sorel et al., 1992). This critical situation, with progressive slowdown of accretionary activity, lasted up to the late Miocene-early Pliocene, when a drastic change of plate mosaic and kinematic pattern allowed other less resisted shortening processes to occur. This reorganization started by decoupling a large portion of the Adriatic promontory (Adria plate in figure 9c) from Africa, through the activation of a long fracture, the Sicily Channel-Medina-Victor Hensen transtensional fault system. Prior to this tectonic phase, the Hyblean-Adventure promontory was part of the undeformed African foreland (Pelagian zone, e.g. Finetti and Del Ben, 1986, 2005b; Hieke et al., 2003, 2006; Lentini et al., 2006; Fedorik et al., 2018). The new motion trend of Adria required another major decouplig (at least partial) in the northern Adriatic zone, which was achieved by the reactivation of an old weak zone, the Schio-Vicenza sinistral fault system (Fig. 9c). After that decoupling, the northern Adria domain moved roughly NNW ward, as indicated by a change in the orientation of the compressional axis from SW-NE to SSE-NNW in the eastern Southern Alps (e.g., Castellarin and Cantelli, 2000) and by the reactivation of many thrust zones as right lateral strike-slip faults in the Northern Dinarides (Placer et al., 2010). The

489 decoupling of the Adria plate from Africa and the new kinematics of that plate (Fig. 9c) avoided an
490 highly resisted collision with the Anatolian-Aegean-Pelagonian belt.

491 The E-W convergence between southern Adria and the northern African promontory (the
492 present Algeria-Tunisia zone) required the roughly Northward expulsion of an African fragment
493 (the Adventure block), guided by the Egadi and Sciacca fault systems (Fig. 9c,d), and the E-W
494 shortening of the Hyblean domain. The extrusion of the Adventure block in the early Pliocene may
495 explain the thrusting recognized at the outer front of that wedge (in the Maghrebian belt, Pepe et al.,
496 2005), the formation of the Pantelleria trough at the inner side of the same block (Civile et al., 2010,
497 2018) and the dextral shear recognized in the Sciacca fault system (Civile et al., 2018). The E-W
498 shortening of the Hyblean domain, constituted by thick and thin zones (Fig. 9c,d), was
499 accommodated by the southward bending of the Sicilian Apennines, with the consequent formation
500 of the Gela nappe (at the expense of the thinned Hyblean domain), and by fracturation and
501 redistribution of small ridge fragments in the Sicily channel zone (see i.e. Finetti and Del Ben,
502 1986). The occurrence of local extension between diverging fragments caused the formation of the
503 Malta and Linosa troughs, as tentatively reconstructed in figure 10.
504
505 The roughly northward displacement of Pelagian blocks is also suggested by the fact that the
506 sector of the Alpine (Kabylo-Calabrides)-Maghrebian belt which lies in front of them (e.g., Ben
507 Avraham et al., 1990; Lentini et al., 1994; Sulli, 2000; Gueguen et al., 2002) shows a northward
508 shift with respect to the lateral North African and Calabrian sectors (see fig. 10 of Mantovani et al.,
509 2007b and references therein).
510 The complex evolution of the Sicily channel tectonic zone may have favored the ambiguity that
511 still surrounds the shear sense in that transcurrent fault system (e.g., Grasso and Pedley, 1985;
512 Boccaletti et al., 1987; Cello, 1987; Reuther et al., 1993; Kim et al., 2003; Finetti and Del Ben,
513 1986, 2005b; Catalano et al., 2009). Since the break of the thick continental African domain is a
514 major tectonic event in the evolution of the central Mediterranean region, any hypothesis advanced

about the nature (dextral or sinistral strike slip or pure passive rifting) and the genetic mechanism of that discontinuity should also provide a geodynamic justification for the coeval deformations observed in the surrounding zones. For instance, one should consider that a dextral movement at the Sicily Channel fault system would imply a SE ward motion of the Hyblean domain with respect to Africa. However, the deformation predicted by such kinematics in the surrounding zones (shortening in the Ionian zone facing the Hyblean block and extension in the zone lying between the Hyblean promontory and Sardinia) are not compatible with the observed features. No evidence of shortening is recognized in the Ionian area, especially at the Syracuse escarpment, as shown by CROP seismic sections (e.g., Finetti and Del Ben, 2005c) and a considerable shortening, instead of the expected extension, is evidenced by the CROP section crossing the Sardinia and Sicily Channels (Finetti et al., 2005c). We would like to remark that the geodynamic framework we propose as responsible for the formation of the Sicily Channel tectonic zone may provide plausible explanations for the major coeval late Miocene-Pliocene tectonic events in the central Mediterranean region (as discussed in the text), whereas this result cannot be achieved with the very general alternative interpretations so far proposed.

We suppose that the occurrence of crustal extension in the Sicily Channel tectonic zone may be an effect of pull-apart troughs at step-overs of the main transcurrent faults (as suggested by Reuther et al., 1993). This hypothesis is compatible with the occurrence of volcanic activity. In fact, the most plausible genetic mechanism of magma uprise through the crust is generally considered the one involved by pull-apart troughs (e.g., Tamburelli et al., 2000; Gudmundsson, 2001; Acocella and Funiciello, 2006). The normal faults generated by a pure extensional regime (in diverging plate boundaries, for instance) are stressed by a very high compression, which does not allow uprising of magmas.

The northward displacement of the Adventure block played an important role in the formation of the Central and Southern Tyrrhenian basins and in the subsequent evolution of the Apennine belt, since the indentation of that continental fragment onto the Alpine-Apenninic orogenic material

541 which lay south of the Selli fault caused the lateral escape of wedges, at the expense of the Ionian
542 Tethys and the thinned margin of the Adriatic domain (Fig. 9c,d). This process may explain why
543 since the late Miocene intense accretionary activity occurred in the Apennine belt, in front of the
544 extruding wedges (Viti et al., 2006 and references therein), and crustal extension developed in the
545 wake of those wedges, with the formation of the central Tyrrhenian basin. The relatively high
546 velocity (up to 5-8 cm/y) of the consequent trench retreat recognized at the related consuming
547 boundary, during the Pliocene (Patacca and Scandone, 1989; Finetti et al., 2005b; Guillaume et al.,
548 2010), might be due to the contemporaneous actions of two opposite kinematic boundary
549 conditions, i.e. the roughly NW ward motion of southern Adria and the roughly ESE ward
550 migration of the Alpine-Apennines wedges (Fig.9c,d).

551 During this phase, belt-parallel compression also stressed the central-northern Apennines,
552 causing the formation of arcs, with in-sequence thrusting at external fronts (e.g., Calamita et al.,
553 1994; Costa, 2003, Ghielmi et al., 2013; Brancolini et al., 2019), out-of-sequence thrust
554 reactivations (e.g., Boccaletti et al., 1999) and extensional to transtensional tectonics in the internal
555 side of the arcs, accompanied by regional uplift (e.g., Martini and Sagri, 1993; Bossio et al., 1998).
556 The formation of arcs is also suggested by the fact that during this phase the geometry of foredeeps
557 in the Northern Apennines changed from continuous, cylindrical, to highly fragmented, non
558 cylindrical (e.g., Amadori et al., 2019). The hypothesis that in the Pliocene the belt underwent a
559 compressional regime is also suggested by the shortening evidenced by CROP seismic sections
560 (e.g., Finetti et al., 2005a).

561 The geodynamic interpretation here proposed may explain why the coeval occurrence of
562 compression in the outer belt and extension in the inner side of the northern Apennines has
563 developed since the late Miocene (e.g., Elter et al., 1975; Bossio et al., 1993). The fact that the
564 above deformation pattern was accompanied by anatectic magmatism (e.g., Peccerillo, 2003) could
565 be explained by considering that the most physically plausible genetic mechanism of the crustal
566 pathways that allow magmas to uprise through the upper crust is the transtensional regime that is

567 expected to occur at the inner side of extruding wedges (e.g., Tamburelli et al., 2000;
568 Gudmundsson, 2001; Acocella and Funiciello, 2006).

569 The fact that lateral escape of wedges may have contemporaneously involved belt-parallel
570 shortening and perpendicular extension in the northern Apennines could explain why both types of
571 strain styles are recognized in that zone (e.g., Bonini et al., 2014; Brogi et al., 2013; Liotta et al.,
572 2015). Anyway, the geodynamics here proposed provides that extensional deformation in the above
573 Apennine sector is the dominant deformation, as indicated by most structural and morphological
574 evidence (e.g., Brogi et al., 2013).

575 During the Late Miocene-Pleistocene interval, a complex system of elongated foredeeps
576 developed in the eastern sector of the Po Plain and in the northern Adriatic area and severe tectonic
577 activity affected the northern Apennines and the Padanian area (Ghielmi et al., 2013).

578 The peculiar fact that the Apennine belt has contemporaneously undergone thrusting, uplift
579 and transtensional deformations is compatible with the strain pattern expected from belt-parallel
580 compression, as suggested by the results of numerical experiments (Viti et al., 2004; Mantovani et
581 al., 2007b), which show that the deformation pattern observed in the Tyrrhenian-Apennines system
582 since the latest Miocene can be reproduced as an effect of the kinematic boundary conditions
583 shown in Fig. 9c,d.

584

585 *Southern Tyrrhenian basin (Early Pleistocene)*

586 When the extruding Southern Apennine wedge reached the continental Adriatic domain, the
587 resistance against such consuming process underwent a significant increase, which led to the
588 progressive stop of that wedge, revealed by the end of thrusting at its outer front and of crustal
589 stretching in the Central Tyrrhenian basin, at the inner side of the wedge.

590 After that trench suture, the convergence of the confining plates was mainly accommodated by the
591 fast lateral escape of the Calabria-Peloritani (CP) wedge, i.e. the only sector of the belt that was
592 still facing an oceanic domain (the Ionian Tethys, Fig. 9d).

593 The acceleration of the CP wedge is mainly testified by the strengthening of accretionary
594 activity along its outer front, which built up the External Calabrian Arc, and by the occurrence of
595 crustal stretching in the wake of that wedge, which has generated the Marsili basin. In this regard, it
596 can be noted that the width of this basin is comparable with the internal (Tyrrhenian) side of the CP
597 wedge (Fig.9d). The lateral guides of the above extrusion process were constituted by the Taormina
598 and Palinuro transcurrent fault systems, as suggested by acceleration of tectonic and volcanic
599 activity along those faults.

600 Nicolosi et al. (2006) suggest that the spreading of the Marsili basin has mainly developed
601 from 2.1 to 1.6 Ma (with a very high rate, 19 cm/y) and that since 0.78 Ma spreading has
602 undergone a considerable slow down. The initial high spreading rate could be due to the fact that
603 such wedge had to extrude through a very narrow corridor between lateral continental domains
604 (Africa and Adria). The subsequent slowdown of the CP wedge could be due to the collision of
605 northern Calabria with the continental Adria domain (e.g., Del Ben et al., 2008, Zecchin et al.,
606 2004, 2011, 2015; Mantovani et al., 2019 and references therein).

607 The strong belt-parallel compressional regime that stressed the CP wedge during its extrusion
608 can explain the fast uplift, the horizontal bowing and the strong fragmentation of that structure,
609 with the formation of transpressional fault systems, troughs and sphenocasms (e.g. Ghisetti and
610 Vezzani, 1981; Van Dijk and Schepeers, 1995; Monaco and Tortorici, 2000; Tansi et al., 2007; Del
611 Ben et al., 2008; Zecchin et al., 2004, 2010, 2011, 2015; Roda-Boluda and Whittaker, 2017 and
612 references therein).

613 The fast uplift of the CP wedge has been alternatively explained as an effect of isostatic
614 rebound in response to breakings of the underlying Ionian slab (e.g., Westaway, 1993; Wortel and
615 Spackman, 2000) or as due to the decoupling of the Calabrian arc from the underlying slab by
616 convective removal of the deep root (e.g., Gvirtzman and Nur, 2001). However, these
617 interpretations cannot easily account for the reconstruction of the Apennine-Maghrebian slab
618 geometry based on several studies (e.g., Massari and Prosser, 2013 and references therein), which

suggests that since about 10 My two large slab windows have developed beneath that belt, one in the sector running from Tunisia to eastern Sicily and the other beneath the central and southern Apennines. If this reconstruction is reliable, one can hardly understand why the most intense uplift expected from isostatic rebound has affected the Calabrian Arc.

Since the suture of the Southern Apennines consuming boundary, in the Late Pliocene-Early Pleistocene, the strong compressional regime induced by the surrounding plates in the southern Adria domain, being not anymore absorbed by the consumption of thinned domains, was accommodated by upward flexure, which accelerated the formation of the Apulian swell. The uplift of southern Adria could have induced an opposite vertical effect in the northern edge of that plate, which might explain the subsidence that affected such zone in the Early Pleistocene (Ghielmi et al., 2013; Zecchin et al., 2017; Brancolini et al., 2019).

Recent present tectonic setting in the Apennine belt (Middle-Late Pleistocene)

Since the middle Pleistocene, the gravitational energy accumulated by the southern Adria domain in the previous Early Pleistocene phase favored the northward displacement of that plate (Fig. 9e), as suggested by the resumption of thrusting and strike-slip tectonics in the eastern (Dinarides) and northern (Eastern Alps) boundaries of Adria (Viti et al., 2006 and references therein). A more complex effect of Adria's acceleration has developed along its western boundary, the Apennine belt. Since the external (eastern) sector of that chain was more closely connected with the underlying Adriatic domain, it underwent a more efficent drag from Adria, with respect to the inner belt (which was overlying a much deeper Adriatic lithosphere). This drag has resulted in a greater mobility, stronger deformation and uplift of the outer belt, which has progressively separated from the inner belt, forming the series of troughs located in the axial chain. This mechanism may explain why since the middle Pleistocene the compressional tectonic style in the Apennines was replaced by a dominant left lateral transcurrent regime (Viti et al., 2006 and references therein). The compressional features previously developed in the two main Pliocenic

645 extruding wedges, the Molise-Sannio (MS) and the Romagna-Marche-Umbria (RMU), were cut by
646 a series of longitudinal trastensional fault sytems, as the Irpinia-Benevento-Matese (e.g., Pantosti
647 and Valensise, 1990; Cinque et al., 2000; Ascione et al., 2007) in the MS wedge and the Norcia-
648 Colfiorito-Val Tiberina (Calamita et al., 2002; Pizzi and Galadini, 2009) in the RMU wedge,
649 connected by longitudinal transtensional fault systems (Aquila and Fucino) in the Lazio-Abruzzi
650 platform (Fig. 10, Viti et al., 2006 and references therein). This dynamics also induced belt-parallel
651 shortening in the outer belt, accommodated by the formation of arcs (e.g., Maiella, Gran Sasso,
652 Laga Mt.) and a generalized uplift (e.g., Pizzi and Galadini, 2009; Elter et al., 2012; Blumetti et al.,
653 2013).

654 In the northernmost belt (Romagna-Emilia Apennines), shortening has been accommodated by
655 the formation of various arcs, also involving the buried folds beneath the Po valley, and progressive
656 outward migration of thrust fronts (Cerrina Feroni et al., 2001; Costa, 2003; Vannoli et al., 2004,
657 2015; Boccaletti et al., 2011; Ghielmi et al., 2013; Chicco et al., 2019).

658 The relative motion between the southernmost sector of the mobile outer belt (MS wedge) and
659 Calabria has been accommodated by a system of transcurrent faults in the Lucanian Apennines (Fig.
660 11, Viti et al., 2006 and references therein).

661 The recent/present kinematic pattern of the Apennine belt inferred from Pleistocene
662 deformation pattern, involving a faster motion of the outer Apennine belt with respect to the inner
663 belt (Fig.11), fairly well agrees with the velocity field derived by geodetic observations (Fig. 7).

664 It can be noted as well that the motion trend of Adria indicated by several GPS velocity vectors
665 in the Apulia zone and in the Venetian plain (Fig.7) is compatible with the NNEward Africa-
666 Eurasia convergence trend suggested by Mantovani et al. (2007a), which can explain the absence of
667 major decoupling zones inside the Adriatic domain. If conversely, one adopts the roughly NNW
668 ward convergence trend provided by global kinematic models (e.g., Calais et al., 2003) it becomes
669 necessary to identify major tectonic discontinuities able to decouple the Adria plate from Africa.
670 However, the very different solutions so far suggested by the numerous attempts in this sense (see

671 Babbucci et al., 2004 and references therein) clearly indicate the scarce significance of the available
672 evidence about possible decoupling zones inside the Adria plate.

673 The lower GPS velocities of the northern Adriatic zone (2-3 mm/y) with respect to the southern
674 Adriatic domain (5-6 mm/y) could be connected with a non rigid behaviour of the Adria plate, due
675 to the non uniform time distribution of decoupling earthquakes along the circum Adriatic
676 boundaries (Mantovani et al., 2015a). In this regard, one could suppose that the Pleistocene
677 compressional deformations recognized in the inner part of Adria (such as the Middle Adriatic
678 Ridge, Fig. 6, e.g., Scisciani and Calamita, 2009) may have developed during such transitory
679 phases, characterized by an accelerated motion of southern Adria and a low mobility of the northern
680 Adria (Mantovani et al., 2016).

681

682 The belt-parallel compression and the sinistral shear that affected the Apennine belt during the
683 last evolutionary phase may have emphasized transtensional stresses in the inner side of the two
684 main Pliocenic wedges, the Molise-Sannio and the Romagna-Marche Umbria (Fig.11), favouring
685 the uprise of magmas in the Roman and Campanian magmatic provinces (e.g., Argnani and Savelli,
686 1999; Tamburelli et al., 2000; Peccerillo, 2005; Finetti, 2006; Tibaldi et al., 2010). This could
687 provide a possible explanation for the two major features, i.e. the location and timing, of the most
688 intense Quaternary volcanic episodes that has developed in the Apennine belt during this
689 evolutionary phase.

690 Around the middle Pleistocene, a major reorganization of the tectonic setting has been caused
691 by the collision of southern Calabria with the continental Adriatic domain (Fig. 9e). Such obstacle
692 was overcome by the activation of the Sibari fault system (e.g., Volpi et al., 2017) which has
693 allowed the Calabrian wedge to gain a new extrusion trend, more parallel to the Adriatic border
694 (Fig. 9e). This change also had major effects in the interaction zone with the Hyblean domain,
695 favoring the activation of the Vulcano-Syracuse fault system, which has become the main
696 decoupling zone between the Calabrian wedge and the Hyblean block, that were moving in almost

opposite directions. After the activation of such decoupling, the Hyblean block may have accelerated its northward motion, as suggested by the fact that the border with the Ventura block (Sciacca fault) has become a sinistral shear zone (Civile et al., 2018). The fact that magmatic activity in the Sciacca fault system has mainly developed in the Pleistocene could suggest that the inversion of the strike slip sense at that fault may have favoured the development of pull-apart mechanisms. The hypothesis that the Syracuse fault is an active decoupling zone is supported by the occurrence of great damages in the eastern side of Sicily (e.g., 1169 and 1683, Rovida et al., 2016).

The reliability of the evolution here proposed may also be supported by the fact that major features of the spatio-temporal distribution of main earthquakes in the periAdriatic regions are compatible with the short-term implications of the present tectonic setting, as discussed in a number of papers (Mantovani et al., 2010, 2012, 2015b, 2016, Viti et al., 2006, 2012, 2013, 2015a,b, 2016).

4. Conclusions

Discussions about Mediterranean geodynamics mainly concern the driving mechanism of the ATBA systems. To overcome the apparent difficulty raised by the occurrence of crustal extension in zones of plate convergence, some authors suggest that such tectonic process may be produced by deep seated forces, mainly induced by gravitational sinking of subducted lithosphere. However, the implications of that genetic mechanism can hardly be reconciled with the observed deformation pattern. In particular, the development of the northern, central and southern Tyrrhenian basins in three well distict phases, would require a discontinuos, very peculiar and scarcely plausible action of gravity on the Adriatic subducted margin. This and other major problems discussed in the text suggest that slab-pull forces can hardly be invoked to explain the surface deformation pattern.

So far, many laboratory and numerical experiments, both 2D and 3D, have investigated the dynamics of slab-pull. However, the parameters adopted in various experiments vary over wide ranges, making it difficult to compare among results provided by different works. For instance, one may consider the density difference between the slab and surrounding mantle ($\Delta\rho$), which crucially affects the magnitude of the slab-pull force. Capitanio et al. (2010) use the values $\Delta\rho$ = 30, 60 and 90 kg m^{-3} for their numerical experiments. Schellart and Moresi (2013) instead adopt the value $\Delta\rho$ = 80 kg m^{-3}. Meyer and Schellart (2013) consider the significantly larger value $\Delta\rho$ = 103 kg m^{-3}, which obviously provides a larger slab-pull force. Furthermore, some attempts disregard basic features of the subduction systems, in particular the role of the upper/overriding plate (e.g., Funiciello et al., 2003a,b, 2004, 2006; Schellart, 2010; Schellart et al., 2011). On the other hand, the upper plate may effectively resist slab retreat and back-arc extension (Shemenda, 1993; Hassani et al., 1997; Capitanio et al., 2010). Thus, the conclusions drawn by the above works cannot be considered as definitive ones, because they depend on questionable experimental settings or on still poorly constrained subduction parameters.

Furthermore, one should consider that the geodynamic interpretations based on the slab-pull genetic mechanism, advanced to explain the development of ATBA systems, can hardly account for the occurrence of other coeval major tectonic events in the central Mediterranean area, such as the activation of important fractures in the African foreland (the Sicily Channel-Medina-Victor Hensen fault system) and in the northern Adriatic region (the Giudicarie and the Schio-Vicenza fault systems), along with some peculiar variations of tectonic style in the periAdriatic belts.

In this and previous papers (Mantovani et al., 2006, 2009, 2019; Viti et al., 2006, 2009) it is suggested that the observed deformation pattern in the central-western Mediterranean has been driven the convergence of the confining plates (Africa, Eurasia and Anatolian-Aegean-Pelagonian system). The migration of arcs and the consequent occurrence of crustal extension in their wake can plausibly be explained as effects of extrusion processes, that develop where orogenic belts lying aside oceanic domains are stressed by plate convergence (Fig. 8). Some considerations about why

748 the proposed geodynamic interpretation may better explain the evolution of the study area are given
749 in the following.
750 - It takes into account a very large set of tectonic features, more complete than the ones considered
751 in other attempts.
752 - All deformations considered are tentatively explained as effects of a unique driving mechanism
753 (the convergence of the confining plates). In particular, it is taken into account the influence of an
754 important kinematic boundary condition, the westward motion of the Anatolian-Aegean-Pelagonian
755 belt, that has been often neglected by other attempts.
756 - It is based on clear tectonic concepts, related to the well known least-action principle, which
757 allows finding plausible explanations for coeval tectonic events, even if located very far from each
758 other. For instance, this scheme may allow to understand which was the possible connection
759 between the activation of the Sicily Channel-Medina-Victor Hensen fault system in the
760 Pelagian/Ionian zone and the reactivation of the Schio-Vicenza fault in the northern Adriatic
761 domain.
762 - The present tectonic setting resulting from the proposed evolutionary reconstruction allows
763 identifying plausible explanations for major features of the spatio-temporal distribution of major
764 earthquakes in the last centuries.

765 In literature, many works concerning the Mediterranean region start by citing the various
766 geodynamic interpretations so far proposed for this area, underlying the considerable ambiguity that
767 still surrounds this problem. This uncertainty involves many negative implications, even concerning
768 important social problems, as for instance the mitigation of seismic hazard. In fact, it is well known
769 that a reliable attempt at recognizing the zones most prone to next strong shocks can only be made
770 if one can rely on a deep knowledge of the ongoing tectonic processes. Exploiting this knowledge
771 and the seismic history of the zone considered, one could try to recognize the perturbation of the
772 strain/stress fields that may be caused by major earthquakes and how this effect may influence the
773 subsequent spatio-temporal distribution of seismicity (Mantovani et al., 2015b, 2016, 2017; Viti et

al., 2015a, 2016). However, since the reliability of the above results is heavily conditioned by the reliability of the adopted tectonic model, it is crucially important to exploit the information now available on the past deformations in order to recognize which driving mechanisms are actually stressing the Italian region. Thus, the Scientific Community should make all efforts to overcome the presumed ambiguity about this problem. This work aims at providing a contribution in this direction, by analyzing the plausibility of the geodynamic interpretations so far proposed, first focusing attention on the models most often cited in literature.

References

Acocella, V., Funiciello, R. 2006. Transverse systems along thee Tyrrhenian margin of Central Italy and their influence on volcanism. Tectonics, 25, TC2003. http://dx.doi.org/10.1029/2005tc001845

Alagna, K., E., Peccerillo, A., Martin S., Donati C., 2010. Tertiary to Present Evolution of orogenic magmatism in Italy, in: Beltrando, M., Peccerillo, A., Mattei, M., Conticelli, S., Doglioni, C. (Eds.) The geology of Italy: tectonics and life along plate margins. J. Virtual Explorer, 36, 18. doi:10.3809/jvirtex.2010.00233.

Altamimi, Z., Rebischung, P., Metivier, L.,, Collilieux, X., 2016. ITRF2014: A new release of the International Terrestrial Reference Frame modeling nonlinear station motions, J. Geophys. Res., Solid Earth, 121, 6109-6131. doi:10.1002/2016JB013098

Amadori, C., Toscani, G., Di Giulio, A., Maesano, F.E., D'Ambrogio, C., Ghielmi, M., Fantoni, R., 2019. From cylindrical to non-cylindrical foreland basin:Pliocene–Pleistocene evolution of the Po Plain–Northern Adriatic basin (Italy). Basin Res., 2019, 1–25. DOI: 10.1111/bre.12369

Argnani, A., Savelli, C., 1999. Cenozoic volcanism and tectonics in the southernTyrrhenian sea: space-time distribution and geodynamic significance. J. Geodyn., 27, 409-432.

799 Argnani, A., Barbacini, G., Bernini, M., Camurri, F., Ghielmi, M., Papani, G., Rizzini, F., Rogledi, S.,
800 Torelli L., 2003. Gravity tectonics driven by Quaternary uplift in the Northern Apennines: insights
801 from the La Spezia-Reggio Emilia geo-transect. Quatern. Int., 101-102, 13-26.
802 Ascione, A., Cinque, A., Improta, L., Villani, F. 2003. Late Quaternary faulting within the Southern
803 Apennines seismic belt: new data from Mt. Marzano area (southern Italy). Quatern. Int., 101-102, 27-
804 41. https://doi.org/10.1016/S1040-6182(02)00127-1
805 Ascione, A., Caiazzo, C., Cinque, A., 2007. Recent faulting in Southern Apennines (Italy): geomorphic
806 evidence, spatial distribution and implications for rates of activity. Ital. J. Geosci., 126, 293-305.
807 Ascione, A., Cinque, A., Miccadei, E., Villani, F., 2008. The Plio-Quaternary uplift of the Apennines Chain:
808 new data from the analysis of topography and river valleys in Central Italy. Geomorphology, 102, 105-
809 118, http://dx.doi.org/10.1016/j.geomorph.2007.07.022
810 Babbucci, D., Tamburelli, C., Viti,M., Mantovani, E., Albarello, D., D'Onza, F., Cenni, N., Mugnaioli, E.
811 2004. Relative motion of the Adriatic with respect to the confining plates: seismological and geodetic
812 constraints. Geophys. J. Int., 159, 765-775.
813 Bartolini, C., 2003. When did the Northern Apennine become a mountain chain? Quatern. Int., 101-102, 75-
814 80.
815 Becker, T.W., Faccenna, C.,, O'Connel, R.J., 1999. The development of slabs in the upper mantle: Insights
816 from numerical and laboratory experiments. J. Geophys. Res., Solid Earth, 104, 15,207-15,226, doi:
817 10.1029/ 1999JB900140
818 Ben Avraham, Z., Boccaletti, M., Cello, G., Grasso, M., Lentini, F., Torelli, L., Tortorici, L., 1990. Principali
819 domini strutturali originatisi dalla collisione neogenico-quaternaria nel mediterraneo
820 centrale.Mem.Soc.Geol. It., 45, 453-462.
821 Benaouali-Mebarek, N., Frizon de Lamotte, D., Roca, E., Bracene, R., Jean-Luc Faure, J.L., Sassi, W.,
822 Roure, F., 2006. Post-Cretaceous kinematics of the Atlas and Tell systems in central Algeria: Early
823 foreland folding and subduction-related deformation. Comptes Rendus Geoscience, 338, 1–2, 115-
824 125.

825 Blumetti, A.M., Guerrieri, L., Vittori, E., 2013. The primary role of the Paganica-San Demetrio fault system
826 in the seismic landscape of the Middle Aterno Valley basin (Central Apennines). Quatern. Int., 288,
827 183-194. doi:10.1016/j.quaint.2012.04.040
828 Boccaletti, M., Cello, G., Tortorici, L., 1987. Transtensional tectonics in the Sicily Channel. J. Struct. Geol.,
829 9, 869-876.
830 Boccaletti, M., Bonini, M., Moratti, G., Sani, F., 1999. Compressive Neogene-Quaternary tectonics in the
831 hinterland area of the Northern Apennines. In: Boccaletti, M., Dahmani, M. (Eds.), Neogene
832 sedimentation and tectonics in the Western Mediterranean. J. Petrol. Geol., 22, 37-60.
833 Boccaletti, M., Corti, G., Martelli, L. 2011. Recent and active tectonics of the external zone of the Northern
834 Apennines (Italy). Geologische Rundschau, 100, 1331–1348, https://doi.org/10.1007/s00531-010-
835 0545-y
836 Boncio, P., Lavecchia, G., 2000. A structural model for active extension in Central Italy. J. Geodyn., 29,
837 233-244.
838 Boncio, P., Bracone, V. 2009. Active stress from earthquake focal mechanisms along the Padan–Adriatic
839 side of the Northern Apennines (Italy), with considerations on stress magnitudes and pore-fluid
840 pressures. Tectonophysics, 476. 180-194, https://doi.org/10.1016/j.tecto.2008.09.018.
841 Bonini, M., Sani, F., Stucchi, E.M., Moratti, G., Benvenuti, M., Menanno, G., Tanini, C.,2014. Late Miocene
842 shortening of the Northern Apennines back-arc. J. Geodyn.74, 1–31.
843 Bossio, A., Costantini, A., Lazzarotto, A., Liotta, D., Mazzanti, R., Mazzei, R.,Salvatorini, G.F., Sandrelli,
844 F., 1993. Rassegna delle conoscenze sulla stratigrafiadel Neoautoctono toscano. Mem. Soc. Geol. Ital.
845 49, 17–98.
846 Bossio, A., Costantini, A., Foresi, L.M., Lazzarotto, A., Mazzanti, R., Mazzei, R., Pascucci, V., Salvatorini,
847 G., Sandrelli, F., Terzuoli, A., 1998. Neogene–Quaternary sedimentary evolution in the western side
848 of the Northern Apennines (Italy). Mem. Soc. Geol. Ital., 52, 513–525.
849 Boutelier, D., Gagnon, L., Johnston, S., Cruden, A., 2018. In press. Buckling of orogens: insights from
850 analogue modelling. J. Struct. Geol., corrected proof online 27 February 2018.
851 https://doi.org/10.1016/j.jsg.2018.02.005

852 Brancolini, G., Civile, D., Donda, F., Tosi., L., Zecchin, M., Volpi, V., Rossi, G., Sandron, D., Ferrante,
853 G.M., Forlin, E., 2019. New insights into the Adria plate geodynamics from the northern Adriatic
854 perspective. Mar. Pet. Geo., 109, 687-697. https://doi.org/10.1016/j.marpetgeo.2019.06.049
855 Bressan, G., Snidarcig, A., Venturini, C., 1998. Present state of tectonic stress of the Friuli area (eastern
856 Southern Alps). Tectonophysics, 292, 211-227.
857 Brogi, A., Fidolini, F., Liotta, D., 2013. Tectonic and sedimentary evolution of theUpper Valdarno Basin:
858 new insights from the lacustrine S. Barbara Basin. Ital. J.Geosci. 132, 81–97.
859 Brozzetti, F. 2011. The Campania–Lucania Extensional Fault System, southern Italy: a suggestion for a
860 uniform model of active extension in the Italian Apennines. Tectonics, 30, TC5009.
861 https//doi.org/10.1029/2010TC002794
862 Buiter, S. J. H., Govers R., Wortel, M.J.R., 2001. A modelling study of vertical surface displacements at
863 convergent plate margins. Geophys. J. Int.., 147, 415-427.
864 Calais, E., Demets, D., Nocquet, J.M., 2003. Evidence for a post-3.16 Ma change in Nubia-Eurasia- North
865 America plate motions? Earth Planet. Sci. Lett., 216, 81-92.
866 Calamita, F., Cello, G., Deiana, G., Paltrinieri, W., 1994. Structural styles, cronology-rates of deformation
867 and time space relationships in the Umbria-Marche thrust system (Central Apennines, Italy).
868 Tectonics, 13, 873-881.
869 Calamita, F., Scisciani, V., Montefalcone, R., Paltrinieri, W., Pizzi A., 2002. L'ereditarietà del paleomargine
870 dell'Adria nella geometria del sistema orogenico centro-appenninico: l'area abruzzese esterna. Mem.
871 Soc. Geol. Ital., 57, 355-368.
872 Capitanio, F.A. Stegman, D.R., Moresi L.N., Sharples, W., 2010. Upper plate controls on deep subduction,
873 trench migrations and deformations at convergent margins. Tectonophysics, 483, 80-92.
874 doi:10.1016/j.tecto.2009.08.020
875 Carmignani, L., Decandia, F.A., Fantozzi, P.L., Lazzarotto, A., Liotta, D., Meccheri, M. 1994. Tertiary
876 extensional tectonics in Tuscany (northern Apennines, Italy). Tectonophysics, 238, 295–315.
877 Carmignani, L., Conti, P., Cornamusini, G., Meccheri, M., 2004. The internal northern Apennines, the
878 northern Tyrrhenian sea and the Sardinia-Corsica block, in: Crescenti, V., D'Offizi, S., Merlino S.,

879 Sacchi, L. (Eds.) Geology of Italy: Spec. Vol. Ital. Geol. Soc. for the IGC 32 Florence, Società
880 Geologica Italiana, Roma, 59-77.
881 Carminati E., Doglioni , C., Gelabert, B., Panza, G.F.,Raykova, R.B., Roca, E., Sabat, F., Scrocca, D., 2012.
882 Evolution of theWestern Mediterranean, in: Roberts, D.G., Bally, A.W. (Eds.) Regional Geology and
883 Tectonics: Phanerozoic Passive Margins, Cratonic Basins and Global Tectonic Maps. 437-470.
884 Elsevier, Amsterdam. DOI:10.1016/B978-0-444-56357-6.00011-1
885 Castellarin, A., Cantelli, L., 2000. Neo-Alpine Evolution of the Southern Eastern Alps. J. Geodyn., 30, 251-
886 274. http://dx.doi.org/10.1016/S0264-3707(99)00036-8.
887 Castellarin, A., TRANSALP Working Group, 20 04. Structural synthesis of the Eastern Alps: a collisional
888 orogenic chain. In: Crescenti, V., D'Offizi, S., Merlino S., Sacchi, L. (Eds.) Geology of Italy: Spec.
889 Vol. Ital. Geol. Soc for the IGC 32 Florence, Società Geologica Italiana, Roma, 3-13.
890 Catalano, R., Infuso, S., Sulli, A., 1994. The submerged alpidic chain from southern Sardinia shelf to the
891 Pelagian rifting: tectonic history. Boll. Geof. Teor. Appl., 36, 139-158.
892 Catalano, R., Di Stefano, P., Sulli, A., Vitale, F.P., 1996. Paleogeography and structure of the central
893 Mediterranean: Sicily and its offshore area. Tectonophysics, 260, 291-323.
894 Catalano, S., Monaco, C., Tortorici, L., 2004. Neogene-Quaternary tectonic evolution of the Southern
895 Apennines. Tectonics, 23, TC2003, doi:10.1029/2003TC001512
896 Catalano, S., De Guidi, G., Lanzafame, G., Monaco, C., Tortorici, L., 2009. Late Quaternary deformation on
897 the island on Pantelleria: new constraints for the recent tectonic evolution of the Sicily Channel Rift
898 (southern Italy). J.Geodyn. 48, 75–82. doi: 10.1016/j.jog.2009.06.005.
899 Cello, G., 1987. Structure and deformation processes in the Strait of Sicily "rift zone". Tectonophysics, 141,
900 237-247.
901 Cello, G., Mazzoli, S., 1999. Apennine tectonics in southern Italy: a review. J. Geodyn., 27, 191-211.
902 Cello, G., Mazzoli, S., Tondi, E., 1998. The crustal fault structure responsible for the 1703 earthquake
903 sequence of central Italy. J. Geodyn., 26, 443-460.
904 Cello, G., Deiana, G., Ferreli, L., Marchigiani, L., Maschio, L., Mazzoli, S., Michetti, A., Serva, L., Tondi,
905 E., Vittori, E., 2000. Geological constraints for earthquake faulting analysis in the Colfiorito area. J.
906 Seismol., 4, 357-364.

907 Cello, G., Tondi, E., Micarelli, L., Mattioni, L., 2003. Active tectonics and earthquake sources in the
908 epicentral area of the 1857 Basilicata earthquake (southern Italy). J. Geodyn., 36, 37-50.
909 Cenni, N., Mantovani,E., Baldi, P., Viti, M., 2012. Present kinematics of Central and Northern Italy from
910 continuous GPS measurements. J. Geodyn., 58, 62-72, doi:10.1016/j.jog.2012.02.004.
911 Cenni, N., Viti, M., Baldi, P., Mantovani, E., Bacchetti, M., Vannucchi, A., 2013. Present Vertical
912 Movements in Central and Northern Italy from GPS Data: Possible Role of Natural and
913 Anthropogenic Causes. J. Geodyn., 71, 74-85. http://dx.doi.org/10.1016/j.jog.2013.07.004
914 Ceriani, S., Schmid, S.M., 2004. From N-S collision to WNW-directed post-collisional thrusting and folding:
915 structural study of the Frontal Penninic Units in Savoie (Western Alps, France). Eclogae Geologicae
916 Helvetiae, 97, 47–369.
917 Cerrina Feroni, A., Leoni, L., Martelli, L., Martinelli, P. Ottria, G., Sarti G., 2001. The Romagna Apennines,
918 Italy: an eroded duplex. Geological Journal, 36, 39-54.
919 Cerrina Feroni, A., Ottria, G., Ellero, A., 2004. The northern Apennine, Italy: geological structure and
920 transpressive evolution, in: Crescenti, V., D'Offizi, S., Merlino S., Sacchi, L. (Eds.) Geology of Italy:
921 Special Volume of the Italian Geological Society for the IGC 32 Florence, Società Geologica Italiana,
922 Roma, 15-32.
923 Chiarabba, C., Jovane, L., Di Stefano, R., 2005. A new view of Italian seismicity using 20 years of
924 instrumental recordings. Tectonophysics, 395, 251-268.
925 Chicco, J.M., Pierantoni, P.P., Costa, M., Invernizzi, C., 2019. Plio-Quaternary tectonics and possible
926 implications for geothermal fluids in the Marche Region (Italy). Tectonophysics, 755, 21–34.
927 https://doi.org/10.1016/j.tecto.2019.02.005
928 Cinque, A., Ascione, A., Chiazzo, C., 2000. Distribuzione spaziotemporale e caratterizzazione della
929 fagliazione quaternaria in Appennino meridionale, in: Galadini, F., Meletti, C., Rebez , A. (Eds.),
930 Ricerche del GNDT nel campo della pericolosità sismica (1996-1999), Special Publication of CNR-
931 GNDT, Gruppo Nazionale per la Difesa dai Terremoti, Roma, 203-218.
932 Civile, D., Lodolo, E., Accettella, D.,Geletti, R., Ben-Avraham, Z., Deponte, M., Facchin, L., Ramella, R.,
933 Romeo, R., 2010. The Pantelleria graben (Sicily Channel, Central Mediterranean): An example of
934 intraplate 'passive' rift. Tectonophysics, 490, 173-183, doi:10.1016/j.tecto.2010.05.008

935 Civile, D., Lodolo, E., Alp, H., Ben-Avraham, Z., Cova, A., Baradello, L., Accettella, D., Burca,M.,
936 Centonze, J., 2014. Seismic stratigraphy and structural setting of the Adventure Plateau (Sicily
937 Channel). Mar. Geophys. Res. 35, 37–53.
938 Civile, D., Lodolo, E., Accaino, F.,Geletti, R., Schiattarella, M., Giustiniani, M., Fedorik, J., Zecchin, M.,
939 Zampa, L., 2018. Capo Granitola-Sciacca Fault Zone (Sicilian Channel, Central Mediterranean):
940 Structure vs magmatism. Mar. Pet. Geo., 96, 627-644, https://
941 doi.org/10.1016/j.marpetgeo.2018.05.016
942 Cloos, M. 1993. Lithospheric buoyancy and collisional orogenesis: subduction of oceanic plateaus,
943 continental margins, island arcs, spreading ridges, and seamounts. GSA Bulletin, 105, 715–737.
944 https://doi.org/10.1130/0016-7606(1993)105<0715:LBACOS>2.3.CO;2
945 Copley, A., Avouac, J.-P., Royer, J.-Y., 2010. India-Asia collision and the Cenozoic slowdown of the Indian
946 plate: Implications for the forces driving plate motions. J. Geophys. Res., 115,
947 B03410.doi:10.1029/2009JB006634ù
948 Cornamusini, G., Foresi, L.M., Dall'Antonia, B., Bossio, A., Mazzei, R., Salvatorini, G., 2014. The
949 Miocene of Pianosa Island: key to understanding the opening of the Northern Tyrrhenian back-arc
950 basin (Central Mediterranean). J. Geol. Soc. London, 171, 801–819.
951 http://dx.doi.org/10.1144/jgs2013-112
952 Costa, M. 2003. The buried, Apenninic arcs of the Po Plain and northern Adriatic Sea (Italy): a new model.
953 Boll. Soc. Geol. Ital., 122, 3-23.
954 Coward, M.P., De Donatis, M., Mazzoli, S., Paltrinieri, W., Wezel, F.C., 1999. Frontal part of the northern
955 Apennines fold and thrust belt in the Romagna-Marche area (Italy): shallow and deep structural styles.
956 Tectonics, 18, 559-574.
957 Davy, P., Hansen, A., Bonne,t E., Zhang, S.Z., 1995. Localisation and fault growth in layered brittle-ductile
958 systems: implications for deformations of the continental lithosphere. J. Geophys. Res., 100, 6281-
959 6294.
960 Del Ben, A., 1993. Calabrian Arc tectonics from seismic exploration. Boll. Geof. Teor. Appl., 139, 339-347.

961 Del Ben, A, Barnaba, C., Toboga, A., 2008. Strike-slip systems as the main tectonic features in the Plio-
962 Quaternary kinematics of the Calabrian Arc. Mar. Geophys. Res., 29, 1-12. doi:101007/s11001-007-
963 9041-6

964 Dercourt, J., Zonenshain, L.P., Ricou, L.E., Kazmin, V.G., Le Pichon, X., Knipper, A.L., Grandjacquet C.,
965 Sborshchikov, I.M., Geyssant, J., Lepvrier, C., Pechersky, D.H., Boulin, J., Sibuet, J.C., Savostin,
966 L.A., Sorokhtin, O., Westphal, M., Bazhenov, M.L., Lauer, J.P., Biju-Duval, B., 1986. Geological
967 evolution of the Tethys Belt from the Atlantic to the Pamirs since the Lias, in: Aubouin, J., Le Pichon,
968 X., Monin, A.S. (Eds.), Evolution of the Tethys. Tectonophysics, 123, 241–315.

969 Di Domenica, A., Turtù, A., Satolli, S., Calamita, F., 2012. Relationships between thrusts and normal faults
970 in curved belts: New insight in the inversion tectonics of the Central-Northern Apennines (Italy). J.
971 Struct. Geo., 42, 104-117. http://dx.doi.org/10.1016/j.jsg.2012.06.008

972 Driehaus, L., Nalpas, T., Cobbold, P.R., Gelabert, B. , Sàbat F., 2013. Effects of margin-parallel shortening
973 and density contrasts on back-arc extension during subduction: Experimental insights and possible
974 application to Anatolia. Tectonophysics, 608, 288-302. http://dx.doi.org/10.1016/j.tecto.2013.09.028

975 Elter, P., Giglia, G., Tongiorgi, M., Trevisan L., 1975. Tensional and compressional areas in the recent
976 (Tortonian to Present) evolution of the Northern Apennine. Boll.Geof.Teor.Appl. 65, 3-18.

977 Elter, F.M., Elter, P., Eva, C., Eva, E., Kraus, R.K., Padovano, M., Solarino, S. 2012. An alternative model
978 for the recent evolution of Northern–Central Apennines (Italy). J. Geodyn., 54, 55-63.
979 https://doi.org/10.1016/j.jog.2011.11.001

980 Etheve, N., Frizon de Lamotte, D., Mohn, G, Martos, R., Roca E.,, Christian Blanpied, C., 2016. Extensional
981 vs contractional Cenozoic deformation in Ibiza (Balearic Promontory, Spain): Integration in the West
982 Mediterranean back-arc setting. Tectonophysics, 682, 35–55

983 Faccenna, C., Davy, P., Brun, J.P., Funiciello, R., Giardini, D., Mattei, M., Nalpas, T., 1996. The dynamics
984 of back-arc extension: an experimental approach to the opening of the Tyrrhenian Sea. Geophys. J.
985 Int., 126, 781-795.

986 Faccenna, C., Giardini, D., Davy, P., Argentieri, A., 1999. Initiation of subduction at Atlantic-type margins:
987 insights from laboratory experiments. J. Geophys. Res., 104, 2749-2766.

988 Faccenna, C., Funiciello, F., Giardini, D., Lucente P., 2001, Episodic back-arc extension during restricted
989 mantle convection in the central Mediterranean. Earth Planet. Sci. Lett., 187, 105-116.
990 Faccenna, C., Piromallo, C., Crespo-Blanc, A., Jolivet L., 2004. Lateral slab deformation and the origin of
991 the western Mediterranean arcs. Tectonics, 23, TC1012. doi:10.1029/2002TC001488
992 Faccenna, C., Funiciello, F., Civetta, L., D'Antonio, M., Moroni, M., Piromallo, C., 2007. Slab disruption,
993 mantle circulation, and the opening of the Tyrrhenian basins, in: Beccaluva, L., G. Bianchini, G.,
994 Wilson, M. (Eds.), Cenozoic Volcanism in the Mediterranean Area, Geol. Soc. Am. Spec. Pap., 41,
995 153-169, AGU, Washington, D. C.. doi:10.1130/2007.2418(08)
996 Faccenna, C., Becker, T.W., Auer, L., Billi, A., Boschi, L., Brun, J.P., Capitanio, F.A., Funiciello, F.,
997 Horvàth, F., Jolivet, L., Piromallo, C., Royden, L., Rossetti, F., Serpelloni, E., 2014a. Mantle
998 dynamics in the Mediterranean, Rev. Geophys., 52, 283–332, doi:10.1002/2013RG000444.
999 Faccenna, C., Becker, T.W., Miller, M.S., Serpelloni, E., Willett, S.D. 2014b. Isostasy, dynamic topography,
1000 and the elevation of the Apennines of Italy. Earth Planet. Sci. Lett., 407, 163-174,
1001 https://doi.org/10.1016/j.epsl.2014.09.027
1002 Favaro, S., Handy, M. R., Scharf, A.,, Schuster, R., 2017. Changing patterns of exhumation and denudation
1003 in front of an advancing crustal indenter, Tauern Window (Eastern Alps). Tectonics, 36, 1053–1071.
1004 https://doi.org/10.1002/2016TC004448
1005 Fedorik, J., Toscani, G., Lodolo, E., Civile, D., Bonini, L., Seno, S., 2018. Structural analysis and Miocene-
1006 to-Present tectonic evolution of alithospheric-scale, transcurrent lineament: The Sciacca Fault
1007 (SicilianChannel, Central Mediterranean Sea). Tectonophysics, 722, 342-355.
1008 Fellin, M.G., Martin, S., Massironi, M., 2002. Polyphase tertiary fault kinematics and Quaternary
1009 reactivation in the central-eastern Alps (western Trentino). J. Geodyn., 34, 31–46.
1010 Ferranti, L., Santoro, E., Mazzella, M.E., Monaco, C., Morelli, D. 2009. Active transpression in the northern
1011 Calabria Apennines, southern Italy. Tectonophysics, 476, 226-251,
1012 https://doi.org/10.106/j.tecto.2008.11.010
1013 Ferranti, L., Burrato, P., Pepe, F., Santoro, E., Mazzella, M.E., Morelli, D., Passaro, S., Vannucci., G., 2014.
1014 An active oblique contractional belt at the transition between the Southern Apennines and Calabrian

Arc: The Amendolara Ridge, Ionian Sea, Italy. Tectonics, 33, 2169–2194, https://doi.org/10.1002/2014TC003624

Finetti, I.R., 2005a. Geodynamic Evolution of the Mediterranean Region from the Permo-Triassic Ionian Opening to the Present, Constrained by New Lithospheric CROP Seismic Data, in: Finetti, I.R. (Ed.), CROP PROJECT: Deep Seismic Exploration of the Central Mediterranean and Italy. Elsevier, Amsterdam, Chapter 34, 767-776.

Finetti, I.R., 2005b. Crustal Tectono-Stratigraphic Sections across the Western and Eastern Alps from ECORS-CROP and Transalp Seismic Data, in: Finetti, I.R. (Ed.), CROP PROJECT: Deep Seismic Exploration of the Central Mediterranean and Italy, Elsevier, Chapter 7, Elsevier, 109-118.

Finetti, I.R. 2006. Basic regional crustal setting and superimposed local pluton-intrusion-related tectonics in the Larderello-M. Amiata geothermal province, from integrated CROP seismic data. Boll. Soc. Geol. Ital., 125, 117-146.

Finetti, I.R., Del Ben, A., 1986. Geophysical study of the Tyrrhenian opening. Boll. Geof. Teor. Appl., 110, 75-156.

Finetti, I.R., Del Ben, A., 2005a. Ionian Tethys Lithosphere Roll-Back Sinking and Back-Arc Tyrrhenian Opening from New CROP Seismic Data, in: Finetti, I.R. (Ed.) CROP PROJECT: Deep Seismic Exploration of the Central Mediterranean and Italy, Elsevier, Chapter 21, 483-504.

Finetti, I.R., Del Ben, A., 2005c. Crustal tectono-stratigraphy of the Ionian Sea from new integrated CROP seismic data, in: Finetti I.R.(Ed.), CROP PROJECT: Deep Seismic Exploration of the Central Mediterranean and Italy, Elsevier, Chapter 19, 447- 470.

Finetti, I.R., Del Ben, A., 2005b. Crustal Tectono-Stratigraphic Setting of the Pelagian Foreland from New CROP Seismic Data, in: Finetti, I.R. (Ed.), CROP PROJECT: Deep Seismic Exploration of the Central Mediterranean and Italy, Elsevier, Chapter 26, 581-596.

Finetti, I., Lentini, F., Carbone, S., Catalano, S., Del Ben, A., 1996. Il sistema Appenninico Meridionale-Arco Calabro-Sicilia nel Mediterraneo Centrale: studio geofisico-geologico. Boll. Soc. Geol. Ital., 115, 529-559.

1041 Finetti, I.R., Boccaletti, M., Bovini, M., Del Ben, A., Geletti, R., Pipan, M., Sani, F., 2001. Crustal section
1042 based on CROP seismic data across the North Tyrrhenian-Northern Apennines-Adriatic Sea.
1043 Tectonophysics, 343, 135-163.
1044 Finetti, I.R., Boccaletti, M., Bonini, M., Del Ben, A., Pipan, M., Prizzon, A., Sani, F., 2005a. Lithospheric
1045 Tectono-Stratigraphic Setting of the Ligurian Sea-Northern Apennines-Adriatic Foreland from
1046 Integrated CROP Seismic Data, in: Finetti I.R. (Ed.), CROP PROJECT: Deep Seismic Exploration of
1047 the Central Mediterranean and Italy, Elsevier, Chapter 8, 119-158.
1048 Finetti, I.R., Lentini, F., Carbone, S., Del Ben, A., Di Stefano, A., Guarnieri, P., Pipan, M., Prizzon, A.,
1049 2005b. Crusta1 Tectono-Stratigraphy and Geodynamics of the Southern Apennines from CROP and
1050 other Integrated Geophysical-Geological Data, in: Finetti, I.R. (Ed.), CROP PROJECT: Deep Seismic
1051 Exploration of the Central Mediterranean and Italy Elsevier, Chapter 12, 225-262.
1052 Finetti, I.R., Lentini, F., Carbone, S., Del Ben, A., Di Stefano, A., Forlin, E., Guarnieri, P., Pipan, M.,
1053 Prizzon, A., 2005c. Geological Outline of Sicily and Lithospheric Tectono-Dynamics of its Tyrrhenian
1054 Margin from New CROP Seismic Data, in: Finetti, I.R. (Ed.), CROP PROJECT: Deep Seismic
1055 Exploration of the Central Mediterranean and Italy, Elsevier, Chapter 15, 319-376.
1056 Frisch, W., Dunkl, I., Kuhlemann, J., 2000. Post-collisional orogen-parallel large-scale extension in the
1057 Eastern Alps. Tectonophysics, 327, 239–265. http://dx.doi.org/10.1016/S0040-1951(00)00204-3
1058 Frizon de Lamotte, D., Leturmy, P., Missenard, Y., Khomsi, S., Ruiz, G., Saddiqi, O., Guillocheau., F.,
1059 Michard, A., 2009. Mesozoic and Cenozoic vertical movements in the Atlas system (Algeria,
1060 Morocco, Tunisia): An overview. Tectonophysics, 475, 9–28.
1061 Funiciello, F., Morra, G., Regenauer-Lieb, K., Giardini, D. 2003a. Dynamics of retreating slabs: 1. Insights
1062 from two-dimensional numerical experiments. J. Geophys. Res., 108, 2206,
1063 doi:10.1029/2001JB000898.
1064 Funiciello, F., Faccenna, C., Giardini, D., Regenauer-Lieb, K. 2003b. Dynamics of retreating slabs: 2.
1065 Insights from three-dimensional laboratory experiments. J. Geophys. Res., 108, 2207,
1066 doi:10.1029/2001JB000896
1067 Funiciello, F., Faccenna, C., Giardini D. 2004. Role of lateral mantle flow in the evolution of subduction
1068 system: Insights from 3-D laboratory experiments. Geophys. J. Int., 157, 1393-1406.

1069 Funiciello, F., Moroni, M., Piromallo, C., Faccenna, C., Cenedese, A., Bui H.A. 2006. Mapping mantle flow
1070 during retreating subduction: Laboratory models analyzed by feature tracking. J. Geophys. Res., 111,
1071 B03402, doi:10.1029/2005JB003792
1072 FurlanI, S., Antonioli, F., Biolchi, S., Gambin, T., Gauci, R., Lo Presti, V., Anzidei, A., Devoto, S.,
1073 Palombo, M., Sulli, A., 2013. Holocene sea level change in Malta. Quaternary International, 288, 146-
1074 157. doi:10.1016/j.quaint.2012.02.038
1075 Galadini, F., 1999. Pleistocene change in the central Apennine fault kinematics, a key to decipher active
1076 tectonics in central Italy. Tectonics, 18, 877-894.
1077 Galadini, F., Messina, P., 2004. Early-middle Pleistocene eastward migration of the Abruzzi Apennine
1078 (central Italy) extensional domain. J. Geodyn., 37, 57-81.
1079 Galadini, F., Poli, M.E., Zanferrari, A., 2005. Seismogenic sources potentially responsible for earthquakes
1080 with M≥6 in the eastern southern Alps (Thiene-Udine sector, NE Italy). Geophys. J. Int., 161, 739-
1081 762, doi:10.1111/j.1365-246X.2005.02571.x.
1082 Gaspar-Escribano, J. M., Garcia-Castellanos, D., Roca, E., Cloetingh, S., 2004. Cenozoic vertical motions
1083 of the Catalan Coastal Ranges (NE Spain): The role of tectonics, isostasy, and surface transport.
1084 Tectonics, 23, tc1004, doi:10.1029/2003tc001511
1085 Gattacceca, J., Deino, A., Rizzo, R., Jones, D.S., Henry, B., Beaudoin, B., Vadeboin, F., 2007. Miocene
1086 rotation of Sardinia: new paleomagnetic and geochronological constraints and geodynamic
1087 implications. Earth Planet. Sci. Lett., 258, 359–377. http://dx.doi.org/10.1016/j.epsl.2007.02.003
1088 Ghielmi, M., Minervini, M., Nini, C., Rogledi, S., Rossi, M., 2013. Late Miocene–Middle Pleistocene
1089 sequences in the Po Plain–Northern Adriatic Sea (Italy): the stratigraphic record of modification
1090 phases affecting a complex foreland basin. Mar. Pet. Geo., 42, 50-81.
1091 http://dx.doi.org/10.1016/j.marpetgeo.2012.11.007
1092 Ghisetti, F., Vezzani, L., 1981. Contribution of structural analysis to understanding thegeodynamic evolution
1093 of the Calabrian Arc (southern Italy). J. Struct. Geo., 3, 371–381.
1094 Ghisetti, F.C., Gorman, A.R., Grasso, M., Vezzani, L., 2009. Imprint of foreland structure on the
1095 deformation of a thrust sheet: the Plio-Pleistocene Gela Nappe (southern Sicily, Italy). Tectonics 28
1096 (4). http://dx.doi.org/10.1029/2008TC002385. TC4015.

1097 Grasso, M., Pedley, H.M., 1985. The Pelagian Islands: a new geological interpretation from sedimentological
1098 and tectonic studies and its bearing on the evolution of the Central Mediterrranean Sea (Pelagian
1099 Block). Geologica Romana, 24, 13-34.
1100 Guillaume, B., Funiciello, F., Faccenna, C., Martinod, J., Olivetti, V., 2010. Spreading pulses of the
1101 Tyrrhenian Sea during the narrowing of the Calabrian slab. Geology 38, 819–822.
1102 Guarnieri, P., 2004. Structural evidence for deformation by block rotation in the context of transpressive
1103 tectonics, northwestern Sicily (Italy). J. Struct. Geol., 26, 207-219.
1104 Guarnieri, P., 2006. Plio-Quaternary segmentation of the south Tyrrhenian forearc basin International
1105 Journal of Earth Sciences (Geologische Rundschau), 95, 107-118, doi:101007/s00531-005-0005-2
1106 Gudmundsson, A., 2001. Fluid Overpressure and Flow in Fault Zones: Field Measurements and Models.
1107 Tectonophysics, 336, 183-197.
1108 Gueguen, E., Tavarnelli, E., Renda, P., Tramutoli, M., 2002. The geodynamics of the southern Tyrrhenian
1109 Sea margin as revealed by integrated geological, geophysical and geodetic data. Boll. Soc. Geol. It.,
1110 Vol. Spec., 1, 77-85.
1111 Gvirtzman, Z., Nur, A. , 2001. Residual topography, lithospheric structure and sunken slabs in the central
1112 Mediterranean. Earth Planet. Sci. Lett., 187, 117–130.
1113 Hall, R., 2018. The subduction initiation stage of the Wilson cycle. From: Wilson, R.W., Houseman, G.A.,
1114 Mccaffrey, K.J.W., Doré, A.G., Buiter, S.J.H. (eds) Fifty Years of the Wilson Cycle Concept in Plate
1115 Tectonics. Geol. Soc. London, Spec. Public., 470. https://doi.org/10.1144/SP470.3
1116 Hampel, A., Pfiffner, A., 2006. Relative importance of trenchward upper plate motion and friction along the
1117 plate interface for the topographic evolution of subductionrelated mountain belts. In: Buiter, S. J. H.,
1118 Schreurs, G. (Eds.) Analogue and Numerical Modelling of Crustal-Scale Processes. Geological
1119 Society London , Special Publication 253, 105-115.
1120 http://dx.doi.org/10.1144/GSL.SP.2006.253.01.05

1121 Handy, M. R., Ustaszewski, K., Kissling, E., 2015. Reconstructing the Alps–Carpathians–Dinarides as a key
1122 to understanding switches in subduction polarity, slab gaps and surface motion. Int. J. Earth Sci., 104,
1123 1–26. DOI 10.1007/s00531-014-1060-3
1124 Hassani, R., Jongmans, D., Chery, J., 1997. Study of plate deformation and stress in subduction processes
1125 using two dimensional numerical models. J. Geophys. Res., 102, 17951-17965.
1126 Hieke, W., Hirschleber, H.B., Deghani, G.A. 2003. The Ionian Abyssal Plain (central Mediterranean Sea):
1127 morphology, sub-bottom structures and geodynamic history – an inventory. Mar. Geophys. Res., 24,
1128 279–310, https://doi.org/10.1007/s11001-004-2173-z
1129 Hieke, W., Cita, M.B., Forcella, F., Muller, C. 2006. Geology of the Victor Hensen Seahill (Ionian Sea,
1130 eastern Mediterranean): insights from the study of cored sediment sequences. Boll. Soc. Geol. Ital.,
1131 125, 245–257.
1132 Hippolyte, J.C., Angelier, J., Barrier, E., 1994. A major geodynamic change revealed by Quaternary stress
1133 patterns in the southern Apennines (Italy). Tectonophysics, 230, 199-210.
1134 Hippolyte, J.C., Murovskaya, A., Volfman, Y., Yegorova T., Gintov, O., Kaymakci, N., Sangu, E., 2018.
1135 Age and geodynamic evolution of the Black Sea Basin: Tectonic evidences of rifting in Crimea. Mar.
1136 Pet. Geo., 93, 298-314. https://doi.org/10.1016/j.marpetgeo.2018.03.009
1137 Horváth, F., Musitza, B., Balázsb, A., Véghd, A., Uhrine, A., Nádorc, A., Koroknaia, B., Papd, N., Tótha,
1138 T., Wóruma, G., 2015. Evolution of the Pannonian basin and its geothermal resources. Geothermics,
1139 53, 328–352. https://doi.org/10.1016/j.geothermics.2014.07.009,
1140 Husson, L. 2006. Dynamic topography above retreating subduction zones. Geology, 34, 741-744,
1141 doi:10.1130/G22436.1
1142 ISIDe working group, 2016. Version 1.0, DOI: 10.13127/ISIDe.
1143 Jolivet, L., Faccenna, C., 2000. Mediterranean extension and the Africa-Eurasia collision. Tectonics, 19, 6,
1144 1095-1106.
1145 Kim, Y.S., Peacock, D.C.P., Sanderson, D.J., 2003. Mesoscale strike-slip faults and damage zones at
1146 Malsalform, Gozo Island, Malta. J. Struct. Geol., 25, 793-812.
1147 Lentini, F., Carbone, S., Catalano S., 1994. Main structural domains of the central Mediterranean region and
1148 their tectonic evolution. Boll. Geofis. Teor. Appl., 36, 103-125.

1149 Lentini, F., Carbone, S., Guarnieri, P., 2006. Collisional and postcollisional tectonics of the Apenninic-
1150 Maghrebian orogen (southern Italy), in: Dilek, Y., Pavlides, S. (Eds.) Postcollisional tectonics and
1151 magmatism in the Mediterranean region and Asia Geol. Soc. Am. Spec. Pap., 409, p. 57–81, doi:
1152 10.1130/2006.2409(04)
1153 Liotta, D., Brogi, A., Meccheri, M., Dini, A., Bianco, C., Ruggieri, G., 2015. Coexistenceof low-angle
1154 normal and high-angle strike-to oblique-slip faults during Late Miocene mineralization in eastern Elba
1155 Island (Italy). Tectonophysics, 660, 17-34.
1156 Lodolo E., Civile D., Zanolla C., Geletti R., 2012. Magnetic signature of the Sicily channel volcanism. Mar.
1157 Gephys. Res., 33, 33-44.
1158 Lucente, F.P., Chiarabba, C., Cimini, G., Giardini, D., 1999. Tomographic constraints on the geodynamic
1159 evolution of the Italian region, J. Geophys. Res.,104, 20307–20327.
1160 Lustrino, M., Wilson, M., 2007. The Circum-Mediterranean Anorogenic Cenozoic Igneous Province. Earth-
1161 Sci. Rev., 81, 1-65. doi:10.1016/j.earscirev.2006.09.002
1162 Lustrino, M., Morra, V., Fedele, L., Franciosi. L., 2009. Beginning of the Apennine subduction system in
1163 central western Mediterranean: Constraints from Cenozoic ‘‘orogenic’’ magmatic activity of Sardinia,
1164 Italy. Tectonics, 28, TC5016, doi:10.1029/2008TC002419
1165 Maillard, A., Mauffret, A., 1999. Crustal structure and riftogenesis of the Valencia through (North-Western
1166 Mediterranean Sea). Basin Res., 11, 357–379.
1167 Makris, J., Egloff, F., Nicolich, R., Rihma, R., 1999. Crustal structure from the Ligurian Sea to the Northern
1168 Apennines-a wide angle seismic transect. Tectonophysics, 301, 305-319.
1169 Malinverno, A., Ryan, W.B.F., 1986. Extension in the Tyrrhenian sea and shortening in the Apennines as
1170 result of arc migration driven by sinking of the lithosphere. Tectonics, 5, 227-245.
1171 Malusà, M.G., Danišík, M., Kuhlemann, J., 2016.Tracking the Adriatic-slab travel beneath the Tethyan
1172 margin of Corsica–Sardinia by low-temperature thermochronometry. Gondwana Res., 31, 135–149.
1173 Mantovani, E., 2005. Evolutionary reconstruction of the Mediterranean region: extrusion tectonics driven by
1174 plate convergence, in: Finetti I.R. (Ed.) CROP PROJECT: Deep Seismic Exploration of the Central
1175 Mediterranean and Italy, Elsevier, Chapter 32, 705-746.

1176 Mantovani, E., Viti, M., Albarello, D., Tamburelli, C., Babbucci, D., Cenni, N., 2000. Role of kinematically
1177 induced forces in Mediterranean tectonics: insights from numerical modelling. J. Geodyn., 30, 287–
1178 320.
1179 Mantovani, E., Viti, M., Babbucci, D., Tamburelli, C., Albarello, D. 2001a. Back arc extension: which
1180 driving mechanism? J. Virtual Explorer , 3, 17-44.
1181 Mantovani, E., Cenni, N., Albarello, D., Viti, M., Babbucci, D., Tamburelli, C., D'Onza, F., 2001b.
1182 Numerical simulation of the observed strain field in the central-eastern Mediterranean region. J.
1183 Geodyn., 3, 519-556.
1184 Mantovani, E., Albarello, D., Babbucci, D., Tamburelli, C., Viti, M. 2002. Trench-Arc-BackArc systems in
1185 the Mediterranean area: examples of extrusion tectonics. J. Virtual Explorer , 8, 125-141.
1186 Mantovani, E., Viti, M., Babbucci, D., Tamburelli, C., Albarello, D., 2006. Geodynamic connection between
1187 the indentation of Arabia and the Neogene tectonics of the central-eastern Mediterranean region, in:
1188 Dilek, Y., Pavlides, S. (Eds.) Post-Collisional Tectonics and Magmatism in the Mediterranean region
1189 and Asia. Geol. Soc. Am. Spec. Pap., 490, 15-49. doi: 10.1130/2006.2409(02)
1190 Mantovani, E., Viti, M., Babbucci, D., Albarello, D., 2007a. Nubia-Eurasia kinematics: an alternative
1191 interpretation from Mediterranean and North Atlantic evidence. Ann. Geophys., 50, 311-336.
1192 Mantovani E., Viti M., Babbucci D.,, Tamburelli C., 2007b. Major evidence on the driving mechanism of the
1193 Tyrrhenian-Apennines arc-trench-back-arc system from CROP seismic data. Boll. Soc. Geol. Ital.,
1194 126, 459-471
1195 Mantovani, E., Babbucci, D., Tamburelli, C., Viti, M., 2009. A review on the driving mechanism of the
1196 Tyrrhenian–Apennines system: implications for the present seismotectonic setting in the Central-
1197 Northern Apennines. Tectonophysics, 476, 22-40. doi:10.1016/j.tecto.2008.10.032.
1198 Mantovani, E., Viti, M., Babbucci, D., Albarello, D., Cenni, N., Vannucchi, A., 2010. Long-term earthquake
1199 triggering in the southern and Northern Apennines. J. Seismol., 14, 53-65. doi:10.1007/s10950-008-
1200 9141-z.
1201 Mantovani ,E., Viti, M., Babbucci, D., Cenni, N., Tamburelli, C., Vannucchi, A., 2012. Middle term
1202 prediction of earthquakes in Italy: some remarks on empirical and deterministic approaches. Boll.
1203 Geof. Teor. Appl., 53, 89-111. doi: 10.4430/bgta00XX.

1204 Mantovani, E., Viti, M., Babbucci, D., Tamburelli, C., Cenni, N., Baglione, M., D'Intinosante, V., 2014.
1205 Generation of back-arc basins as side effect of shortening processes: Examples from the Central
1206 Mediterranean. Int. J. Geosciences, 5, 1062-1079. http://dx.doi.org/10.4236/ijg.2014.510091.
1207 Mantovani E., Viti M., Cenni N., Babbucci D., Tamburelli C., 2015a. Present velocity field in the Italian
1208 region by GPS data: geodynamic/tectonic implications. Int. J. Geosciences, 6, 1285-1316.
1209 http://dx.doi.org/10.4236/ijg.2015.612103.
1210 Mantovani E., Viti M., Babbucci D., Tamburelli C., Cenni N., Baglione M., D'Intinosante V. 2015b.
1211 Present tectonic setting and spatio-temporal distribution of seismicity in the Apennine belt. Int. J.
1212 Geosciences, 6, 429-454. http://dx.doi.org/10.4236/ijg.2015.64034.
1213 Mantovani, E., Viti, M., Babbucci, D., Tamburelli, C., Cenni, N., Baglione, M., D'Intinosante, V., 2016.
1214 Recognition of periAdriatic seismic zones most prone to next major earthquakes: insights from a
1215 deterministic approach, in: D'Amico, S. (Ed.) Earthquakes and Their Impact on Society. Springer
1216 Natural Hazard Series, Springer International Publishing, Switzerland, pp.43-80. DOI 10.1007/978-3-
1217 319-21753-6_2
1218 Mantovani, E., Viti, M., Babbucci, D., Tamburelli, C., Cenni, N., 2017. Possible Location of the Next Major
1219 Earthquakes in the Northern Apennines: Present Key Role of the Romagna-Marche-Umbria Wedge.
1220 IJG, 8, 1301-1314. https://doi.org/10.4236/ijg.2017.811075.
1221 Mantovani, E., Viti, M., Babbucci D., Tamburelli, C. , Cenni, N., 2019. How and why the present tectonic
1222 setting in the Apennine belt has developed. J. Geol. Soc. London, Published Online.
1223 https://doi.org/10.1144/jgs2018-175
1224 Martin, S., Bigazzi, G., Zattin, M., Viola, G., Balestrieri, M.L., 1998. Neogene kinematics of the Giudicarie
1225 fault (Central-Eastern Alps, Italy): new apatite fission-track data. Terra Nova, 10, 217–221.
1226 Martini, I.P., Sagri, M., 1993. Tectono-sedimentary characteristics of late Miocene-Quaternary extensional
1227 basins of the Northern Apennines, Italy. Earth Sci. Rev., 34, 197-233.
1228 Maschio, L., Ferranti, L., Burrato, P. 2005. Active extension in Val d'Agri area, southern Apennines, Italy:
1229 implications for the geometry of the seismogenic belt. Geophys. J. Int., 162, 591-609,
1230 https://doi.org/10.1111/j.1365-246X.2005.02597.x

1231 Mascle, J., Rehault, J.P., 1990. A revised seismic stratigraphy of the Tyrrhenian Sea: implications for the
1232 basin evolution, in: Mastens, K.A., Mascle, J. et al. (Eds.), Proceedings ODP Scientific Results,
1233 College Station, Tx (Ocean Drilling Program), 107, 617-636.
1234 Massari, F., Prosser, G., 2013. Late Cenozoic tectono-stratigraphic sequences of the Crotone Basin: insights
1235 on the geodynamic historyof the Calabrian arc and Tyrrhenian Sea. Basin Res., 25, 26–51,
1236 doi:10.1111/j.1365-2117.2012.00549.x
1237 Massironi, M., Zampieri, D., Caporali, A., 2006. Miocene to present major fault linkages through the
1238 Adriatic indenter and the Austroalpine-Penninic collisional wedge (Alps of NE Italy). Geol. Soc.
1239 London, Spec. Public., 262, 245-258, http://dx.doi.org/10.1144/GSL.SP.2006.262.01.15.
1240 Mauffret, A., Contrucci, I., Brunet, C., 1999. Structural evolution of the Northern Tyrrhenian Sea from new
1241 seismic data. Mar. Pet. Geol., 16, 381-407.
1242 Mercier, J., Sorel, D., Simeakis, K., 1987. Changes in the state of stress in the overriding plate of a
1243 subduction zone: the Aegean Arc from the Pliocene to the Present. Annales Tectonicae, 1, 20-39.
1244 Meulenkamp, J.E., Sissingh, W., 2003.Tertiary palaeogeography and tectonostratigraphic evolution of the
1245 Northern and Southern Peri-Tethys platforms and the intermediate domains of the African–Eurasian
1246 convergent plate boundary zone. Palaeogeogr. Palaeocl, 196, 209-22, https://doi.org/10.1016/S0031-
1247 0182(03)00319-5
1248 Meyer, C., Schellart W.P., 2013. Three-dimensional dynamic models of subducting plate-overriding plate-
1249 upper mantle interaction. J. Geophys. Res., Solid Earth, 118, 775-790, doi:10.1002/jgrb.50078
1250 Milia, A., Torrente, M.M., 2003. Late-Quaternary volcanism and transtensional tectonics in the Bay of
1251 Naples, Campanian continental margin, Italy. Mineralogy and Petrology, 79, 49-65,
1252 http://dx.doi.org/10.1007/s00710-003-0001-9
1253 Moeller, S., Grevemeyer, I., Ranero, C.R., Berndt, C., Klaeschen, D., Sallares, V., Zitellini, N., de Franco,
1254 R., 2013. Early-stage rifting of the northern Tyrrhenian Sea Basin: Results from a combined wide-
1255 angle and multichannel seismic study. Geochem. Geophy. Geosy., 14, 3032–3052,
1256 doi:10.1002/ggge.20180

1257 Molli G., 2008. Northern Apennine-Corsica Orogenic System: an Updated Overview, in: Siegesmund, S.,
1258 Fugensheu, B., Froitzheim, N. (Eds.) Tectonic aspects of the Alpine-Dinarides-Carpathians System.
1259 Geol. Soc. London. Spec. Publ., 298, 413–442.
1260 Monaco, C., Tortorici L., 2000. Active faulting in the Calabrian arc and eastern Sicily. J. Geodyn., 29, 407-
1261 424.
1262 Mosca, P., Polino, R., Rogledi, S., Rossi, M., 2010. New data for the kinematic interpretation of the Alps-
1263 Apennines junction (Northwestern Italy). Int. J. Earth Sci., 99, 833-849. doi:10.1007/s00531-009-
1264 0428-2.
1265 Nicolosi, I., Speranza, F., Chiappini, M., 2006. Ultrafast oceanic spreading of the Marsili basin, southern
1266 Tyrrhenian Sea. Evidence from magnetic anomaly analysis. Geology 34, 717–720.
1267 Oggiano, G., Funedda, A., Carmignani, L., Pasci, S., 2009. The Sardinia-Corsica microplate and its role in
1268 the Northern Apennine geodynamics: new insights from the Tertiary intraplate strike-slip tectonics of
1269 Sardinia. Boll. Soc. Geol. Ital.. 128, 527-539, DOI: 10.3301/IJG.2009.128.2.527
1270 Pantosti, D., Valensise, G., 1990. Faulting mechanism and complexity of the November 23, 1980, Campania-
1271 Lucania earthquake inferred from surface observations. J. Geophys. Res., 95, 319-341.
1272 Parotto, M., Praturlon, A., 2004. The southern Apennini arc, in: Crescenti, V., D'Offizi, S., Merlino S.,
1273 Sacchi, L. (Eds.) Geology of Italy: Spec. Vol. Ital. Geol. Soc. for the IGC 32 Florence, Società
1274 Geologica Italiana, Roma, 33-58.
1275 Patacca, E., Scandone, P., 1989. Post-Tortonian mountain building in the Apenines. The role of the passive
1276 sinking of a relict lithospheric slab. In: Boriani, A., Bonafede, M., Piccardo, G.B., Vai, G.B. (Eds.),
1277 The Lithosphere in Italy. Advances in Earth Science Research, Accademia Naz.le dei Lincei, Roma,
1278 80, pp.157-176.
1279 Patacca, E., Scandone, P., 2004. The Plio-Pleistocene thrust belt-foredeep system in the southern Apennines
1280 and Sicily (Italy). In: Crescenti, V., D'Offizi, S., Merlino S., Sacchi, L. (Eds.) Geology of Italy: Spec.
1281 Vol. Ital. Geol. Soc.for the IGC 32 Florence, Società Geologica Italiana, Roma, 93-129.
1282 Patacca, E., Scandone, P., 2007. Geology of the Southern Apennines. In: Mazzotti, A., Patacca, E.,
1283 Scandone, P. (Eds.) CROP-04. Boll.Soc. Geo. Ital., Special Issue, 7, 75-119.

1284 Patacca, E., Sartori, R., Scandone, P., 1990. Tyrrhenian basin and Apenninic arcs: kinematic relations since
1285 Late Tortonian times. Mem. Soc. Geol. Ital., 45, 425-451.
1286 Peccerillo, A., 2003. Plio-quaternary magmatism in Italy. Episodes 26, 222–226.
1287 Peccerillo, A., 2005. Plio-Quaternary Volcanism in Italy. Petrology, Geochemistry, Geodynamics. Springer-
1288 Verlag,Berlin-Heidelberg, ISBN 978-3-540-25885-8.
1289 Pepe, F., Sulli, A., Bertotti, G., Catalano, R., 2005. Structural highs formation and their relationship to
1290 sedimentary basins in the north Sicily continental margin (southern Tyrrhenian Sea): Implications for
1291 the Drepano Thrust Front. Tectonophysics, 409, 1-18, doi:10.1016/j.tecto.2005.05.009
1292 Peresson, H., Decker, K., 1997. The Tertiary dynamics of the northem Eastem Alps (Austria): changing
1293 palaeostresses in a collisional plate boundary. Tectonophysics, 272, 125-157.
1294 Piccardi, L., Gaudemer, Y., Tapponnier, P., Boccaletti, M., 1999. Active oblique extension in the central
1295 Apennines (Italy): evidence from the Fucino region. Geophys. J. Int., 139, 499-530.
1296 Piccardi, L., Tondi, G., Cello, G., 2006. Geo-structural evidence for active oblique extension in South-
1297 Central Italy, in: Pinter, N., Grenerczy, G., Weber, J., Stein, S., Medak, D. (Eds), The Adria
1298 microplate: GPS Geodesy, Tectonics and Hazard. Springer, pp. 95-108.
1299 Piromallo, C., Morelli, A., 2003. P wave tomography of the mantle under the Alpine-Mediterranean area. J.
1300 Geophys. Res., 108 (B2), 2065, doi:10.1029/2002jb001757.
1301 Pizzi, A., 2003. Plio-Quaternary uplift rates in the outer zone of the central Apennines fold-and-thrust belt,
1302 Italy. Quatern. Int., 101-102, 229-237.
1303 Pizzi, A., Galadini, F., 2009. Pre-existing cross-structures and active fault segmentation in the northern-
1304 central Apennines (Italy). Tectonophysics , 476, 304–319. doi:10.1016/j.tecto.2009.03.018
1305 Placer, L., Vrabec, M., Celarc, B., 2010. The bases for understanding of the NW Dinarides and Istria
1306 Peninsula tectonics. Geologija 53 (1), 55–86.
1307 Pola, M., Ricciato, A., Fantoni, R., Fabbri, P., Zampieri, D. 2014. Architecture of the western margin of the
1308 north Adriatic foreland: The Schio–Vicenza Fault System. Ital. J. Geosci., 133, 223-234,
1309 https://doi.org/10.3301/IJG.2014.04
1310 Ratschbacher, L., Frisch, W., Linzer, H.G., Merle, O., 1991. Lateral Extrusion in the Eastern Alps, Part 2:
1311 Structural Analysis. Tectonics, 10, 257-271. http://dx.doi.org/10.1029/90TC02623.

1312 Réka, L., Harangi, S., Guillong, M., Bachmann, O., László Fodor, L., Buret, Y., Dunkl, I., Sliwinski, J., von
1313 Quadt, A., Peytcheva, I., Zimmerer, M., 2018. Early to Mid-Miocene syn extensional massive silicic
1314 volcanism in the Pannonian Basin (East-Central Europe): Eruption chronology, correlation potential
1315 and geodynamic implications. Earth Sci. Rev., 179, 1-19.
1316 Reuther, D., Ben Avraham, Z., Grasso, M., 1993. Origin and role of major strike-slip transfers during plate
1317 collision in the Central Mediterranean. Terra Nova, 5, 249-257.
1318 Robertson, A., Shallo, M., 2000. Mesozoic-Tertiary tectonic evolution of Albania in its regional eastern
1319 Mediterranean context. Tectonophysics, 316, 197-254.
1320 Robl, J., Stuwe, K., 2005. Continental collision with finite indenter strength: 2. European Eastern Alps.
1321 Tectonics, 24, TC4014, doi:10.1029/2004TC001741
1322 Roda-Boluda, D.C., Whittaker, A.C. 2017. Structural and geomorphic constraints on active normal faulting
1323 and landscape evolution in Calabria, Italy. J. Geol. Soc. London, 174, 701-720,
1324 https://doi.org/10.1144/jgs2016-097
1325 Rosenberg, CL., Kissling, E., 2013. 3D insight into Central Alpine Collision: lower plate or upper plate
1326 indentation? Geology , 41,1219–1222.
1327 Rovida A., Locati M., Camassi R., Lolli B., Gasperini P. Eds., 2016. CPTI15, the 2015 version of the
1328 Parametric Catalogue of Italian Earthquakes. Istituto Nazionale di Geofisica e Vulcanologia,
1329 doi:http://doi.org/10.6092/INGV.IT-CPTI15.
1330 Royden, L.H., 1993. Evolution of retreating subduction boundaries formed during continental collision.
1331 Tectonics, 12, 629-638.
1332 Royden, L.H., Burchfiel, B.C., 1989. Are systematic variations in thrust belt style related to plate boundary
1333 processes? (The western Alps versus the Carpathians). Tectonics, 8, 51-61.
1334 Royden, L., Horvath, F., Rumpler, J., 1983. Evolution of the Pannonian Basin system. Tectonics, 2, 63–90.
1335 Rusciadelli, G., 2005. Provenance of Upper Pliocene conglomerates of the Mutignano Formation (Abruzzo,
1336 Italy): implications for unraveling the uplift history of the Central Apennines external zones. Boll.
1337 Soc. Geol. Ital.,124, 645-659.

1338 Santangelo, N., Di Donato, V., Lebreton, V., Romano, P., Russo Ermolli, E., 2012. Palaeolandscapes of
1339 Southern Apennines during the late Early and the Middle Pleistocene. Quatern. Int., 267, 20-29.
1340 doi:10.1016/j.quaint.2011.02.036
1341 Sartori, R., 1990. The main results of ODP Leg 107 in the frame of Neogene to recent geology of
1342 perityrrhenian areas. In: Mastens, K.A., Mascle, J. et al. (Eds.), Proceedings ODP Scientific Results,
1343 College Station, Tx (Ocean Drilling Program), 107, 715–730.
1344 Sartori, R., 2005. Bedrock geology of the Tyrrhenian sea insights on Alpine paleogeography and magmatic
1345 evolution of the basin, in: Finetti, I.R. (Ed.), CROP PROJECT, Deep Seismic Exploration of the
1346 Central Mediterranean and Italy. Elsevier, 4, 69–80.
1347 Sartori, R., Carrara, G., Torelli, L., Zitellini, N., 2001. Neogene evolution of the southwestern Tyrrhenian
1348 Sea (Sardinia basin and western Bathyal plain). Mar. Geol., 175, 47–66.
1349 Sartori, R., Torelli, L., Zitellini, N., Carrara, G., Magaldi, M., Mussoni, P., 2004. Crustal features along a W-
1350 E Tyrrhenian transect from Sardinia to Campania margins (Central Mediterranean). Tectonophysics,
1351 383, 171-192.
1352 Savelli, C., 2002. Time–space distribution of magmatic activity in the western Mediterranean and peripheral
1353 orogens during the past 30 Ma (a stimulus to geodynamic considerations. J. Geodyn., 34, 99–126.
1354 Scafidi, D., Solarino, S., 2012. Can local earthquake tomography settle the matter about subduction in the
1355 Northern And Central Apennines? Response from a new high resolution P velocity and Vp/Vs ratio 3-
1356 D model. Tectonophysics, 554-557, 63-73. https://doi.org/10.1016/j.tecto.2012.06.007
1357 Scafidi, D., Solarino, S., Eva, C., 2009. P wave seismic velocity and Vp/Vs ratio beneath the Italian
1358 peninsula from local earthquake tomography. Tectonophysics , 465, 1-23.
1359 doi:10.1016/j.tecto.2008.07.013
1360 Schellart, W.P., 2005. Influence of the subducting plate velocity on the geometry of the slab and migration of
1361 the subduction hinge. Earth Planet. Sci. Lett., 231, 197-219. https://doi.org/10.1016/j.epsl.2004.12.019
1362 Schellart, W.P., 2010. Evolution of subduction zone curvature and its dependence on the trench velocity and
1363 the slab to upper mantle viscosity ratio. J. Geophys. Res., 115, B11406, doi:10.1029/2009JB006643

1364 Schellart, W.P., Moresi, L., 2013. A new driving mechanism for backarc extension and backarc shortening
1365 through slab sinking induced toroidal and poloidal mantle flow: Results from dynamic subduction
1366 models with an overriding plate. J. Geophys. Res., Solid Earth, 118, doi:10.1002/jgrb.50173
1367 Schellart, W.P., Freeman, J., Stegman, D.R., Moresi, L., May, D., 2007. Evolution and diversity of
1368 subduction zones controlled by slab width. Nature, 446, 308-311.
1369 Schellart, W.P., Stegman, D.R., Farrington, R.J., Moresi, L., 2011. Influence of lateral slab edge distance on
1370 plate velocity, trench velocity, and subduction partitioning. J. Geophys. Res., 116, B10408,
1371 doi:10.1029/2011JB008535
1372 Schettino, A., Turco, E., 2006. Plate kinematics of theWestern Mediterranean region during the Oligocene
1373 and Early Miocene .Geophys. J. Int. , 166, 1398–1423, doi: 10.1111/j.1365-246X.2006.02997.x
1374 Schiattarella, M., Di Leo, P., Beneduce, P., Giano, S.I., 2003. Quaternary uplift vs tectonic loading: a case
1375 study from the Lucanian Apennine, southern Italy. Quatern. Int., 101-102, 239-251.
1376 Schmid, S.M., Pfiffner, O.A., Froitzheim, N., Schonborn, G.E., 1996. Geophysical–geological transect and
1377 tectonic evolution of the Swiss-Italian Alps. Tectonics, 15, 1036–1064.
1378 Schmid, S.M., Fugenschuh, B., Schuster, R., 2004. Tectonic map and overall architecture of the Alpine
1379 orogen. Eclogae Geol. Helv., 97, 93-118.
1380 Scisciani, V., Calamita, F. 2009. Active intraplate deformation within Adria: Examples from the Adriatic
1381 region. Tectonophysics, 476, 57-72, https://doi. org/10.1016/j.tecto.2008.10.030
1382 Shemenda, A.I., 1993. Subduction of the lithosphere and back-arc dynamics: insights from physical
1383 modeling. J. Geophys. Res., 98, 167-185.
1384 Sorel, D., Bizon, G., Aliaj, S., Hasani, L., 1992. Calage stratigraphique de l'age et de la durée des phases
1385 compressives des Hellénides externes (Grèce nord-occidentale et Albanie) du Miocène à l'Actuel.
1386 Bull. Soc. Géol. France, 163, 447-454.
1387 Spakman, W., Wortel, R., 2004. A tomographic view on Western Mediterranean geodynamics, in: Cavazza,
1388 W., Roure, F.M., Spakman, W., Stampfli, G.M., Ziegler, P.A. (Eds.), The TRANSMED atlas: the
1389 Mediterranean region from Crust to Mantle, geological and geophysical framework of the
1390 Mediterranean and the surrounding areas. A publication of the Mediterraean consortium for the 32nd
1391 international geological congress. Springer, Berlin, Heidelberg, 31–52.

1392 Speranza, F., Villa, I.M., Sagnotti, L., Florindo, F., Cosentino, D., Cipollari, P., Mattei, M., 2002. Age of
1393 the Corsica–Sardinia rotation and Liguro–Provencal Basin spreading: new paleomagnetic and Ar/Ar
1394 evidence. Tectonophysics, 347, 231– 251.
1395 Spina, V, Tondi, E., Mazzoli, S., 2011. Complex basin development in a wrench-dominated back-arc area:
1396 Tectonic evolution of the Crati Basin, Calabria, Italy. J. Geodyn., 51, 90-109.
1397 doi:101016/jjog201005003
1398 Stampfli, G.M., Mosar, J., Marquer, D., Marchant, R., Baudin, T., Borel, G., 1998. Subduction and
1399 obduction processes in the Swiss Alps. Tectonophysics, 296, 159–204.
1400 Stern, R.J., 2004. Subduction initiation: spontaneous and induced. Earth Planet. Sci. Lett., 226, 275-292.
1401 doi:10.1016/j.epsl.2004.08.007
1402 Sulli, A., 2000. Structural framework and crustal characteristics of the Sardinia Channel Alpine transect in
1403 the central Mediterranean. Tectonophysics, 324, 321-336.
1404 Sulli, A., Lo Presti, V., Morticelli, M.G., Antonioli, F., 2013. Vertical movements in NE Sicily and its
1405 offshore: Outcome of tectonic uplift during the last 125 ky. Quaternary International, 288, 168-182.
1406 doi:10.1016/j.quaint.2012.01.021
1407 Tamburelli, C., Babbucci, D., Mantovani, E., 2000. Geodynamic implications of subduction related
1408 magmatism: insights from the Tyrrhenian-Apennines region. J. Volcanol. Geotherm. Res., 104, 33-43.
1409 Tansi, C., Muto, F., Critelli, S., Iovine, G., 2007. Neogene-Quaternary strike-slip tectonics in the central
1410 Calabria Arc (southern Italy). J. Geodyn., 43, 397-414. doi:101016/jjog200610006
1411 Tari, V., Pamic, J., 1998. Geodynamic evolution of the northern Dinarides and the southern part of the
1412 Pannonian Basin. Tectonophysics, 297, 269–281.
1413 Tesòn, E., Teixell, A., 2008. Sequence of thrusting and syntectonic sedimentation in the eastern Sub-Atlas
1414 thrust belt (Dade` s and Mgoun valleys, Morocco). Int. J. Earth Sci., 97, 103–113. DOI
1415 10.1007/s00531-006-0151-1
1416 Tibaldi, A., Pasquarè, F., Tormey, T. 2010. Volcanism in reverse and strike-slip fault settings, in: Cloetingh,
1417 S., Negendank, J. (Eds.), New Frontiers in Integrated Solid Earth Sciences, Springer Netherlands,
1418 Berlin, 315-348. http://dx.doi.org/10.1007/978-90-481-2737-5_9

1419 Tondi, E., Cello, G., 2003. Spatiotemporal evolution of the central Apennines fault system (Italy). J.
1420 Geodyn., 36, 113-128.
1421 Tripodi, V., Muto, F., Brutto, F., Perri, F., Critelli, S., 2018. Neogene-Quaternary evolution of the forearc
1422 and backarc regions between the Serre and Aspromonte Massifs, Calabria (southern Italy). Mar. Pet.
1423 Geo., 95, 328-343. https://doiorg/101016/jmarpetgeo201803028
1424 Tropeano, M., Sabato, L., Pieri, P., 2002. Filling and cannibalization of a foredeep: the Bradanic Trough
1425 (Southern Italy), in: Jones, S.C., Frostick L.E. (Eds.) Geol Sediment Flux to Basins: Causes, Controls
1426 and Consequences. Geol. Soc., London, Spec. Publ., 191, 55-79.
1427 Van Dijk, J.P., Scheepers, P.J.J., 1995. Neotectonic rotations in the Calabrian Arc; implications for a
1428 Pliocene-Recent geodynamic scenario for the Central Mediterranean. Earth Sci. Rev., 39, 207-246.
1429 van Hinsbergen, D.J.J., Reinoud, L.M.V., Spakman, W., 2014. Origin and consequences of western
1430 Mediterranean subduction, rollback, and slab segmentation. Tectonics, 33, 393-419,
1431 doi:10.1002/tect.20125
1432 Vannoli, P., Basili, R.,Valensise, G., 2004. New geomorphic evidence for anticlinal growth driven by blind-
1433 thrust faulting long the northern Marche coastal belt (central Italy), J. Seismol., 8, 3, 297-312.
1434 Vannoli, P., Burrato, P., Valensise, G., 2015. The Seismotectonics of the Po Plain (Northern Italy): Tectonic
1435 Diversity in a Blind Faulting Domain. Pure Appl. Geophys., 172, 1105–1142
1436 Ventura, G., Vilardo, G., Milano, G., Pino, N.A., 1999. Relationships among crustal structure, volcanism and
1437 strike-slip tectonics in the Lipari - Vulcano Volcanic complex (Aeolian Islands, southern Tyrrhenian
1438 sea, Italy). Phys. Earth Planet. In., 116, 31-52.
1439 Viola, G., Mancktelow, N.S., Seward, D., 2001. Late Oligocene–Neogene evolution of Europe Adria
1440 collision: new structural and geochronological evidence from the Giudicarie fault system (Italian
1441 Eastern Alps). Tectonics, 20, 999–1020.
1442 Viti, M., Albarello, D., Mantovani, E., 1997. Rheological profiles in the central-eastern Mediterranean.
1443 Annali di Geofisica, 40, 849-864
1444 Viti M., De Luca J., Babbucci D., Mantovani E., Albarello D., D'Onza F., 2004. Driving mechanism of
1445 tectonic activity in the northern Apennines: Quantitative insights from numerical modeling. Tectonics,
1446 23, 1-16, TC4003, doi:10.1029/2004TC001623

1447 Viti, M., Mantovani, E., Babbucci, D., Tamburelli, C., 2006. Quaternary geodynamics and deformation
1448 pattern in the Southern Apennines: implications for seismic activity. Boll. Soc. Geol. Ital., 125, 273-
1449 291.
1450 Viti, M., Mantovani, E., Tamburelli, C., Babbucci, D., 2009. Generation of trench-arc-backarc systems in the
1451 Western Mediterranean region driven by plate convergence. Boll. Soc. Geol. Ital., 128, 1, 89-106.
1452 Viti, M., Mantovani, E., Babbucci, D., Tamburelli, C., 2011. Plate kinematics and geodynamics in the
1453 Central Mediterranean. J. Geodyn., 51, 190-204.
1454 Viti M., Mantovani E., Cenni N., Vannucchi, A. 2012. Post seismic relaxation: an example of earthquake
1455 triggering in the Apennine belt (1915-1920). J. Geodyn., 61, 57-67,
1456 http://dx.doi.org/10.1016/j.jog.2012.07.002.
1457 Viti M., Mantovani E., Cenni N., Vannucchi, A. 2013. Interaction of seismic sources in the Apennine belt.
1458 Phys. Chem. Earth, 63, 25-35, http://dx.doi.org/10.1016/j.pce.2013.03.005.
1459 Viti M., Mantovani E., Babbucci D., Cenni N., Tamburelli C., 2015a. Where the Next Strong Earthquake in
1460 the Italian Peninsula? Insights by a Deterministic Approach. Boll. Geof. Teor. Appl., 56, 329-350, doi:
1461 10.4430/bgta0137
1462 Viti, M., Mantovani, E., Babbucci, D., Tamburelli, C., Cenni, N., Baglione, M., D'Intinosante, V., 2015b.
1463 Belt-parallel shortening in the Northern Apennines and seismotectonic implications. Int. J.
1464 Geosciences, 6, 938-961, http://dx.doi.org/10.4236/ijg.2015.68075.
1465 Viti, M., Mantovani, E., Babbucci, D., Tamburelli, C., Cenni N., 2016. Seismotectonics of the Padanian
1466 region and surrounding belts: Which driving mechanism? Int. J. Geosciences, 7, 1412-1451,
1467 http://dx.doi.org/10.4236/ijg.2016.712100.
1468 Volpi V., Del Ben A., Civile D., Zgur F. 2017. Neogene tectono-sedimentaty interaction between the
1469 Calabrian Accretionary Wedge and the Apulian foreland in the northern Ionian Sea. Mar. Pet.Geol.,
1470 83, 246-260.
1471 Westaway, R., 1993. Quaternary uplift in Southern Italy. J. Geophys. Res., 98, 21741-21772.
1472 Wölfler, A., Kurz, W., Fritz, H., Stüwe, K., 2011. Lateral extrusion in the Eastern Alps revisited: Refining
1473 the model by thermochronological, sedimentary, and seismic data. Tectonics, 30, TC4006,
1474 doi:10.1029/2010TC002782

Wortel, M.J.R., Spakman, W., 2000. Subduction and slab detachment in the Mediterranean–Carpathian region. Science, 290, 1910–1917.

Zampieri, D., Massironi, M., Sedea, R., Saracino, V., 2003. Strike-slip contractional stepovers in the Southern Alps (northeastern Italy). Eclogae Geol. Helv., 96, 115-123.

Zecchin, M., Nalin, R., Roda, C., 2004. Raised Pleistocene marine terraces of the Crotone peninsula (Calabria, southern Italy): facies analysis and organization of their deposits. Sediment. Geol., 172, 165–185, https:// doi.org/10.1016/j.sedgeo.2004.08.003

Zecchin, M., Caffau, M., Tosi, L., Civile, D., Brancolini, G., Rizzetto, F., Roda, C., 2010. The impact of Late Quaternary glacio-eustasy and tectonics on sequence development: evidence from both uplifting and subsiding settings in Italy. Terra Nova 22, 324–329.

Zecchin, M., Civile, D., Caffau, M., Sturiale, G., Roda, C., 2011. Sequence stratigraphy in the context of rapid regional uplift and high-amplitude glacio-eustatic changes: the Pleistocene Cutro Terrace (Calabria, southern Italy). Sedimentology 58, 442–477.

Zecchin, M., Praeg, D., Ceramicola, S., Muto, F., 2015. Onshore to offshore correlation of regional unconformities in the Plio-Pleistocene sedimentary successions of the Calabrian Arc (central Mediterranean). Earth Sci. Rev., 142, 60–78. http://dx.doi.org/10.1016/j.earscirev.2015.01.006

Zecchin, M., Donda, F., Forlin, E., 2017. Genesis of the Northern Adriatic Sea (Northern Italy), since early Pliocene. Mar. Pet. Geo., 79, 108-130.

Figure captions

Fig. 1. Comparison between the presumed Oligocene (A) and Present (B) tectonic settings in the Mediterranean area. Al-Ka-Pe-Ca=tentative assemblage of the Alboran, Kabylides, Peloritani and Calabrian fragments of the Alpine belt. 1, 2) Continental and thinned continental Eurasian domains 3, 4) Continental and thinned continental African and Adriatic domains 5) Old oceanic domains 6) Alpine belt 7) Other orogenic belts 8,9) Tectonically thinned and oceanized zones 10) Outer fronts

of belts 11, 12, 13) Compressional, extensional and transcurrent features. BP, CS=Balearic Promontory and Corsica-Sardinia fragments of the Iberian foreland. VH=Victor-Hensen fault. Blue arrows indicate the kinematics of the Africa/Adriatic domain and the Anatolian-Aegean-Pelagonian belt with respect to Eurasia (Mantovani et al., 2007a; Viti et al., 2009, 2011).

Fig. 2. A) Late Oligocene-Early Miocene. The outward migration of the Norhern Al-Ka-Pe-Ca Arc (NA) and the Corsica-Sardinia block (CS), at the expense of the Tethyan domain, builds up the Apennine accretionary belt, while crustal extension occurs in the wake of the Arc, forming the Balearic basin (BB). A more limited rotation is undergone by the Southern Al-Ka-Pe-Ca arc (SA) and the Balearic Promontory (BP), generating the relatively small Valencia trough (VT). The relative motion between NA and SA is accommodated by the North Balearic fault (NBF). **B**) **Middle Miocene.** The NA stops rotating after its collision with the continental Adriatic domain, causing the end of crustal extension in the Balearic basin, while the SA continues its migration at the expense of the Tethyan domain, until reaching the continental African domain. Back-Arc extension develops in the wake of SA, generating the Eastern Algerian basin (EAB). (From Viti et al., 2009, modified). Colours and symbols as in figure 1.

Fig. 3. Formation of the Northern Tyrrhenian basin (NTB), from the middle-upper Miocene (**A**) to the late Miocene (**B**). ESA=Eastern Southern Alps, Gi= Giudicarie fault system, NAp=Northern Apennines, SAp=Southern Apennines, TW=Tauern window, WPa= Western Padanian protuberance. Colours and symbols as in figure 1.

Fig. 4. A) Late Miocene-Early Pliocene (about 5 My). Crustal stretching occurs in the Alpine-Apennine sector lying south of the Selli fault (SF), generating the Central Tyrrhenian basin (CT). See text for the description of the other major coeval tectonic events. AB=Adventure block, Eg=Egadi fault, HB=Hyblean domain, Me=Medina fault, Pd=Padanian, Pl=Pantelleria trough,

Sci=Sciacca- fault, SV=Schio-Vicenza fault, VH=Victor-Hensen fault. **B) Late Pliocene** (2 My). Ge=Gela nappe, Li=Linosa trough, Ma=Malta trough Ta=Taormina fault. Colours, symbols and other abbreviations as in figures 1 and 3**.**

Fig. 5. Formation of the Southernmost Tyrrhenian basin (ST) in the early Pleistocene. AS=Apulian Swell, CP=Calabria-Peloritani wedge, ECA=External Calabrian Arc, Pa=Palinuro fault, SAp=Southern Apennines wedge. Colours, symbols and other abbreviations as in figures 1, 3 and 4.

Fig. 6. Present tectonic setting. 1) Africa-Adriatic continental domains, 2) Quaternary magmatism, Ca=Catanzaro trough, Cam=Campanian magmatic province, Cr=Crati trough, CV=Capo Vaticano fault, LuAp=Lucanian Apennines, MAR= Middle Adriatic Ridge, Me=Mesima trough, NAp, CAp, and SAp=Northern, Central and Southern Apennines, Rom=Roman magmatic province, Si=Sibari fault, Sy=Syracuse fault, Vu=Vulcano fault. Other colours, symbols and abbreviations as in figures 1-5.

Fig. 7. Horizontal velocity field (vectors) with respect to a fixed Eurasian frame in the ITRF2014 reference frame (Altamimi et al., 2016), obtained by GPS measurements. Scale in the bottom. Colours of GPS sites indicate velocities, in agreement with the chromatic scale given on the left. See Cenni et al. (2012, 2013) for details about the network and data analysis.

Fig. 8. Sketch of the tectonic process that is supposed to generate a Arc-Trench-Back Arc system. **A)** An orogenic belt, flanked by an oceanic domain, is longitudinally stressed by a continental indenter. **B**) The stressed belt undergoes uplift and bowing, through the lateral escape of crustal wedges, at the expense (subduction) of the adjacent oceanic domain. **C**) The separation of the migrating Arc from the stable plate induces crustal extension in the interposed zone (Back Arc

1555 basin). See Driehaus et al. (2013) and Boutelier et al. (2018) for laboratory modeling of the above
1556 process.
1557
1558 **Fig. 9.** Proposed evolutionary reconstruction. **A) Middle-Upper Miocene (15-9 My).** Blue arrows
1559 indicate the the proposed kinematic pattern (Mantovani et al., 2007a, 2009, 2015a; Viti et al., 2009,
1560 2011) **B) Upper Miocene (9-6 My**). After the reactivation of the Giudicarie fault system, the
1561 northern Adriatic promontory undergoes a NE ward motion and clockwise rotation, releasing its
1562 previous internal deformation. The divergence between that promontory and the stable Corsica-
1563 Sardinia block induces crustal extension in the interposed Alpine-Apennine belt, with the formation
1564 of the Northern Tyrrhenian basin. **C) Pliocene (5-2 My).** A large part of the Adriatic promontory
1565 decouples from Africa, by the activation of major discontinuities in the Pelagian and Ionian zones,
1566 the Sicily Channel-Medina (Me)-Victor Hensen (VH) fault systems and from its Padanian sector
1567 (Pd), by the reactivation of an old fracture in the northern Adriatic domain (the Schio-Vicenza fault
1568 system=SV). The E-W compression induced by the convergence between the southernmost Adria
1569 block and the northern African foreland (Tunisia) causes the roughly NW ward extrusion of a
1570 continental fragment, the Adventure block (AB), guided by the Egadi (Eg) and the Sciacca (Sci)
1571 fault systems. The northward indentation of such block causes eastward escape of wedges from the
1572 Alpine-Apennine belt lying south of the Selli fault, at the expense of the Tethyan domain and the
1573 thinned Adriatic western margin. Thrusting develops in front of the extruding wedges (Southern
1574 Apennines). Crustal stretching takes place in the wake of such wedges, forming the central
1575 Tyrrhenian basin (CT). **D) Early Pleistocene.** After the stop of the Southern Apennines wedge
1576 against the continental Adriatic domain, the convergence of the confining plates is accommodated
1577 by the outward extrusion of the CP wedge, at the expense of the Ionian domain, and the upward
1578 flexure of the southern Adriatic platform, forming the Apulian Swell (AS). Thrusting at the outer
1579 front of the Calabrian wedge forms the External Calabrian Arc (ECA), while extension at the inner
1580 side forms the Southern Tyrrhenian basin (ST). The Calabrian wedge undergoes strong uplift,

bowing and fragmentation. **E) Middle-Upper Pleistocene.** The potential gravitational energy accumulated by the southern Adriatic favors the northward displacement of that plate, which induces a longitudinal compression in the outer sector of the Apennine belt. Such regime is accommodated by the outward extrusion of wedges, which separate from the inner belt. After the contact with the Adriatic continental domain, the extrusion of the Calabrian wedge is guided by new lateral guides, the Sibari (Si) and Vulcano-Syracuse (Vu-Sy) faults. Colours, symbols and other abbreviations as in figures 1-6.

Fig. 10. Tentative reconstruction of the E-W shortening processes in the Hyblean-Adventure domain that determined the formation of troughs in the Sicily Channel. **A)** Late Miocene. Configuration of the Hyblean-Adventure domain just after the activation of the Victor-Hensen-Medina-Sicily Channel discontinuity, when the shortening of that zone, induced by the E-W convergence between the decoupled Adria plate and the Tunisian protuberance (Fig.9c) started to develop. Present geographical contours are reported for reference. **B)** Present tectonic setting (modified after Finetti and Del Ben, 1986). The Adventure block, decoupled from the Hybelan domain by the Sciacca fault system, has undergone a roughly NNWward escape, forming the Pantelleria trough. Contemporaneously, the heterogeneous Hyblean domain has undergone E-W shortening, at the expense of its thinned central part. In the northern side, the Maghrebian belt has been forced to bend southward, forming the Gela nappe. In the southern side, the original NW-SE plateau, in the Sicily channel zone, has undergone a considerable fragmentation, in order to accommodate E-W shortening. Local crustal extensions, induced by the divergence between fragments, have formed the Malta and Linosa grabens and other minor troughs. Toothed lines indicate the fronts of belts. Geological-geophysical data from Finetti and Del Ben, 1986, 2005b; BenAvraham et al., 1990; Lentini et al., 1994). Lit, Mat, Plt =Linosa, Malta and Pantelleria troughs Em, Gi, Li, Ma, Se, Ur =Empedocle, Girgenti, Linosa, Malta, Selinunte, Urialo plateau fragments.

1607 **Fig. 11.** Outer mobile sector of the Apennine belt (green), stressed by the Adria plate**.**
1608 Ben=Benevento, Ca=Calabria, Ir=Irpinia, LA=Lazio-Abruzzi wedge, Lu=Lucania Apennines, Ma=
1609 Matese, MS=Molise-Sannio, No-Cf=Norcia-Colfiorito fault system, OA=Olevano-Antrodoco thrust
1610 front, RMU=Romagna-Marche-Umbria wedge, SVo= Sangro-Volturno thrust front, TE=Toscana-
1611 Emilia wedge. The buried external folds in the Northern Apennines are light green. Red arrows
1612 indicate the kinematic pattern, compatible with the Pleistocene deformation pattern and geodetic
1613 data (Fig. 7). Other symbols and abbreviations as in figures 1-6.
1614


A
EURASIA
Carpathians
Alps
Pelagonian
Dinarides
ADRIATIC
Balcanides
Pontides
Pyrenees
ATLANTIC - IBERIA
Ligurides
Hellenides
Aegean
Anatolia
Taurides
Bitlis-Zagros
Alpine Tethys
Ionian Tethys
Levantine Tethys
AFRICA
B
Carpathians
Pannonian
Alps
Dinarides
ADRIATIC
Pyrenees
Apennines
Balkans
Caucasus
ATLANTIC
Iberia
Valencia trough
Balearic
Tyrrhenian
Pelagonian
Hellenides
Anatolia
Alboran
Maghrebides
Aegean
Sicily Channel
ARABIA
Trans-Morocco fault system
AFRICA
5 mm/y
1 2 3 4 5 6 7 8 9 10 11 12 13


A
Alpine Maghrebian belt
Egadi
Adventure plateau
Sciacca
Hyblean thinned domain
Hyblean plateau
Tunisia
Ionian
Selinunte
Empedocle
Malta
Girgenti
Linosa
Urialo
Lampedusa plateau
Medina plateau
Medina fault


B
VENTURA PLATEAU
MAGHREBIAN CHAIN
ATLASIC FOLDS AND THRUSTS
TUNISIAN PLATFORM
TUNISIAN - LAMPEDUSAN PLATEAU
MEDINA PLATEAU
PLATEAU
Se
Em
Gi
Ma
Li
Ur


SV
Adriatic
TE
RMU
No-CP
AQ
LA
L'Aquila
Fucino
Rom
MS
Svo
Ma
Ben
Ir
Calm
Lu
Ca
5 mm/y
Adriatic plate
Calabrian units
Internal apennines
Quaternary magmatism










EURASIA
Insubric line
ESA
SV
ADRIA
Dinarides
NAp
MAR
Rom
CAp
Cam
TYRRHENIAN
SAp
LuAp
Si
Cr
Ca
Me
CV
VU
External Calabrian Arc
Hellenides
Sicily
AB
HB
Maghrebides
Sicily Channel
Sy
Ionian
AFRICA
1
2
46
44
42
40
38
36
34
8
10
12
14
16
18
20


47°
46°
45°
44°
43°
42°
41°
40°
39°
38°
37°
36°
35°
6°
7°
8°
9°
10°
11°
12°
13°
14°
15°
16°
17°
18°
19°
Vel. (mm/y)
6
5
4
3
2
1
0
5 mm/y


Orogenic belt
S
S'
Lithosphere
Asthenosphere
A
Continental plate
S
S'
Continental plate
OROGENIC BELT
Oceanic zone
Back-arc basin
Arc
B
ARC
Back-arc basin
C
BASIN
ARC


A
B
C
D
E
ADRIATIC
Apennines
Dinarides
Pelagonian
Hellenides
Rhodope
Aegean
Ionian zone
Ionian Tethys
Maghrebides
Pelagian zone
AFRICA
Corsica - Sardinia
TW
ESA
WPa
Pannonian
NAp
NTB
SAp
Sella
Calabria
PLATE
AB
HB
Sicily Channel
Me
IH
Pd
SV
CT
ST
AS
Ta
Ge
ECA
ALPS
MAR
Ionian
5 mm/y

The observed deformation pattern in the study area is not compatible with slab pull forces, whereas it can be coherently and plausibly interpreted as an effect of plate convergence

Crustal extension can develop in the framework of extrusion processes

The subduction of oceanic lithosphere at convergent boundaries may require the triggering favoured by an extrusion process

Declaration of interests

☒The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: