

Ozonated oils as functional dermatological matrices: Effects on the wound healing process using SKH1 mice

This is a pre print version of the following article:
Original:
Valacchi, G., Zanardi, I., Lim, Y., Belmonte, G., Miracco, C., Sticozzi, C., et al. (2013). Ozonated oils as functional dermatological matrices: Effects on the wound healing process using SKH1 mice. INTERNATIONAL JOURNAL OF PHARMACEUTICS, 458(1), 65-73 [10.1016/j.ijpharm.2013.09.039].
Availability:
This version is availablehttp://hdl.handle.net/11365/45325 since 2016-11-19T10:36:41Z
Published:
DOI:10.1016/j.ijpharm.2013.09.039
Terms of use:
Open Access The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license. For all terms of use and more information see the publisher's website.

(Article begins on next page)

AUTHOR QUERY FORM

	Journal: IJP	Please e-mail or fax your responses and any corrections to:
		E-mail: corrections.esch@elsevier.thomsondigital.com
ELSEVIER	Article Number: 13673	Fax: +353 6170 9272

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult http://www.elsevier.com/artworkinstructions.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.

Location in article	Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof
$\frac{Q1}{Q2}$ $Q3$	Please confirm that given names and surnames have been identified correctly. The citation "Ali et al. (2012)" has been changed to match the author name/date in the reference list. Please check here and in subsequent occurrences, and correct if necessary. Figs. 4 and 5 will appear in black and white in print and in colour on the web. Based on this, the respective figure captions have been updated. Please check, and correct if necessary.
	Please check this box or indicate your approval if you have no corrections to make to the PDF file

Thank you for your assistance.

SEV

ARTICLE IN PRESS

International Journal of Pharmaceutics xxx (2013) xxx-xxx

Contents lists available at ScienceDirect

International Journal of Pharmaceutics

journal homepage: www.elsevier.com/locate/ijpharm

A THE ERATORAL OUR ALL OF PHARMAC BAR OF PHARMAC BA

Graphical Abstract



International Journal of Pharmaceutics xxx (2013) xxx-xxx



Contents lists available at ScienceDirect

International Journal of Pharmaceutics



journal homepage: www.elsevier.com/locate/ijpharm

Ozonated oils as functional dermatological matrices: Effects on the wound healing process using SKH1 mice

Q1 , G. Valacchi^{a,b,**}, J. Zanardi^c, Y. Lim^b, G. Belmonte^d, C. Miracco^e, C. Sticozzi^a, V. Bocci,^f, V. Travagli,^{c,*}

^a <mark>Pipartimento</mark> di Scienze della Vita e Biotecnologie, Università degli Studi di Ferrara, Italy

^b Department of Food and Nutrition, Kyung Hee University, Seoul, South Korea

Dipartimento di Biotecnologie, Chimica e Farmacia, Università degli Studi di Siena, Italy

^d Dipartimento di Scienze Biomediche, Università degli Studi di Siena, Italy

^e <mark>Pipartime</mark>nto di Scienze Mediche, Chirurgiche e Neuroscienze, Università degli Studi di Siena, Italy

^f Dipartimento di Fisiologia, Università degli Studi di Siena, Italy

ARTICLE INFO

Article history: Received 10 July 2013 Accepted 30 September 2013 Available online xxx

Keywords:

10 11

12 13

14

15

16

17

18

19

20

23

24

25

26

27

28

29

30

31

32

33

34

Wound healing 1,2,4-Trioxolane moiety

Prodrugs

21 Vegetable matrices 22

ABSTRACT

Wound tissue repair is a complex and dynamic process of restoring cellular structures and tissue layers. Improvement of this process is crucial for several pathologies characterized by chronic delayed wound closure such as diabetes, and the investigation of new approaches aimed to ameliorate the wound healing process is under continuous evolution. Recently, the usage of vegetable matrices in the form of ozonated oils has been proposed and several researchers have shown a positive effect in the wound, based on their bactericidal, antiviral, and antifungal properties. The present study was undertaken to compare the effect that different ozonated oils (olive, sesame and linseed) with the same level of ozonation have on wound healing rate in SKH1 mice. Several histological parameters and the level of key proteins such as VEGF and PCNA have been analyzed. Only treatment with ozonated sesame oil shows a faster wound closure in the first 7 days. This effect paralleled with the increased VEGF and PCNA levels, NFkB nuclear translocation and 4-HNE formation. The present study shows that not only the ozonation grade is of importance for the improvement of wound healing process but also the typical composition of the oil.

© 2013 Published by Elsevier B.V.

1. Introduction

In normal skin, the epidermis (outermost layer) and dermis (inner or deeper layer) exists in a steady-state equilibrium, forming a protective barrier against the external environment. Once the protective barrier is broken, a set of complex biochemical events takes place in a closely orchestrated cascade to repair the damage. Such a process defined wound healing (WH) is immediately set in motion to prevent infectious events. The closure of cutaneous wounds involves complex tissue movements such as haemorrhage. inflammation, re-epithelization (proliferation), granulation tissue formation, and the late remodelling phase of repair (Werner and Grose, 2003).

* Corresponding author at: Dipartimento di Biotecnologie, Chimica e Farmacia, Università degli Studi di Siena, Viale Aldo Moro, 2, 53100 Siena, Italy. Tel.: +39 0577 234317; fax: +39 0577234254.

** Corresponding author at: Dipartimento Scienze della Vita e Biotecnologie, Università degli Studi di Ferrara, Via Borsari, 26, 44100 Ferrara, Italy.

Tel.: +39 0532455482; fax: +39 0532455550.

E-mail addresses: giuseppe.valacchi@unife.it (G. Valacchi), valter.travagli@unisi.it (V. Travagli).

0378-5173/\$ - see front matter © 2013 Published by Elsevier B.V. http://dx.doi.org/10.1016/j.ijpharm.2013.09.039

Previous studies have demonstrated that endogenous factors are able to modulate and coordinate the healing process, such as vascular endothelial growth factor (VEGF), transforming growth 37 factor β (TGF β), platelet-derived growth factor (PDGF), cellular 38 cycle proteins and even reactive oxygen species (ROS). All these factors are released and expressed from different cells involved in the WH process such as macrophages, fibroblasts, and keratinocytes at the site of injury and they participate in the regulation of reepithelization, granulation tissue formation, collagen synthesis and neovascularization (Barrientos et al., 2008; de Melo et al., 2011).

The ability to modulate the levels and the release of the mentioned growth factors can influence the WH process. In addition, infection of the wound will delay the wound closure (Edwards and Harding, 2004).

Ozone (O_3) is widely recognized as one of the best bactericidal, 49 antiviral, and antifungal molecules used in the therapy of chronic 50 wounds. The beneficial effects of O₃ on WH might be assumed to be 51 due to decreased bacterial infection, ameliorated dermal WH rate 52 and increased oxygen tension by O₃ exposure in the wound area 53 (Valacchi et al., 2012). In an aqueous environment, O₃ does not pen-54 etrate the cells but it instantaneously reacts with the double bonds 55 of polyunsaturated fatty acids (PUFA) leading to the formation of

G Model IJP 13673 1–9

2

57

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

80

81

82

83

84

85

86

87

88

89

92

93

95

97

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

ARTICLE IN PRESS

G. Valacchi et al. / International Journal of Pharmaceutics xxx (2013) xxx-xxx

ROS and bioactive products such as aldehydes (Pryor and Church, 1991). These "second messenger" of O_3 reactivity, via the activation of redox transcription factors such as nuclear factor-kappaB (NF κ B), can induce the synthesis of growth factors and accelerate the cell cycle.

Recently, it has been suggested that ozonated oils could be functional matrices to slowly deliver O₃ messengers and improve WH due to the ability of the oil to stabilize O3 in a suitable chemical form (Travagli et al., 2010). Moreover, in an experimental, well-characterized skin lesion in mice it had been demonstrated that the repeated application of ozonated sesame oil (ozSO) was able to significantly accelerate the first phase of the wound closure which is more susceptible to infections (Valacchi et al., 2011). This effect was mainly observed at a certain level of ozonation, emphasizing that the peroxidation grade is an essential parameter to keep under consideration in the modulation of wound closure process. In fact, the peroxide value is an indirect outcome of the 1,2,4-trioxolane moiety that represents the active compound of the ozonated vegetable matrices (Travagli et al., 2009). Apparently, when this annular ozonide comes into contact with the altered skin in the presence of exudates at the body temperature, it decomposes to reactive ozone derivatives, which readily dissolve in the aqueous biological milieux generating both hydrogen peroxide and a variety of oxidized compounds that improve the local metabolism and enhances healing, even in the presence of skin infectious diseases. In the present paper, the relevance of three different oils such as olive oil, sesame oil and linseed oil at a specific degree of ozonation in a wound healing model using SKH1 mice had been studied in terms of: (i) wound closure rate; (ii) skin structure; (iii) activation of the transcription factor NFkB, together with the expression of key proteins involved in the wound healing process such as VEGF (yascularization) and proliferating cell nuclear antigen (PCNA) as well as the levels of oxidative stress markers such as 4-hydroxynonenal (4-HNE).

2. Materials and methods

2.1. Animals

Hairless female SKH-1 mice (6 weeks old) were purchased from Orient Bio Inc. (Gyeonggi-do, Korea) and lodged in individual plastic cages at temperature- and humidity-controlled conditions $(22 \pm 1 \,^{\circ}C, RH 50-60\%, 12 h light/dark cycle)$ with allowed access to distilled water and food. Mice were acclimated for 10 days before initiation of the treatment. Mice (n = 36) were divided in 4 groups, one group (n = 9) for each time-point (d0, d3, d7 and d14, as reported in Section 2.3). The animals were randomly selected at the start of the experiment and they were used in accordance with animal protocols approved by the Kyung Hee University Institutional Animal Care and Use Committee.

2.2. Wound biopsy and measurement of wound closure

The skin was pinched and folded on the dorsum (anaesthetization with isoflurane, sterile biopsy punch 3.5 mm diameter, Miltex Instrument Company, York, PA), below the shoulder so as to obtain height circular wounds identical in size, as previously described (Lim et al., 2006).

This area was selected for wounding as it cannot be reached by the mice thereby preventing self-licking and wound irritation. Wounds in individual mouse were photographed digitally every day, beginning on the day of wounding (d0) with a standard dot equivalent to the initial wound area placed beside the wound. The quantification of wound closure was performed as previously described (Lim et al., 2006). Wound closure was quantified by Canvas 11SE software (Deneba, Miami, FL). The rate of wound closure was expressed as the % ratio of wound area (each day after wounding) compared with the initial wound area. Decreased wound ratio indicates the amelioration of wound closure.

As to concern the immunohistochemical analysis skin samples were harvested at the following time points: days 0, 3, 7, and 14 (d0, d3, d7, and d14, respectively).

2.3. Ozonated oil and topical application

Olive oil (OO), sesame oil (SO), and linseed oil (LO) of pharma-125 ceutical grade were purchased from Galeno Srl (Comeana, Italy). 126 Table 1 summarizes the main chemical-physical properties of the 127 different samples used. Specifically, the oils have been selected for 128 their compositions in terms of n-3, n-6 as well as n-9 fatty acids. 129 The real percentages as stated in the manufacturer's data sheet 130 compared to the intervals reported in the literature (in brackets) 131 have been reported (Crews et al., 2006; European Pharmacopoeia, 132 2010, 2011). Ozonated samples of olive oil (ozOO), sesame oil 133 (ozSO), and linseed oil (ozLO) were obtained according to Zanardi 134 et al. (2008) and Sega et al. (2010). The ozonation has been con-135 ducted until a peroxide value of about 1500 was reached. Physical 136 characterization has been performed by viscosity measurements 137 (Viscomate VM-10AL, CBC Europe) at both the temperatures of 22 138 and 35 ± 0.2 °C. Eight microlitres of untreated or ozonated oil was 139 applied twice a day on the corresponding wound. As control groups, 140 nothing was applied on the seventh and eight wound. 141

2.4. Immunohistochemical analysis

Surgical excision of the wounded area was performed at days 0, 143 3, 7, and 14 (d0, d3, d7, and d14, respectively) in each group. Skin 144 tissue was immersion fixed in 10% NBF (neutral-buffered forma-145 lin) for 24 h at RT and embedding in paraffin block. Sections $(4 \mu m)$ 146 were deparaffinized in xylene and rehydrated in alcohol gradients 147 (Sticozzi et al., 2012). For immunohistochemistry, after dewax-148 ing, sections were incubated overnight at 4°C with the following 149 antibodies: VEGF (polyclonal antibody; Millipore, Milan, Italy; dilu-150 tion factor 1:300; 0.05% trypsin-CaCl₂ buffer pH 7.8 pretreatment 151 for 10 min at 37 °C, for antigen retrieval); NFκB (polyclonal anti-152 body; Santa Cruz, DBA, Milan, Italy; dilution factor 1:100; Na-citrate 153 buffer pH 6 pretreatment for 10 min at 95 °C); PCNA (polyclonal 154 antibody; Biovision, Milan, Italy; dilution factor 1:100), and 4-155 HNE (polyclonal antibody; Millipore, Milan, Italy; dilution factor 156 1:200). Then the slides were washed three times with PBS and 157 endogenous peroxidase was blocked with 3% hydrogen peroxide 158 in distilled water for 10 min at RT. Finally, after aspecific bind-159 ing block, they were incubated with the EnVision+System-HRP 160 (DAKO, Glostrup, Denmark) for 1 h at RT. The reaction products 161 were stained with diaminobenzidine (DAB), counterstained with 162 Mayer's Haematoxylin and after drying were mounted with Eukitt 163 mounting medium. The semi-quantitative analysis was performed 164 by measuring the saturation of reaction by using Thresold colour 165 HSB of Image] software. 166

2.5. Preparation of total and nuclear extracts for Western blotting

For Western blotting analysis, the tissue was homogenized and lysed in radioimmunoprecipitation assay (RIPA) buffer supplemented with protease and phosphatase inhibitors for total protein extracts, as previously described (Valacchi et al., 2009).

For nuclear extracts, tissue was homogenized and then lysed in hypotonic buffer containing 10 mM HEPES (pH 7.9), 10 mM KCl, 1.5 mM MgCl₂, 0.3% Nonidet P-40, 0.5 mM dithiothreitol, 0.5 mM phenylmethylsulphonyl fluoride, protease inhibitor cocktail, 1 mM orthovanadate, 5 mM β -glycerophosphate. The lysates

142

167

121

122

123

124

G. Valacchi et al. / International Journal of Pharmaceutics xxx (2013) xxx-xxx

Table 1

The main chemical-physical properties of the different samples used.

Sample	Chemical composition			Peroxide value (mEq/kg)	Viscosity (mP	as)
	<mark>n-</mark> 3 content	n-6 content	n-9 content		$22\pm0.2^{\circ}\text{C}$	$35\pm0.2^\circ\text{C}$
Olive Oil (OO)	0.6% (max 1.2%)	8.8% (3.5–20%) ^a	72.9% (56.0–85.0%) ^a	143 ± 3	78.7 ± 1.2	37.5 ± 0.5
Sesame Oil (SO)	0.5% (0.2-1.0%) ^b	48.1% (36.9-46.8%) ^b	43.1% (34.1-44.7%) ^b	198 ± 9	59.9 ± 1	34.2 ± 0.3
Linseed Oil (LO)	53.7% (35.0-65.0%) ^c	14.3% (11.0-24.0%) ^c	21.5% (11.0-35.0%) ^c	163 ± 7	55.9 ± 0.5	30.8 ± 0.1
Ozonated Olive Oil (ozOO)	n.a.,d	n.a. ^d	n.a. ^d	1612 ± 146	980 ± 23	470 ± 15
Ozonated Sesame Oil (ozSO)	n.a, <mark>4</mark>	n.a. ^d	n.a. ^d	1543 ± 44	108 ± 1.3	61.3 ± 0.3
Ozonated Linseed Oil (ozLO)	n.a, <mark>d</mark>	n.a. ^d	n.a. ^d	1535 ± 78	81.3 ± 0.5	$\textbf{38.2}\pm\textbf{0.2}$

^A European Pharmacopoeia (2011).

Crews et al. (2006).

European Pharmacopoeia (2010).

^d Not available.

179

181

183

184

186

187

188

189

190

191

102

193



were incubated for 15 min on ice and then centrifuged at $1500 \times g$ 177 for 5 min at 4°C for collection of the supernatants containing 178 cytosolic proteins. Pellets containing crude nuclei were resuspended in extraction buffer containing 20 mM HEPES (pH 7.9), 180 1.5 mM MgCl₂, 0.6 M KCl, 0.2 mM EDTA, 20% glycerol, and 0.5 mM phenylmethylsulphonyl fluoride, protease inhibitor cocktail, 1 mM 182 orthovanadate, $5 \text{ mM} \beta$ -glycerophosphate and then were incubated for 30 min on ice. The samples were centrifuged at $10,000 \times g$ for 15 min to obtain supernatants containing nuclear fractions. 185

2.6. Western blot analysis

Briefly, 60 µg boiled protein was loaded onto 10% sodium dodecyl sulphate, polyacrylamide electrophoresis gels and separated by molecular size. Gels were electro-blotted onto nitrocellulose membranes and then blocked for 1 h in Tris-buffered saline, pH 7.5, containing 0.5% Tween20 and 5% milk. Membranes were incubated overnight at 4°C with either VEGF (Millipore Corporation, Billerica, MA, USA) or β -actin (Cell Signalling; Celbio, Milan, Italy),

and with either p65 subunit (Santa Cruz, CA, USA) or laminin 194 B (Cell Signalling; Celbio, Milan, Italy). The membranes were 195 then incubated with horseradish peroxidase-conjugated secondary 196 antibody for 1 h, and the bound antibodies were detected by chemi-197 luminescence (BioRad, Milan, Italy). β-actin was used as loading 198 control. 4',6-diamidino-2-phenylindole dihydrochloride hydrate 199 (DAPI, 1:10,000, Sigma–Aldrich D9642) was used to counterstain 200 the nuclei. Images of the bands were digitized and the densitometry 201 analysis was performed using Image-J software. 202

2.7. Histological evaluation of wound healing

Histological qualitative assessment of wound healing on skin biopsies was performed using 4 µm thick H&E-stained sections. They were examined for the presence of the following features: ulceration, necrosis, re-epithelization, inflammatory cells, myofibroblasts, vascularization, mastocytes, and collagen fibre bundle organization. VEGF and PCNA-positive cells were counted in immunostained sections in five to ten high power fields (HPFs).



Fig. 1. Effects of ozonated oils on PCNA levels. PCNA staining at d3, d7 and d14 of murine skin after the different treatments. (A) A higher number of PCNA-positive nuclei of keratinocytes are evident in basal/suprabasal epidermal layers in ozSO, when compared to the other groups (original magnification, 200; scale bar: 100 μm). Left diagram reports the number of cell PCNA positive in epidermis. (B) A higher number of PCNA-positive nuclei of fibroblasts/myofibroblasts are evident in dermal layers in ozSO (black arrows) with respect to control and SO treatment, where it is observed a persistence of inflammatory cells (white arrows; original magnification, 400; scale bar: 50 µm). Right diagram reports the number of cell PCNA positive in dermis. Values represent mean ± SEM for three separate experiments. *P<0.05.

Please cite this article in press as: Valacchi, G., et al., Ozonated oils as functional dermatological matrices: Effects on the wound healing process using SKH1 mice. Int J Pharmaceut (2013), http://dx.doi.org/10.1016/j.ijpharm.2013.09.039

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234 235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253 254

255

256

257

258

259

260

261

262 263

264

ARTICLE IN PRESS

G. Valacchi et al. / International Journal of Pharmaceutics xxx (2013) xxx-xxx

2.8. Picrosirius red staining of collagen fibres

Picrosirius red staining was performed on paraffin embedded sections. The sections were deparaffinized and rehydrated through five changes of decreasing gradient ethanol and the nucleus stained with Weigert's haematoxylin. Then, the sections were incubated in a solution containing picrosirius red 0.1% (Sigma_A-Aldrich Inc.) for 1 h at RT. Sections were then washed in two changes of acidified water, dehydrated in absolute ethanol, cleared in xylene, and mounted with Permount (Fisher Scientific Company). The sections were observed under light and polarization microscopy (Zeiss).

2.9. Statistical analysis

The results of quantitative parameters were statistically analyzed. The Kolmogorov–Smirnov test was applied to verify the normal distribution of sample data. In each group, the mean value per field was considered for epidermal thickness and microvessel density, and the median value per field for all the other parameters, for which a normal distribution was not observed. The Kruskal–Wallis and the paired Student's t-test were applied. A significance level of p < 0.05 was chosen.

3. Results

3.1. Effect of different ozonated oils on wound closure rate

Wound area was calculated from day 0 (d0) until complete closure of wound (d14), in terms of original wound size. As shown in Table 2 at d14 a wound closure greater than 90% was reached in all cases. It is evident that ozSO showed a significant increase in closure rate compared to SO already at d1 and it was statistical significant from d3 to d7 (p < 0.05). In detail, the increased WH rate was about 18% at d3, 42% at d4 and d5, 65% at d6 and 57% at d7. As far as ozOO, there was a similar trend observed for ozSO, but the wound closure rate was not statistically significant with respect to both air and OO controls. On the other hand, ozLO did not show any effect on wound closure rate.

3.2. Effect of ozonated oil on PCNA expression

Being WH process characterized by cellular proliferation, we have performed immunohistochemistry (IHC) analysis for PCNA, a marker of cellular proliferation and wound repair (Moldovan et al., 2007). As shown in Fig_A 1, ozSO was able to clearly increase skin PCNA levels as also reported by the semi-quantification of positive keratinocytes (panel A) and fibroblasts/myofibroblasts (panel B). This increase was statistically significant at d3 (2.5-fold increase, p < 0.05) and at d7 (2-fold increase, p < 0.05) for keratinocytes and at d7 for fibroblasts/myofibroblasts (p < 0.05) going back to the control levels in the latest time points. Both ozOO and ozLO did not show any significant effect on PCNA levels, although ozLO seems to induce PCNA levels in the last part of wound healing process (data not shown).

At the end of the wound healing process (d14) while in ozSO the skin presents a normal architecture with only few inflammatory cells, in control the reparative phenomena is still evident with persistence of inflammatory cells as well as of myofibroblasts and fibroblasts (panel B).

3.3. Effect of ozonated oils on VEGF levels

VEGF has been shown to be the main player in the regulation of all the phases of the wound closure process. As shown in Fig_A 2, both ozSO and ozOO increased VEGF production in the first 7 days

Fable 2 Quantific.	ation of wound e	closure in terms	s of % of original	l wound size <mark>.</mark>										
	d1	d2	d3	d4	d5	d6	d7	d8	6P	d10	d11	d12	d13	d14
Air	81.8 ± 2.5	80.7 ± 4.6	72.6 ± 6.7	66.6 ± 2.5	61.3 ± 4.6	50.6 ± 0.7	37.2 ± 4.7	30.8 ± 2.5	19.9 ± 5.4	16.5 ± 4.0	15.3 ± 3.0	11.9 ± 4.4	9.25 ± 2.2	8.47 ± 3.4
00	82.1 ± 4.3	79.7 ± 6.6	63.6 ± 4.5	62.0 ± 3.2	57.5 ± 2.5	50.7 ± 4.6	33.0 ± 3.7	23.8 ± 5.3	17.1 ± 3.5	16.5 ± 3.2	12.1 ± 2.1	9.49 ± 2.2	7.17 ± 1.2	6.02 ± 1.5
00z0	76.8 ± 10.6	69.1 ± 6.7	63.9 ± 4.4	49.7 ± 8.7	44.8 ± 6.6	38.1 ± 10.6	26.0 ± 1.5	18.1 ± 3.0	15.0 ± 4.4	10.3 ± 2.5	9.49 ± 5.4	6.82 ± 3.2	5.39 ± 2.1	4.06 ± 2.4
SO	77.8 ± 6.7	71.3 ± 7.8	68.1 ± 5.5	59.5 ± 5.0	49.6 ± 9.9	41.8 ± 2.5	23.3 ± 8.6	15.9 ± 3.7	9.21 ± 7.8	6.44 ± 2.3	5.70 ± 3.3	5.73 ± 2.5	5.39 ± 2.5	4.97 ± 1.2
OZZO	70.0 ± 6.7	63.9 ± 4.5	56.2 ± 3.6	$34.8\pm4.6^*$	$28.2 \pm 5.6^{*}$	$14.5\pm3.3^{*}$	$9.96\pm4.7^*$	8.51 ± 3.4	7.1 ± 2.4	5.41 ± 2.3	4.62 ± 2.2	3.80 ± 1.56	3.62 ± 2.5	3.04 ± 1.3
ΓO	82.2 ± 8.3	72.6 ± 5.8	70.6 ± 6.0	59.7 ± 6.8	55.8 ± 5.7	41.2 ± 3.7	19.6 ± 5.8	17.4 ± 2.3	13.0 ± 4.4	$\textbf{7.68} \pm \textbf{3.2}$	7.45 ± 2.1	6.18 ± 3.0	5.57 ± 4.4	5.40 ± 2.2
0ZL0	$\textbf{78.8} \pm \textbf{4.6}$	72.3 ± 3.7	63.7 ± 5.7	59.6 ± 4.7	54.9 ± 3.7	42.3 ± 4.6	21.9 ± 4.0	18.7 ± 3.5	12.5 ± 3.2	10.2 ± 2.1	8.30 ± 2.17	7.30 ± 2.45	6.46 ± 4.6	6.08 ± 0.7
Inter re	aresent mean +	SFM for three st	enarate exnerin	nents										

ARTICLE IN PRESS

G. Valacchi et al. / International Journal of Pharmaceutics xxx (2013) xxx-xxx



Fig. 2. Effects of ozonated oil on VEGF levels. (A) Vascular endothelial growth factors (VEGF) staining at d3, d7 and d14 of murine skin after the different treatments. VEGF positivity was observed in all the ozonated oils groups (ozOO, ozSO and ozLO) at d3, and only for ozOO and ozSO for d7 (original magnification, χ 200; scale bar: 100 µm). The diagram reports the number of cell VEGF positive. Values represent mean ± SEM for three separate experiments; *P<0.05. (B) Western blot for VEGF protein at d3, d7 and d14 of murine skin after the different treatments (ozOO, OO, ozSO, SO, ozLO and LO). Shown is a representative blot from three independent experiments; β -actin was used as loading control.

of the wound closure process. The induction of VEGF by ozOO was more evident than the ozSO at d7 (5-fold and 1.7-fold, respectively) while the effect was similar for d3 (circa 2-fold) (bottom panel). On the other hand, ozLO significantly induced VEGF levels at d3 (3fold). These data were confirmed by Western blot analysis for VEGF protein, as shown in panel B.

3.4. Effect of ozonated oils on cutaneous 4-HNE protein adducts

265

266

267

268

269

270

Wound healing is a redox controlled process, where oxidative 272 stress, ROS and lipid peroxidation products play a key role in its 273 modulation (Moseley et al., 2004). Therefore, as a marker of lipid 274 peroxidation and an index of oxidative stress (Uchida et al., 2004), we evaluated the levels of 4-HNE by IHC analysis during the wound 276 closure. As shown in Fig, 3, ozSO showed the highest level of 4-HNE 277 during the 14 days of wound closure. Cytoplasmic positivity to 4-278 279 HNE was mainly observed in suprabasal epidermal layers and this 280 was much evident than in the samples treated with ozOO and ozLO.

Untreated OO, SO and LO did not shown any difference from the controls. 281

3.5. NFkB activation

NFκB is a redox transcription factor, modulated by oxidative 284 stress and peroxidation products and the role of 4-HNE in the acti-285 vation of NFkB has been studied in several cells type (Ruef et al., **O2**86 2001; Ali and Sultana, 2012; Meng et al., 2012). Since the levels of 287 4-HNE were increased only with the ozSO treatment, the activation 288 of NFkB (p65 subunit) has been performed by both IHC and West-289 ern blot analysis only in the samples treated with ozSO. As shown 290 in Fig, 4, ozSO induced a clear activation of NFkB in cutaneous tis-291 sues. NFkB nuclear positivity (short arrow) is detected, mainly in 292 the upper epidermal layers at d3; it decreases at d7, and almost dis-293 appears on d14; in addition, several inflammatory/reparative cells 294 in the dermis also show nuclear positivity to NFkB (long arrow). 295 On the other hand, the control samples did not show a clear NFkB 296

283

6

297

298

200

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

ARTICLE IN PRESS G. Valacchi et al. / International Journal of Pharmaceutics xxx (2013) xxx-xxx



Fig, 3. Effect of ozonated oils on 4-HNE levels. At d3, as well as at d7 and d14, a cytoplasmic positivity to 4-HNE is mainly observable in suprabasal epidermal layers, with ozSO stronger than ozOO and ozLO (original magnification, ×400; scale bar: 50 µm).

nuclear positivity in the epidermis, whereas in the dermis positive nuclei were present in few inflammatory/reparative cells; furthermore, several inflammatory/reparative cell nuclei in the dermis are also positive to NFκB (long arrow). These data were also confirmed by Western blot analysis for p65 nuclear translocation using tissue nuclear extracts.

3.6. Effect of different ozonated sesame oil on skin collagen structure

As shown in Fig_A 5 collagen fibre organization and morphology was examined by picrosirius red stained sections of wound closure skin in both ozSO and control wounds. The wounds treated with ozSO showed a less extension of the injury at d3 and d7 compared to the control (arrow). It was possible to note that in ozSO the wound closure rate is faster than in the control as indicated by the collagen fibres (star symbol). Furthermore, under polarization microscopy, the wounds treated with ozSO showed a most amount of type I collagen fibres (yellow/orange appearance) with respect to control. On the other hand, in the wounds treated with SO it is observed a persistence of type III collagen fibres (green appearance) 315 characterizing the granulation tissue. 316

4. Discussion

This work is the continuance of our previous study where we 318 have shown that only a specific level of peroxidation (expressed 319 as peroxide value about 1500) of ozonated sesame oil is able to 320 significantly accelerate the first phase of wound healing in SKH1 321 mice (Valacchi et al., 2011). Therefore, this study was performed to 322 evaluate the wound healing properties of different ozonated oils, 323 such as olive oil, sesame oil and linseed oil with the same peroxide 324 values. The question is: is the peroxidation level the only factor 325 responsible for the wound healing properties of ozonated oils? 326

In general, our data agreed with our previous report, showing the beneficial effect of ozSO on the first phase of the wound healing process (Valacchi et al., 2011). However, the same effect was not appreciable in the other oils used in the present study. The ozOO treatment has to some extent, a similar trend of ozSO, but it was not statistical significant. This is partially in agreement

317

ARTICLE IN PRESS

G. Valacchi et al. / International Journal of Pharmaceutics xxx (2013) xxx-xxx



Fig, 4. Effect of ozSO on NFκB activation. (A) In ozSO, an NFκB nuclear positivity (short arrow) is detected, mainly in the upper epidermal layers at d3; it decreases at d7, and almost it disappears on d14; several inflammatory/reparative cells in the dermis also show nuclear positivity to NFκB (long arrow). In the control, no NFκB nuclear positivity is observable in the epidermis, whereas in the dermis positive nuclei are detectable in few inflammatory/reparative cells (long arrow). DAPI was used to blue stain the cell nuclei (original magnification, 400; scale bar: 50 μm). (B) Activation of NF-κB was determined by the translocation in the nucleus of p65 subunit after the ozSO or
 SO treatments of murine skin at d3, d7 and d14. Shown is a representative blot from three independent experiments; β-actin was used as loading control. For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

with the work of Kim et al. (2009) where the therapeutic effects of 333 commercial ozonated olive oil was evaluated in cutaneous wound 334 healing using guinea pigs as an animal model, It is anyway very dif-335 ficult to compare the two studies since in the study from Kim et al. 336 (2009) the ozonation levels of the olive oil is not indicated. As pre-337 viously reported (Valacchi et al., 2011), the peroxidation levels of 338 the ozonated oil is one of the main factors that seems to be involved 339 in the wound healing process. However, the data presented in the 340 present paper only partially confirm our idea of the role of perox-341 ides in skin reparative events. In fact, based in our hypothesis, all 342 the ozonated oils should have had the same effect on WH, since 343 they have the same amount of peroxide. 344

One reason of the different effect on wound healing from the 345 ozonated oils used could be their different composition in PUFA. 346 In fact, there is a clear difference in the level of n-6 and n-3 fatty 347 acids among the three oils, with almost 3 times more n-6 in the 348 sesame oil compared to the OO and LO (Table 1). Different rhe-349 ological behaviours due to inherent differences in the molecular 350 configuration of unsaturated acyl chains after ozonation could pro-351 vide further explanation for the differences observed on the wound 352 parameters. It has been demonstrated that ozone easily reacts with 353 PUFA of the vegetable oils firstly forming the trioxolane moiety that 354 when an ozonated oil is applied on the wound containing exudates 355 it decomposes releasing its derivative products such as aldehydes 356 and H₂O₂ (Travagli et al., 2009, 2010). In our study we have shown 357 that the levels of 4-HNE in the skin treated with ozSO was higher 358 compared to the other treatments and this could be the conse-359 quence of the higher level of n-6 present in SO compared to OO 360 and LO (Crews et al., 2006; European Pharmacopoeia, 2010, 2011). 361

In fact, 4-HNE derives mainly from the oxidation of n-6 PUFA, essen-362 tially arachidonic and linoleic acid (Poli et al., 2008). 4-HNE is an 363 unusual compound containing three functional groups that in many 364 cases act in concert explaining its high reactivity and because of its 365 electrophilic nature, 4-HNE can form adducts with cellular protein 366 nucleophiles. Indeed, the reactivity of 4-HNE explains its potential 367 involvement in the modulation of enzymes activity, signal trans-368 duction and gene expression (Poli et al., 2008). 4-HNE is also an 369 efficient cell signalling molecule able to modulate the expression 370 of several genes and therefore it may influence important cellular 371 functions such as cell growth, differentiation, and angiogenesis all 372 process involved in wound healing (Leonarduzzi et al., 2004). An 373 increasing bulk of literature indicates that 4-HNE, depending on 374 its concentration, can have opposite effects: activating or inhibit-375 ing stress response mechanisms, such as mitogen-activated protein 376 kinases (MAPKs), modulating redox-sensitive transcription factors 377 such as NF_KB (Amma et al., 2005; Lee et al., 2008). The activation 378 of NFkB observed in the ozSO treated skin is most likely a conse-379 quence of the increased levels of 4-HNE. Among its several function, 380 NF κ B has been shown to be critical in modulating the healing 381 process in injury (Sen and Roy, 2008). Fakhrzadeh et al. (2004) 382 have reported that overexpression of superoxide dismutase not 383 only prevents O₃ related changes in bronchoalveolar lavage fluid 384 protein, macrophage number and hydroxyalkenal levels but also 385 O3-dependent activation of NFkB. These results associate therefore O₃ exposure to NFkB activation and parallel with our pioneering 387 work where we have demonstrated that ozone exposure is able to 388 activate NFkB in cutaneous tissue using the same animal model 389 (SKH1 mice) (Valacchi et al., 2004). It should be mentioned that 390

8

391

392

393

394

395

396

397

398

399

400

401

ARTICLE IN PRESS

G. Valacchi et al. / International Journal of Pharmaceutics xxx (2013) xxx-xxx



Fig. 5. Wound repair was analyzed by using Picrosirius staining. Under polarization microscopy, type I fibres are radiant yellow/orange while type III fibres are green. In the ozSO at d3, as well as at d7, the extent of injury (the area starting from the arrow) is less than in controls (SO and control), wound healing occurring more rapidly in ozSO (the white star symbol indicates the normal dermis, in which collagen fibres are stained in red by Picrosirius observed under light microscopy). At d14, in ozSO the dermis returns to be normal (green star symbol), with well-oriented Picrosirius red-stained collagen fibres. Instead, in the control the reparative phenomena are still evident, there are less collagen fibres than in ozSO (original magnification, × 100; scale bar: 200 µm). For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

the dose-dependent effect relationship between level of oxidative stress (4-HNE) and NFκB exhibits a biphasic profile, consistent with the hormesis dose, response model (Calabrese, 2013): while a moderate levels of α , β -unsaturated aldehydes can activate NFκB through an IκB kinase independent mechanism, extremely high levels of α , β -unsaturated aldehydes have been shown to inhibit NFκB activation by blocking IκB α phosphorylation (Byun et al., 2002; Valacchi et al., 2005).

This could be the case for the delayed effect of highly ozSO (peroxidation levels of circa 3000), where the high levels of peroxidation had an inhibitory effect on the wound healing process. Such an effect could be a consequence of NF κ B inhibition (Valacchi et al., 2011). This hypothesis would parallel previous work where exposure of aged mice to ozone would delay wound healing process because the high oxidative levels derived from both ozone exposure and the increased physiological oxidative stress present in the old animals (Lim et al., 2006).

NFκB activation has been also associated with VEGF and PCNA induction and this explains the increased levels of both proteins in the tissue treated with ozSO. In general, the events that characterize the wound healing process can be summarized as the inflammation stage (Phase I); proliferation with synthesis of extracellular matrix 412

G. Valacchi et al. / International Journal of Pharmaceutics xxx (2013) xxx–xxx

(Phase II) and the remodelling stage (Phase III) in which the scar 413 is transformed to the healed wound. During this well coordinated 414 process there is the appearance of several cell types that follow 415 a specific order (platelets, neutrophils, macrophages, lymphocytes, 416 fibroblasts and angiogenesis) and it has been shown that VEGF plays 417 a critical role the regulation of all these events (Bates and Jones, 418 2003; Bao et al., 2009). Interestingly, although only the treatment 410 with ozSO had a significant effect on the wound closure rate, VEGF 420 was up-regulated also by ozOO treatment. This is in line with the 421 previous work by Kim et al. (2009) where ozOO was able to clearly 422 induce VEGF together with other growth factors (PDGF, TGFB and 423 FGF). It is possible that, in our experimental procedure, the treat-424 ment with ozOO is not able to activate all the pathways involved in 425 the wound healing process. In addition, the increased level of VEGF 426 did not parallel the eventual new vessels formation in the tissue, 427 since there was not any difference in terms of angiogenesis with 428 the control group. This discrepancy could be a consequence of the 429 low level n-6 present in olive oil, and therefore the lower level of 430 the "bioactive products" such as 4-HNE involved in this process. In 431 fact, only the treatment with ozSO was able to significantly activate 432 the proliferative biomarker PCNA. Re-epithelialization is a process 433 434 of restoring the epidermis and consists of proliferation and migration of cutaneous cells to the wound edge to resurface epidermal 435 defects (Li et al., 2007) and therefore the induction of PCNA explains 436 the increased wound healing rate observed in the animal treated 437 with ozonated sesame oil. In addition, it has been shown a direct 438 correlation between NFkB activation and PCNA regulation and this 439 is in support of our findings where we have detected an increased 440 NFkB nuclear translocation in the tissue treated with ozSO. 441

In conclusion, these results demonstrate that application of ozonated oils, especially the ozonated sesame oil with a peroxidation value around 1500, can improve acute cutaneous wound repair in the SKH1 murine model by affecting the early phases of the process. Taken together, the data within presented suggest that topical application of specific vegetable matrices in the form of ozone derivatives may be considered as an alternative therapeutic modality to enhance cutaneous wound healing.

Acknowledgements

The study was partially supported by MIUR (VG "Programma Rientro Cervelli") and by FAR 2011–2012 of University of Ferrara,

References

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

- Ali, F., Sultana, S., 2012. Repeated short-term stress synergizes the ROS signalling through up regulation of NFkB and iNOS expression induced due to combined exposure of trichloroethylene and UVB rays. Mol. Cell Biochem. 360, 133-145.
- Amma, H., Naruse, K., Ishiguro, N., Sokabe, M., 2005. Involvement of reactive oxygen species in cyclic stretch-induced NF-kappa activation in human fibroblast cells. Br. J. Pharmacol. 145, 364-373.
- Bao, P., Kodra, A., Tomic-Canic, M., Golinko, M.S., Ehrlich, H.P., Brem, H., 2009. The role of vascular endothelial growth factor in wound healing. J. Surg. Res. 153. 347-358.
- Barrientos, S., Stojadinovic, O., Golinko, M.S., Brem, H., Tomic-Canic, M., 2008 Growth factors and cytokines in wound healing. Wound Repair Regen. 16, 585-601.
- Bates, D.O., Jones, R.O., 2003. The role of vascular endothelial growth factor in wound healing. Int. J. Low Extrem. Wounds 2, 107-120.
- Byun, M.S., Jeon, K.I., Choi, J.W., Shim, J.Y., Jue, D.M., 2002. Dual effect of oxidative stress on NF-kappaB activation in HeLa cells. Exp. Mol. Med. 34, 332-339.
- Calabrese, E.L. 2013, Historical foundations of wound healing and its potential for acceleration: dose-response considerations. Wound Repair Regen, 21, 180–193.
- Crews, C., Hough, P., Brereton, P., Godward, J., Lees, M., Guiet, S., Winkelmann, W., 2006. Ouantitation of the main constituents of some authentic sesame seed oils of different origin. J. Agric. Food Chem. 54, 6266-6270.
- de Melo, C.M., Porto, C.S., Melo-Júnior, M.R., Mendes, C.M., Cavalcanti, C.C., Coelho, L.C., Porto, A.L., Leão, A.M., Correia, M.T., 2011. Healing activity induced by

Cramoll 1,4 lectin in healthy and immunocompromised mice. Int. J. Pharm. 408, 113-119

- Edwards, R., Harding, K.G., 2004. Bacteria and wound healing. Curr. Opin. Infect. Dis. 17, 91-96.
- European Pharmacopoeia, 2010. Linseed Oil, Virgin, seventh ed. Directorate for the Quality of Medicines and HealthCare, Strasbourg. Druckerei C.H. Beck, Nördlingen, Germany, pp. 2898-2899.
- European Pharmacopoeia, 2011. Olive oil, Refined, seventh ed. Suppl. 7.2. Directorate for the Quality of Medicines and HealthCare, Strasbourg. Druckerei C.H. Beck, Nördlingen, Germany, pp. 3689.
- Kim, H.S., Noh, S.U., Han, Y.W., Kim, K.M., Kang, H., Kim, H.O., Park, Y.M., 2009. Therapeutic effects of topical application of ozone on acute cutaneous wound healing. J. Korean Med. Sci. 24, 368-374.
- Fakhrzadeh, L., Laskin, J.D., Laskin, D.L., 2004. Ozone-induced production of nitric oxide and TNF-alpha and tissue injury are dependent on NF-kappaB p50. Am. J. Physiol. Lung Cell Mol. Physiol. 287, L279-L285.
- Lee, S.J., Seo, K.W., Yun, M.R., Bae, S.S., Lee, W.S., Hong, K.W., Kim, C.D., 2008. 4-HNE MMP-2 production in vascular smooth muscle cells via mitochondrial ROSmediated activation of the Akt/NF-kappaB signaling pathways. Free Radic. Biol. Med. 45, 1487-1492.
- Leonarduzzi, G., Robbesyn, F., Poli, G., 2004. Signaling kinases modulated by 4hydroxynonenal. Free Radic. Biol. Med. 37, 1694-1702.
- Li, J., Chen, J., Kirsner, R., 2007. Pathophysiology of acute wound healing. Clin. Dermatol. 25, 9-18.
- Lim, Y., Phung, A.D., Corbacho, A.M., Aung, H.H., Maioli, E., Reznick, A.Z., Cross, C.E., Davis, P.A., Valacchi, G., 2006. Modulation of cutaneous wound healing by ozone: differences between young and aged mice. Toxicol. Lett. 160, 127-134.
- Meng, Z., Lou, S., Tan, J., Xu, K., Jia, Q., Zheng, W., 2012. Nuclear factor-kappa B inhibition can enhance apoptosis of differentiated thyroid cancer cells induced by ¹³¹I. PLoS ONE 7, e33597. Moldovan, G.L., Pfander, B., Jentsch, S., 2007. PCNA, the maestro of the replication
- fork. Cell 129, 665-679.
- Moseley, R., Hilton, J.R., Waddington, R.J., Harding, K.G., Stephens, P., Thomas, D.W., 2004. Comparison of oxidative stress biomarker profiles between acute and chronic wound environments. Wound Repair Regen. 12, 419-429.
- Poli, G., Schaur, R.I., Siems, W.G., Leonarduzzi, G., 2008. 4-hydroxynonenal: a membrane lipid oxidation product of medicinal interest. Med. Res. Rev. 28, 569–631.
- Prvor, W.A., Church, D.F., 1991, Aldehydes, hydrogen peroxide, and organic radicals as mediators of ozone toxicity. Free Radic. Biol. Med. 11, 41-46.
- Ruef, J., Moser, M., Bode, C., Kübler, W., Runge, M.S., 2001. 4-hydroxynonenal induces apoptosis, NF-kappaB-activation and formation of 8-isoprostane in vascular smooth muscle cells. Basic Res. Cardiol. 96, 143-150.
- Sega, A., Zanardi, I., Chiasserini, L., Gabbrielli, A., Bocci, V., Travagli, V., 2010. Properties of sesame oil by detailed ¹H and ¹³C NMR assignments before and after ozonation and their correlation with iodine value, peroxide value, and viscosity measurements, Chem, Phys. Lipids 163, 148-156.
- Sen, C.K., Roy, S., 2008. Redox signals in wound healing. Biochim. Biophys. Acta 1780, 1348-1361.
- Sticozzi, C., Belmonte, G., Pecorelli, A., Arezzini, B., Gardi, C., Maioli, E., Miracco, C., Toscano, M., Forman, H.J., Valacchi, G., 2012. Cigarette smoke affects keratinocytes SRB1 expression and localization via H_2O_2 production and HNE protein adducts formation. PLoS ONE, http://dx.doi.org/10.1371/ ournal.pone.0033592, e33592.
- Travagli, V., Zanardi, I., Bocci, V., 2009. Topical applications of ozone and ozonated oils as anti-infective agents: an insight into the patent claims. Recent Pat. Antiinfect. Drug Discov. 4, 130–142. Avagli, V., Zanardi, I., Valacchi, G., Bocci, V., 2010. Ozone and ozonated oils in skin
- diseases: a review. Mediat, Inflamm., http://dx.doi.org/10.1155/2010/610418, 610418
- Uchida, Y., Ohba, K., Yoshioka, T., Irie, K., Muraki, T., Maru, Y., 2004. Cellular carbonyl stress enhances the expression of plasminogen activator inhibitor-1 in rat white adipocytes via reactive oxygen species-dependent pathway. J. Biol. Chem, 279, 4075-4083.
- Valacchi, G., Pagnin, E., Corbacho, A.M., Olano, E., Davis, P.A., Packer, L., Cross, C.E., 2004. In vivo ozone exposure induces antioxidant/stress-related responses in murine lung and skin. Free Radic. Biol. Med. 36, 673-681.
- Valacchi, G., Pagnin, E., Phung, A., Nardini, M., Schock, B.C., Cross, C.E., van der Vliet, A. 2005. Inhibition of NFkappaB activation and IL-8 expression in human bronchial epithelial cells by acrolein. Antioxid. Redox Signal. 7, 25-31.
- Valacchi, G., Pecorelli, A., Mencarelli, M., Carbotti, P., Fortino, V., Muscettola, M., Maioli, E., 2009. Rottlerin: a multifaced regulator of keratinocyte cell cycle. Exp. Dermatol. 18, 516-521.
- Valacchi, G., Lim, Y., Belmonte, G., Miracco, C., Zanardi, I., Bocci, V., Travagli, V., 2011. Ozonated sesame oil enhances cutaneous wound healing in SKH1 mice. Wound Repair Regen. 19, 107-115.
- Valacchi, G., Zanardi, I., Sticozzi, C., Bocci, V., Travagli, V., 2012. Emerging topics in cutaneous wound repair. Ann. N. Y. Acad. Sci. 1259, 136-144.
- Werner, S., Grose, R., 2003. Regulation of wound healing by growth factors and cytokines. Physiol. Rev. 83, 835-870.
- Zanardi, I., Travagli, V., Gabbrielli, A., Chiasserini, L., Bocci, V., 2008. Physico-chemical characterization of sesame oil derivatives. Lipids 43, 877-886.

476

478

479

<mark>48</mark>0

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531 532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556