

SHORT REPORT

Open Access

# Simple clinical indicators for early psoriatic arthritis detection

Francesco Caso<sup>1</sup>, Luisa Costa<sup>1,2</sup>, Mariangela Atteno<sup>2</sup>, Antonio Del Puente<sup>2</sup>, Luca Cantarini<sup>3</sup>, Ennio Lubrano<sup>4</sup> and Raffaele Scarpa<sup>2\*</sup>

## Abstract

**Background:** Diagnosis of psoriatic arthritis (PsA), in a period of 12 months from the onset of the first articular episode, permits of identifying the early form defined as “early PsA”. The recognition of the disease in this phase leads to better outcome.

The aim of this study was to identify peculiar clinical and/or laboratory findings that could be useful for the diagnosis of “early PsA”.

**Findings:** Thirty-five patients with early onset of arthritis were observed. The following data were collected for each patient: family and personal history, physical examination, tender and swollen joint counts (TJC, SJC), tender enthesal count, presence of dactylitis and low back pain (LBP), and laboratory tests.

Among the 35 total patients, 24 showed skin and/or nail psoriasis or a family history of psoriasis.

The remaining 11 patients showed absence of concomitant or previous psoriasis and/or familiarity for psoriasis. The comparison between the two groups showed that patients with psoriasis had a significant presence of LBP, dactylitis and enthesitis than patients with psoriasis.

**Conclusions:** The study confirms that the distinctive clinical findings of PsA is psoriasis, but also LBP, dactylitis and enthesitis have a relevant role in early identification. A low number of SJC and TJC are frequently observed in early phases of PsA than in other forms of early arthritis. These aspects could be mostly helpful when psoriasis is not detected or can follow arthritis in absence of familiar positivity, making difficult PsA diagnosis.

In conclusion, careful medical history, clinical examination and first-level laboratory investigations are useful to characterize early phases of PsA.

**Keywords:** Early psoriatic arthritis; Psoriasis; Low-back pain; Dactylitis; Enthesitis

## Introduction

Psoriatic arthritis (PsA) is a chronic inflammatory arthropathy, belonging to the spondyloarthritic area, usually associated with skin and/or nail psoriasis or with its familiarity (Moll & Wright 1973). Other clinical sites such as the bowel, eye and cardiovascular system be variably involved (Scarpa et al. 2000; Niccoli et al. 2012; Shang et al. 2012; Costa et al. 2012).

The progress in the knowledge of new imaging techniques has changed the diagnostic approach to this condition (Soscia et al. 2012; Tan et al. 2014; Poggenborg et al. 2014; Gladman 2012), even if final diagnosis mainly

relies on clinical evaluation performed by a rheumatologist. It usually refers to a characteristic presentation of joints, spine and enthesitis inflammatory findings, a negative rheumatoid factor in presence of psoriasis or its familiarity (Duarte et al. 2012).

Diagnosis of PsA, in a period of 12 months from the onset of the first articular episode, permits of identifying the early form defined as “early PsA”. The recognition of the disease in this phase leads to better outcome. However, in the context of clinical practice, in early phases it is complex to rule out other inflammatory diseases, such as rheumatoid arthritis, undifferentiated arthritis and other spondyloarthritides (Gladman 2012).

In 2006, a set of classification criteria was developed by the CASPAR (Classification criteria for Psoriatic ARthritis) Study Group, with the aim of identifying

\* Correspondence: rscarpa@unina.it

<sup>2</sup>Rheumatology Unit, Department of Clinical Medicine and Surgery, University Federico II, via S. Pansini 5, 80131 Naples, Italy

Full list of author information is available at the end of the article

standardized and homogeneous cohorts of PsA patients in clinical research context (Duarte et al. 2012). Although CASPAR Criteria do not represent diagnostic criteria, these have had a wide consensus, due to high specificity (98.7%) and good sensitivity (91.4%), confirmed also retrospectively (Taylor et al. 2006; Tillett et al. 2012). Furthermore, CASPAR criteria have been applied in some studies on early PsA showing good sensitivity (88.7%) and high specificity (99.1%), but data are still few (van den Berg et al. 2012; D'Angelo et al. 2009). While a distinctive clinical finding of PsA is psoriasis, dactylitis, enthesitis and inflammatory low-back pain should be present without a pathognomonic role. In Addition, no laboratory findings can be helpful in addressing PsA diagnosis. The aim of this study was to identify peculiar clinical and/or laboratory findings that could be useful for the diagnosis of "early PsA".

### Patients and methods

In 7-month period, 35 patients (M/W: 7/28; mean age: 47.0 years, range: 17–78 years) with early onset of arthritis (within 12 weeks of onset), attending Rheumatology Unit of University Federico II of Naples, were observed. The average onset of arthritis was 5.6 months (range 2–12 months). The following data were collected for each patient: family and personal history, including familiar and/or previous psoriasis, physical examination, tender and swollen joint counts (TJC, SJC), tender enthesal count, presence of dactylitis and low back pain. Evaluation of rheumatoid factor (RF), anti-cyclic citrullinated peptide antibodies (anti-CCP) and inflammatory indices, including erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP), were performed.

Written informed consent was obtained from the patient for the publication of this report.

Statistical analysis was performed using the SPSS software, version 18 (SPSS inc, Chicago, Ill). All variables were normally distributed. Analysis of variance (ANOVA) was used to assess differences between group means.

### Results

Among the 35 total patients, twenty-four showed skin and/or nail psoriasis or a family history of psoriasis (M/W: 5/19, mean age 44.9 years, range 17–73 years; mean duration of articular disease 5.8 months, range 2–12). The remaining eleven patients showed absence of concomitant or previous psoriasis and/or familiarity for psoriasis (M/W: 2/9, mean age 51.5 years, range 28–78 years; mean disease duration 5.0 months, range 2–12).

The comparison between the two groups (Table 1) showed that patients with psoriasis had a significant presence of inflammatory back pain, dactylitis and enthesitis (respectively  $p < 0.001$ ,  $p < 0.001$  and  $p < 0.01$ ), whereas patients without psoriasis showed a greater TJC ( $p < 0.001$ ),

**Table 1 Clinical and laboratory findings of early arthritis patients with and without psoriasis**

	Early arthritis patients with psoriasis (n. 24)	Early arthritis patients without psoriasis (n.11)	<i>p</i>
TJC	6.0 (0–28)	13.45 (8–24)	<0.05
SJC	0.9 (0–5.0)	2.54 (0–10)	<0.05
LBP, n (%)	16; 66.6%	0	<0.001
Dactylitis, n (%)	11; 45.8%	0	<0.001
Enthesitis, n (%)	5; 20.8%	0	<0.01
ESR > 15 mm/1 h, n (%)	6; 25%	11; 100%	ns
CRP > 0.5 mg/dL, n (%)	6; 25%	11; 100%	ns
RF positivity, n (%)	0	9; 81.8%	0.01
Anti-CCP positivity, n (%)	0	4; 36.3%	ns
Uveitis, n (%)	3; 12.5%	0	ns

Data expressed as mean ± range, unless otherwise indicated.

SJC ( $p < 0.001$ ), and RF positivity ( $p < 0.01$ ) than patients with psoriasis. In the group of patients with psoriasis, the application of CASPAR criteria led to a PsA diagnosis in all the patients, while none of the patients in the group without psoriasis was satisfying these criteria.

### Conclusions

For a long time, PsA has been considered a low grade inflammatory condition and only most recently increasing data have provided evidence on the severe impact of this disease, establishing at same time the importance of an early diagnosis and treatment (Gladman 2012; Scarpa et al. 2011).

The CASPAR criteria consist of established inflammatory articular disease with at least a sum of three points from the following: current psoriasis (2 points), a history of psoriasis (1 point), a family history of psoriasis (unless current psoriasis was present or there was a history of psoriasis; 1 point), dactylitis (1 point), juxta-articular new bone formation (1 point), RF negativity (1 point), and nail dystrophy (1 point) (Taylor et al. 2006).

In this study, co-occurrence of inflammatory back pain and dactylitis were present respectively in above 46% and 66% of patients with psoriasis and were very helpful in addressing early PsA diagnosis. Enthesitis was detected in above 21% of patients with psoriasis.

Patients with psoriasis showed a lower number of TJC and SJC, than patients without psoriasis; previous episode of uveitis was referred in three of 24 patients with psoriasis and in no one of patients without psoriasis.

With regard to laboratory findings, all the patients without psoriasis showed increased ESR and C-RP, RF and anti-CCP positivity was found in above 82% and 36% of them. Differently, only 25% of patients with psoriasis showed increase of ESR and CRP and no one RF and anti-CCP positivity.

This study confirms that the distinctive clinical findings of PsA is psoriasis, but also inflammatory low-back pain, dactylitis and enthesitis have a relevant role in early identification. A low number of SJC and TJC, as well anti-CCP and RF negativity and no increase of ESR and C-RP are most frequently observed in early phases of PsA than in other forms of early arthritis. These aspects could be mostly helpful when psoriasis is not detected or can follow arthritis in absence of familiar positivity, making difficult PsA diagnosis.

The study shows several limitations: mainly the number of patients which is low and for this reason the results need to be supported by studies on larger populations.

Although the progress in the knowledge of new imaging techniques has changed the diagnostic approach to PsA (Soscia et al. 2012; Tan et al. 2014; Poggenborg et al. 2014; Gladman 2012), but PsA diagnosis still relies mainly on clinical evaluation.

Therefore, the results of the present observation may be interesting from practical point of view. The key message is that careful medical history, clinical examination and first-level laboratory investigations such as ESR, CRP, anti-CCP and FR are reasonably useful to characterize early phases of PsA. Delay in PsA diagnosis significantly contributes to poor outcome of the patient, so early identification of simple and helpful indicators could allow for the characterization of “early PsA”.

#### Abbreviations

TJC: Tender joint counts; SJC: Swollen joint counts; LBP: Low-back pain; ESR: Erythrocyte sedimentation rate; CRP: C-reactive protein; RF: Rheumatoid factor; anti-CCP: anti-cyclic citrullinated peptide antibodies.

#### Competing interests

The authors declare that they have no competing interest.

#### Authors' contributions

RS conceived of the study, provided medical data and drafted the manuscript. FC participated in the design of the study, and drafted the manuscript. LC participated in the design of the study, and drafted the manuscript. MA interpreted the data and drafted the manuscript. EL performed the statistical analysis and revised the manuscript critically. LC performed the statistical analysis and revised the manuscript critically. ADP interpreted the data and drafted the manuscript. All authors read and approved the final manuscript.

#### Author details

<sup>1</sup>Rheumatology Unit, Department of Medicine DIMED, University of Padova, Via Giustiniani 2, 35128 Padova, Italy. <sup>2</sup>Rheumatology Unit, Department of Clinical Medicine and Surgery, University Federico II, via S. Pansini 5, 80131 Naples, Italy. <sup>3</sup>Interdepartmental Research Center of Systemic Autoimmune and Autoinflammatory Diseases, Rheumatology Unit, Policlinico Le Scotte, University of Siena, Viale Bracci 1, 53100 Siena, Italy. <sup>4</sup>Academic Rheumatology Unit, Department of Medicine and Health Sciences, University of Molise, Campobasso, Italy.

Received: 2 December 2014 Accepted: 3 December 2014

Published: 22 December 2014

#### References

Costa L, Caso F, D'Elia L, Attenu M, Peluso R, Del Puente A, Strazzullo P, Scarpa R (2012) Psoriatic arthritis is associated with increased arterial stiffness in the

absence of known cardiovascular risk factors: a case control study. *Clin Rheumatol* 31:711–715

D'Angelo S, Mennillo GA, Cutro MS, Leccese P, Nigro A, Padula A, Olivieri I (2009) Sensitivity of the classification of psoriatic arthritis criteria in early psoriatic arthritis. *J Rheumatol* 36:368–370

Duarte GV, Faillace C, Freire De Carvalho J (2012) Psoriatic arthritis. *Best Pract Res Clin Rheumatol* 26:147–156

Gladman DD (2012) Early psoriatic arthritis. *Rheum Dis Clin North Am* 38(2):373–386

Moll JM, Wright V (1973) Psoriatic arthritis. *Semin Arthritis Rheum* 3:55–78

Niccoli L, Nannini C, Cassarà E, Kaloudi O, Susini M, Lenzetti I, Cantini F (2012)

Frequency of iridocyclitis in patients with early psoriatic arthritis: a prospective, follow up study. *Int J Rheum Dis* 15:414–418

Poggenborg RP, Eshed I, Ostergaard M, Sørensen JJ, Møller JM, Madsen OR, Pedersen SJ (2014) Enthesitis in patients with psoriatic arthritis, axial spondyloarthritis and healthy subjects assessed by 'head-to-toe' whole-body MRI and clinical examination. *Ann Rheum Dis*, doi:10.1136/annrheumdis-2013-204239

Scarpa R, Manguso F, D'Arienzo A, D'Armiendo FP, Astarita C, Mazzacca G, Ayala F (2000) Microscopic inflammatory changes in colon of patients with both active psoriasis and psoriatic arthritis without bowel symptoms. *J Rheumatol* 27:1241–1246

Scarpa R, Atteno M, Lubrano E, Provenzano G, D'Angelo S, Spadaro A, Costa L, Olivieri I (2011) The effectiveness and safety of TNF-alpha blockers in the treatment of early psoriatic arthritis: an Italian multicentre longitudinal observational pilot study. *Clin Rheumatol* 30:1063–1067

Shang Q, Tam LS, Sanderson JE, Sun JP, Li EK, Yu CM (2012) Increase in ventricular-arterial stiffness in patients with psoriatic arthritis. *Rheumatology (Oxford)* 51:2215–2223

Soscia E, Sirignano C, Catalano O, Attenu M, Costa L, Caso F, Peluso R, Bruner V, Aquino MM, Del Puente A, Salvatore M, Scarpa R (2012) New developments in magnetic resonance imaging of the nail unit. *J Rheumatol Suppl* 89:49–53

Tan AL, Fukuba E, Halliday NA, Tanner SF, Emery P, McGonagle D (2014)

High-resolution MRI assessment of dactylitis in psoriatic arthritis shows flexor tendon pulley and sheath-related enthesitis. *Ann Rheum Dis*, doi:10.1136/annrheumdis-2014-205839

Taylor W, Gladman D, Helliwell P, Marchesoni A, Mease P, Mielants H (2006) Classification criteria for psoriatic arthritis: development of new criteria from a large international study. *Arthritis Rheum* 54:2665–2673

Tillett W, Costa L, Jadon D, Wallis D, Cavill C, McHugh J, Korendowych E, McHugh N (2012) The CIASsification for psoriatic ARthritis (CASPAR) criteria - a retrospective feasibility, sensitivity, and specificity study. *J Rheumatol* 39:154–156

van den Berg R, van Gaalen F, van der Helm-van MA, Huizinga T, van der Heijde D (2012) Performance of classification criteria for peripheral spondyloarthritis and psoriatic arthritis in the Leiden early arthritis cohort. *Ann Rheum Dis* 71:1366–1369

doi:10.1186/2193-1801-3-759

Cite this article as: Caso et al.: Simple clinical indicators for early psoriatic arthritis detection. *SpringerPlus* 2014 **3**:759.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)