

Nucleotide Sequence Analysis of Integrative Conjugative Element Tn 5253 of *Streptococcus pneumoniae*

Francesco Iannelli, Francesco Santoro, Marco R. Oggioni
and Gianni Pozzi

Antimicrob. Agents Chemother. 2014, 58(2):1235. DOI:
10.1128/AAC.01764-13.

Published Ahead of Print 2 December 2013.

Updated information and services can be found at:
<http://aac.asm.org/content/58/2/1235>

SUPPLEMENTAL MATERIAL	<i>These include:</i> Supplemental material
REFERENCES	This article cites 29 articles, 18 of which can be accessed free at: http://aac.asm.org/content/58/2/1235#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Nucleotide Sequence Analysis of Integrative Conjugative Element Tn5253 of *Streptococcus pneumoniae*

Francesco Iannelli, Francesco Santoro, Marco R. Oggioni,* Gianni Pozzi

Laboratory of Molecular Microbiology and Biotechnology (LAMMB), Department of Medical Biotechnologies, University of Siena, Siena, Italy

Conjugative transposon Tn5253, an integrative conjugative element (ICE) of *Streptococcus pneumoniae* carrying the *cat* and *tet(M)* genes, was shown to be 64,528 bp in size and to contain 79 open reading frames, of which only 38 could be annotated. Two distinct genetic elements were found integrated into Tn5253: Tn5251 (18,033 bp), of the Tn916-Tn1545 family of ICEs, and Ω cat(pC194) (7,627 bp), which could not conjugate but was capable of intracellular mobility by excision, circularization, and integration by homologous recombination. The highest conjugation frequency of Tn5253 was observed when *Streptococcus pyogenes* was the donor (6.7×10^{-3} transconjugants/donor).

Comparative analysis of sequenced *Streptococcus pneumoniae* genomes indicates that many genes clustered in integrative and conjugative elements (ICEs) are responsible for pneumococcal genome evolution (1). ICEs, which comprise conjugative transposons, are found integrated into bacterial genomes and can be excised to form covalently closed circular intermediates that can either integrate within the cell at a different genomic site or move to a new bacterial host upon conjugative transfer (2). One of the first ICEs to be identified was Tn5253, originally called Ω (*cat-tet*) BM6001, which is a conjugative transposon found in the chromosome of a serogroup 19 clinical isolate of *S. pneumoniae* (3, 4). Tn5253 was shown to be a genetic element conferring resistance to chloramphenicol and tetracycline that is capable of conjugal transfer and site-specific chromosomal integration in different Gram-positive bacteria belonging to the genera *Streptococcus* and

Enterococcus (5–7). Tn5253 was described as a composite conjugative transposon since it carried a complete copy of another ICE,

Received 16 August 2013 Returned for modification 17 September 2013

Accepted 27 November 2013

Published ahead of print 2 December 2013

Address correspondence to Gianni Pozzi, gianni.pozzi@unisi.it.

* Present address: Marco R. Oggioni, Department of Genetics, University of Leicester, Leicester, United Kingdom.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AAC.01764-13>.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.01764-13

Tn5253 (64,528 bp)




FIG 1 Structure of *S. pneumoniae* Tn5253. Sequence analysis of Tn5253 showed that it is 64,528 bp in size and contains 79 ORFs. Tn5253 contains two distinct integrated genetic elements, the 18,033-bp-long ICE Tn5251 carrying *tet(M)* and the 7,627-bp-long element Ω cat(pC194) containing *cat*. ORFs and their directions of transcription are represented by arrows, and the annotated ORFs are indicated only by their numbers. The regions corresponding to Tn5251 and Ω cat(pC194) are indicated by solid bars, and their ORFs are represented by thinner arrows. Tetracycline and chloramphenicol resistance genes are represented by red arrows. Chromosomal genes flanking the Tn5253 insertion site are represented by thin gray arrows. The different GC contents of the various regions are indicated by dotted bars. Homologies with pneumococcal chromosomal regions are marked by solid lines. The scale is in kilobases.

TABLE 1 Annotated ORFs of Tn5253

ORF (no. of amino acids) ^{a,b}	Annotation and comments; reference	Pfam domain(s) ^c (amino acids) [E value(s)]	Homologous protein ID/origin	No. of amino acids identical/total (%)-no. similar/total (%)	E value ^d
<i>repA/orf2</i> (259)	Plasmid replication initiator protein A, N terminal; 20	RepA_N (13-88) [4.1e-28]			
<i>orf4</i> (452)	DNA cytosine methyltransferase; 21				
<i>orf8</i> (195)	Protease, putative	Abi-CAAX protease self-immunity (103-188) [1.6e-13]			
<i>orf9</i> (196)	Abortive infection protein AbiEi; 22	DUF4095 (29-146) [2.1e-23]	AAB52382.1/pNP40 <i>L. lactis</i>	53/203 (26)-91/203 (44)	6e-06
<i>orf10</i> (278)	Abortive infection protein AbiEii; 22	DUF1814 (37-260) [2.1e-19]	AAB52383.1/pNP40 <i>L. lactis</i>	81/279 (29)-136/279 (48)	3e-18
<i>traG/orf13</i> (625)	TraG-like protein, involved in bacterial conjugal transfer; 23	T4SS-DNA_transf (133-574) [1.1e-66]			
<i>orf15</i> (284)	Membrane protein, putative	TrbL (123-269) [0.00026]	YP_195781.1 (<i>prgH</i>)/pCF10 <i>E. faecalis</i>	88/261 (33)-150/261 (57)	5e-24
<i>orf16</i> (119)	Type IV secretion system protein	PrgI (4-95) [2.1e-22]			
<i>orf17</i> (785)	Type IV secretion protein, VirB4, ATPase	AAA_10 (449-736) [3.8e-44]	YP_195783.1 (<i>prgI</i>)/pCF10 <i>E. faecalis</i>	284/750 (37)-448/750 (59)	5e-134
<i>orf18</i> (937)	Peptidoglycan hydrolase, putative	CHAP (805-926) [8.1e-30]			
<i>orf20</i> (64)	ISSt5, transposase, <i>IS110</i> family, truncated		YP_139721.1/ <i>S. thermophilus</i>	43/64 (67)-47/64 (73)	1e-13
<i>orf43</i> (243)	ABC transporter: ATP-binding protein, putative	ABC_tran (27-173) [2.8e-30]			
<i>orf44</i> (438)	ABC transporter: permease (10 transmembrane helices predicted)	FtsX (55-176) [4.7e-07]			
<i>orf47</i> (2028)	Restriction-modification protein, putative	Methyltransf_26 (504-615) [1e-17], SNF2_N (1253-1600) [2.3e-14], Helicase_C (1698-1768) [2.2e-05]			
<i>orf52</i> (361)	DNA primase, putative	DUF3991 (113-198) [2e-13], Toprim_2 (203-329) [4.7e-10]			
<i>pezA/orf56</i> (158)	Antitoxin (TA system); 18				
<i>pezT/orf57</i> (252)	Toxin (TA system); 18				
<i>cat/orf60</i> (216)	CAT ^e	CAT (1-205) [2.8e-95]	NP_040437.1/pC194 <i>S. aureus</i>	216/216 (100)	
<i>orf63</i> (281)	Plasmid replication protein	Rep_1 (54-250) [1e-48]	NP_040435.3/pC194 <i>S. aureus</i>	194/195 (99)-195/195 (100)	6e-111
<i>nplT/orf65</i> (245)	Truncated neopullulanase (α -amylase); 24	Alpha-amylase (2-141) [5.5e-25]			
<i>pezA/orf67</i> (158)	Antitoxin (TA system); 18				
<i>pezT/orf68</i> (256)	Toxin (TA system); 18				
<i>umuD/orf69</i> (227)	UmuD MucA homolog; 25	HTH_3 (7-61) [2.3e-17], Peptidase_S24 (143-205) [8e-12]			
<i>umuC/orf70</i> (471)	SOS response UmuC protein; 25	IMS (17-207) [2.2e-17], IMS_HHH (223-255) [5.1e-06], IMS_C (268-382) [5.9e-12]			
<i>orf74</i> (121)	Bacterial mobilization protein, putative; 26	MobC (60-107) [5.4e-15]			
<i>orf75</i> (609)	Relaxase; 27	Relaxase (12-264) [1.6e-73]			
<i>orf76</i> (75)	Transcriptional regulator, putative; 28	HTH_3 (5-60) [2.1e-15]			
<i>xis/orf78</i> (77)	DNA excisionase; 29		AAA72427.1/Tn5276 <i>L. lactis</i>	26/60(43)-43/60 (71)	5e-06
<i>int/orf79</i> (502)	DNA integrase; 30	Phage_integrase (188-394) [2.3e-24]	NP_150133.1/bacteriophage MM1 <i>S. pneumoniae</i>	121/413 (29)-193/413 (46)	4e-26

^a The number of amino acids is shown in parentheses.

^b *orf21* through *orf42* correspond to *orf1* to *orf22* of Tn5251 and are annotated elsewhere (11). *orf58* through *orf68* belong to Ω cat(pC194).

^c The numbers in parentheses represent the part of the protein homologous to the Pfam domain.

^d Determined by compositional matrix adjustment.

^e CAT, chloramphenicol acetyltransferase.

named Tn5251, integrated into its sequence (8, 9). Tn5251 is a conjugative transposon of the Tn916-Tn1545 family (10) that was shown to be a fully autonomous ICE capable of conjugal transfer to a variety of bacterial species, including *S. pneumoniae*, *Streptococcus gordonii*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Enterococcus faecalis*, and *Bacillus subtilis* (11). Here we report the complete annotated sequence of Tn5253 and its phenotypic characterization, which allowed the identification of a new genetic element, Ω cat(pC194), integrated into the Tn5253 sequence and containing a linearized copy of *cat*-carrying plasmid pC194 (12).

The nucleotide sequence of Tn5253 (GenBank accession no.

EU351020) was determined by direct sequencing of seven PCR fragments spanning the whole element and obtained from *S. pneumoniae* strain DP1322 (13). PCR primers (see Table S1 in the supplemental material) were designed on the basis of (i) available Tn5253 sequences (8, 14), (ii) Tn5253 chromosomal junction sequences (7), and (iii) the *cat* gene of staphylococcal plasmid pC194 (GenBank accession no. V01277), which is homologous to the *cat* gene of Tn5253 (15). The DNA sequence was confirmed on the other strand by using short PCR fragments as sequencing templates. PCR, DNA sequencing, and sequence analysis were performed as already described (11).

TABLE 2 Conjugation frequencies of Tn5253

Donor strain (species)	Pneumococcal recipient	Transfer frequency ^a	Representative transconjugant
FR67 (<i>S. agalactiae</i>)	FP11 (type 2 rough)	1.1×10^{-6}	FR81
FR43 (<i>S. gordonii</i>)	FP11 (type 2 rough)	8.3×10^{-7}	FR61
FR40 (<i>S. pyogenes</i>)	FP11 (type 2 rough)	6.7×10^{-3}	FR59
FR49 (<i>E. faecalis</i>)	FP11 (type 2 rough)	$<1.8 \times 10^{-8}$	
FR50 (<i>E. faecalis</i>)	FP11 (type 2 rough)	$<2.7 \times 10^{-8}$	
FR22 (<i>S. pneumoniae</i>)	FP11 (type 2 rough)	1.6×10^{-4}	FR58
FR58 (<i>S. pneumoniae</i>)	FP58 (type 2)	2×10^{-5}	FR38
FR58 (<i>S. pneumoniae</i>)	HB565 (type 3)	4.4×10^{-7}	FR39
FR22 (<i>S. pneumoniae</i>)	FP47 (type 4)	3.2×10^{-5}	FR54
FR58 (<i>S. pneumoniae</i>)	FR55 (type 6)	1.3×10^{-5}	FR56

^a The transfer frequency is expressed as the number of transconjugant CFU per donor CFU, and each result is the mean of at least three mating experiments.

Tn5253 was found to be 64,528 bp in size and to contain 79 open reading frames (ORFs). Two distinct genetic elements were found integrated into Tn5253, (i) Tn5251, 18,033 bp long, belonging to the Tn916-Tn1545 family of *tet*(M)-carrying ICEs, and (ii) Ω cat(pC194), 7,627 bp long, an element carrying *cat*-containing plasmid pC194 (Fig. 1). Insertion of Tn5251 and Ω cat(pC194) identified three regions differing in GC content in Tn5253, (i) the left arm, containing ORFs 1 to 20, with a GC content of 36.4%; (ii), the central region, ORFs 43 to 57, with a GC content of 34.7%; and (iii) the right arm, ORFs 69 to 79, with a GC content of 31.3%.

Tn5251 showed the highest GC content (38.8%), the integrated copy of pC194 showed the lowest GC content (28.7%), and the GC content of the rest of Ω cat(pC194) was 35%. (Fig. 1). A further mark of separation between the left arm and the rest of Tn5253 is the presence of *orf20*, coding for a truncated transposase of the *IS110* family, right before the *att* site of Tn5251 (Table 1). Manual homology-based annotation (11) with functional prediction of the hypothetical gene product was possible for only 38 out of the 79 predicted ORFs, whereas 41 ORFs encoded hypothetical proteins that showed no homology to other characterized sequences (Table 1). Tn5253 showed a typical ICE modular organization (16) with an “intercellular mobility module” spanning *orf13* to *orf18* and a “recombination module” spanning *orf74* to *orf79*.

Transconjugants of different bacterial species (*Streptococcus agalactiae*, *Streptococcus gordonii*, *Streptococcus pyogenes*, and *Enterococcus faecalis*) harboring Tn5253 were used as donors in mating experiments with *S. pneumoniae* FP11 (17) as the recipient (Table 2). The frequency of transfer of the element from *S. pyogenes* transconjugant FR40 is, to our knowledge, the highest obtained for an ICE (6.7×10^{-3} transconjugants per donor), while we could not transfer Tn5253 from the enterococcal donor FR49 ($<1.8 \times 10^{-8}$ transconjugants per donor). The frequency of Tn5253 transfer was also investigated in capsulated *S. pneumoniae* recipients with different genetic backgrounds. Conjugal transfer




FIG 2 *S. pneumoniae* Ω cat(pC194) is a 7,627-bp-long element integrated into Tn5253 and contains a copy of plasmid pC194 lacking its replication origin. It is flanked by two 1,169-bp DRs (boxed yellow arrows) that are part of two longer 1,357-bp imperfect DRs. The 5' and 3' ends of the imperfect DRs are represented by red (a and a') and blue (b and b') boxes. Ω cat(pC194) is excised from Tn5253 and produces a circular form and a deletion in Tn5253. Excision of Ω cat(pC194) occurs by recombination between the flanking DRs and leaves in Tn5253 a single DR flanked by a and b', while the DR of the circular form is flanked by a' and b.

of Tn5253 from the standard rough donors to capsulated pneumococci occurred at frequencies ranging from 3.2×10^{-5} to 4.4×10^{-7} transconjugants per donor, while the frequency was 1.6×10^{-4} transconjugants per donor among the rough type 2 derivatives.

Ω cat(pC194) was found to be 7,627 bp long and to carry a copy of *cat*-containing plasmid pC194 harboring a 93-bp deletion involving the replication origin (Fig. 1 and 2). The element in its Tn5253-integrated form was flanked by two 1,169-bp direct repeats (DRs) corresponding to *orf56-orf57* and *orf67-orf68* (Fig. 2). The 1,169-bp DRs were part of two longer 1,357-bp imperfect DRs; 9 out of 13 mismatches (including two deletions) were located at the 5' end, clustered in the first 146 nucleotides (designated a and a' in Fig. 2), whereas 4 out of 13 are located at the 3' end, clustered in the last 42 nucleotides (designated b and b' in Fig. 2). PCR and sequencing analysis showed that the Ω cat(pC194) element excised from Tn5253 produced a circular intermediate and a deletion in Tn5253 (Fig. 2). The recombination event leading to the excision of Ω cat(pC194) from Tn5253 occurred within the 1,169-bp segment, leaving in Tn5253 a copy of the 1,169-bp segment flanked by imperfect DRs a and b', whereas in the circular intermediate it was flanked by imperfect DRs a' and b. The element contained 11 ORFs (*orf58* to *orf68*), 5 out of 11 belonging to plasmid pC194, while 4 out of 11, *orf65* to *orf68*, showed a high level of homology (>90%) to region *spr0948* to *spr0952* of R6 *S. pneumoniae* (Table 1). When investigating Ω cat(pC194) mobility, we found that under our experimental conditions, the element could not be transferred by conjugation ($<4.1 \times 10^7$ transconjugants per donor), but it was capable of intracellular mobility. In fact, we found and sequenced a copy of Ω cat(pC194) integrated into the chromosomal *spr0948*-to-*spr0952* region (GenBank accession no. GU808561). DNA sequence analysis of three different transconjugants containing Tn5253 indicated that ectopic integration of Ω cat(pC194) occurred by insertion duplication at different positions between *spr0948* and *spr0952* (data not shown), as expected for homologous recombination. This ectopic integration generated two long DRs that flank the element and are homologous to the long DRs that also flank Ω cat(pC194) when it is inserted into Tn5253. It is interesting that the long DRs at both sites each contain a complete copy of the genes coding for a toxin-antitoxin system (18), the presence of which may contribute to the stable maintenance of these genetic elements in the bacterial host.

In conclusion, sequencing of the whole element allowed us to clearly define the composite nature of Tn5253, which was shown to contain two independent genetic elements, (i) fully functional conjugative transposon Tn5251 (11) and (ii) the newly recognized *cat*-containing element Ω cat(pC194). It is noteworthy that a large fraction (52%) of the Tn5253 ORFs code for putative proteins of unknown function, confirming that previously undescribed genes are preferentially carried by mobile genetic elements and reinforcing the notion that investigation of the mobilome is essential for understanding bacterial genomes (19).

ACKNOWLEDGMENTS

This work is dedicated to the memory of Walter Guild, who, by working on Tn5253, pioneered the field of research on integrative and conjugative elements. We also acknowledge the decisive contribution to the understanding of Tn5253 of the people from Walter Guild's lab: Nadja Shoemaker, Mike Smith, Scott Priebe, Judith Hageman, and Moses Vijayakumar.

The research leading to these results has received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement 241446 (project ANTIRESEDEV).

REFERENCES

- Croucher NJ, Harris SR, Fraser C, Quail MA, Burton J, van der Linden M, MacGee L, von Gottberg A, Song JH, Ko KS, Pichon B, Baker S, Parry CM, Lamberts LM, Shahinas D, Pillai DR, Mitchell TJ, Dougan G, Tomasz A, Klugman KP, Parker J, Hanage WP, Bentley SD. 2011. Rapid pneumococcal evolution in response to clinical interventions. *Science* 331:430–434. <http://dx.doi.org/10.1126/science.1198545>.
- Mullany P, Roberts AP, Wang A. 2002. Mechanism of integration and excision in conjugative transposons. *Cell. Mol. Life Sci.* 59:2017–2022. <http://dx.doi.org/10.1007/s000180200001>.
- Shoemaker NB, Smith MD, Guild WR. 1979. Organization and transfer of heterologous chloramphenicol and tetracycline resistance genes in pneumococcus. *J. Bacteriol.* 139:432–441.
- Vijayakumar MN, Priebe SD, Pozzi G, Hageman JM, Guild WR. 1986. Cloning and physical characterization of chromosomal conjugative elements in streptococci. *J. Bacteriol.* 166:972–977.
- Shoemaker NB, Smith MD, Guild WR. 1980. DNase-resistant transfer of chromosomal *cat* and *tet* insertions by filter mating in pneumococcus. *Plasmid* 3:80–87. [http://dx.doi.org/10.1016/S0147-619X\(80\)90036-0](http://dx.doi.org/10.1016/S0147-619X(80)90036-0).
- Pozzi G, Musmanno RA, Renzoni EA, Oggioni MR, Cusi MG. 1988. Host-vector system for integration of recombinant DNA into chromosomes of transformable and nontransformable streptococci. *J. Bacteriol.* 170:1969–1972.
- Vijayakumar MN, Ayalew S. 1993. Nucleotide sequence analysis of the termini and chromosomal locus involved in site-specific integration of the streptococcal conjugative transposon Tn5252. *J. Bacteriol.* 175:2713–2719.
- Provedi R, Manganelli R, Pozzi G. 1996. Characterization of conjugative transposon Tn5251 of *Streptococcus pneumoniae*. *FEMS Microbiol. Lett.* 135:231–236. <http://dx.doi.org/10.1111/j.1574-6968.1996.tb07994.x>.
- Ayoubi P, Kiliç AO, Vijayakumar MN. 1991. Tn5253, the pneumococcal Ω (*cat tet*) BM6001 element, is a composite structure of two conjugative transposons, Tn5251 and Tn5252. *J. Bacteriol.* 173:1617–1622.
- Rice LB. 1998. Tn916 family of conjugative transposons and determination of antimicrobial resistance determinants. *Antimicrob. Agents Chemother.* 42:1871–1877.
- Santoro F, Oggioni MR, Pozzi G, Iannelli F. 2010. Nucleotide sequence and functional analysis of the *tet*(M)-carrying conjugative transposon Tn5251 of *Streptococcus pneumoniae*. *FEMS Microbiol. Lett.* 308:150–158. <http://dx.doi.org/10.1111/j.1574-6968.2010.02002.x>.
- Widdowson CA, Adrian PV, Klugman KP. 2000. Acquisition of chloramphenicol resistance by the linearization and integration of the entire staphylococcal plasmid pC194 into the chromosome of *Streptococcus pneumoniae*. *Antimicrob. Agents Chemother.* 44:393–395. <http://dx.doi.org/10.1128/AAC.44.2.393-395.2000>.
- Smith MD, Hazum S, Guild WR. 1981. Homology among *tet* determinants in conjugative elements of streptococci. *J. Bacteriol.* 148:232–240.
- Kiliç AO, Vijayakumar MN, Al-khaldi SF. 1994. Identification and nucleotide sequence analysis of a transfer-related region in the streptococcal conjugative transposon Tn5252. *J. Bacteriol.* 176:5145–5150.
- Pozzi G, Guild WR. 1988. Two genes for chloramphenicol resistance common to staphylococci and streptococci. *Eur. J. Epidemiol.* 4:20–24. <http://dx.doi.org/10.1007/BF00152687>.
- Burrus V, Waldor MK. 2004. Shaping bacterial genomes with integrative and conjugative elements. *Res. Microbiol.* 155:376–386. <http://dx.doi.org/10.1016/j.resmic.2004.01.012>.
- Iannelli F, Pozzi G. 2007. Protocol for conjugal transfer of genetic elements in *Streptococcus pneumoniae*, p 525–529. In Hakenbeck R, Chhatwal GS (ed), *Molecular biology of streptococci*. Horizon Bioscience, Norfolk, United Kingdom.
- Khoo SK, Loll B, Chan WT, Shoeman RL, Ngo L, Yeo CC, Meinhart A. 2007. Molecular and structural characterization of the PezAT chromosomal toxin-antitoxin system of the human pathogen *Streptococcus pneumoniae*. *J. Biol. Chem.* 282:19606–19618. <http://dx.doi.org/10.1074/jbc.M701703200>.
- Frost LS, Leplae R, Summers AO, Toussaint A. 2005. Mobile genetic

- elements: the agents of open source evolution. *Nat. Rev. Microbiol.* 3:722–732. <http://dx.doi.org/10.1038/nrmicro1235>.
20. Díaz-López T, Lages-Gonzalo M, Serrano-López A, Alfonso C, Rivas G, Diaz-Orejas R, Giraldo R. 2003. Structural changes in RepA, a plasmid replication initiator, upon binding to origin DNA. *J. Biol. Chem.* 278: 18606–18616. <http://dx.doi.org/10.1074/jbc.M212024200>.
 21. Sampath J, Vijayakumar MN. 1998. Identification of a DNA cytosine methyltransferase gene in conjugative transposon Tn5252. *Plasmid* 39: 63–76.
 22. Garvey P, Fitzgerald GF, Hill C. 1995. Cloning and DNA sequence analysis of two abortive infection phage resistance determinants from the lactococcal plasmid pNP40. *Appl. Environ. Microbiol.* 61:4321–4328.
 23. Schröder G, Krause S, Traxler B, Yeo HJ, Lurz R, Waksman G, Lanka E. 2002. TraG-like proteins of DNA transfer systems and of the *Helicobacter pylori* type IV secretion system: inner membrane gate for exported substrates? *J. Bacteriol.* 184:2767–2789. <http://dx.doi.org/10.1128/JB.184.10.2767-2779.2002>.
 24. Lee HS, Kim MS, Cho HS, Kim JI, Kim TJ, Choi JH, Park C, Lee HS, Oh BH, Park KH. 2002. Cyclomaltodextrinase, neopullulanase, and maltogenic amylase are nearly indistinguishable from each other. *J. Biol. Chem.* 277:21891–21897. <http://dx.doi.org/10.1074/jbc.M201623200>.
 25. Munoz-Najar U, Vijayakumar MN. 1999. An operon that confers UV resistance by evoking the SOS mutagenic response in streptococcal conjugative transposon Tn5252. *J. Bacteriol.* 181:2782–2788.
 26. Zhang S, Meyer R. 1997. The relaxosome protein MobC promotes conjugative plasmid mobilization by extending DNA strand separation to the nick site at the origin of transfer. *Mol. Microbiol.* 25:509–516. <http://dx.doi.org/10.1046/j.1365-2958.1997.4861849.x>.
 27. Srinivas P, Kiliç AO, Vijayakumar MN. 1997. Site-specific nicking *in vitro* at *oriT* by the DNA relaxase of Tn5252. *Plasmid* 37:42–50. <http://dx.doi.org/10.1006/plas.1996.1268>.
 28. Brennan RG, Matthews BW. 1989. The helix-turn-helix DNA binding motif. *J. Biol. Chem.* 264:1903–1906.
 29. Rauch PJG, De Vos WM. 1994. Identification and characterization of genes involved in excision of the *Lactococcus lactis* conjugative transposon Tn5276. *J. Bacteriol.* 176:2165–2171.
 30. Gindreau E, Lopez R, Garcia P. 2000. MM1, a temperate bacteriophage of the type 23F Spanish/USA multiresistant epidemic clone of *Streptococcus pneumoniae*: structural analysis of the site-specific integration system. *J. Virol.* 74:7803–7813. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC112310/pdf/jv007803.pdf>.

TABLE S1 Oligonucleotide primers^a

Name	Sequence (5' to 3')	GenBank ID: nucleotides	Amplicon length (bp)
Tn1	TGA CCG ATG AGC TTG TTT TG	AE007317: 1,035,500-1,035,519	7,683
IF330	ATT TGG GAT TTA GCC TCA TCA A	AF295925: 9,098-9,119	
IF329	CCT TAA TCG TCT CAG AAT ATC A	L29323: 129-108	13,493
IF334	TGA ATA AAG AGA AGC AAC AGG A	X90940: 116-137	
IF487	TTC GCT GAA GAC CTT TAT TCG	X90940: 378-358	5,518
IF393	TTC TGC CGA AAT TGT AAT CA	X90939: 2,541-2,560	
IF394	GCT ATA GTA TAA GCC ATA CT	X90939: 3,621-3,602	13,743
IF488	TCC TCC TGA TTC CAG TGT CA	X90941: 52-71	
IF333	GAC TTT ATC CTC GCC AGC AA	X90941: 558-539	16,755
IF353	CTC TCC GTC GCT ATT GTA AC	V01277: 1,487-1,468	
IF184	AAC TTT AAT AAA ATT GAT TTA GA	V01277: 1,263-1,285	6,576
IF367	GAA TAG TTC CGA TTG AGT GAA A	L29324: 7,511-7,532	
IF368	AAC AGA ATG CGA AGG TTG TAG A	L29324: 7,841-7,820	8,664
Tn2	TTA ATC GTT GCC ATC GAG GT	AE007317:1,037,162-1,037,143	
IF460	TGT AGA AAT CAA AGT CCT AGC	AE007317: 931746-931766	12,314
IF461	ATC ATA AGA AAC AGG TGT ATG C	AE007317: 936264-936243	

^aPrimers are listed in pairs, and the length of amplicons produced is reported