

Pleural plaques: markers of asbestos exposure or independent risk factor for pleural mesothelioma? a case report

This is the peer reviewed version of the following article:

Original:

Sartorelli, P., D'Hauw, G., Sisinni, A.g., Romeo, R., Paolucci, V., Spina, D. (2018). Pleural plaques: markers of asbestos exposure or independent risk factor for pleural mesothelioma? a case report. OCCUPATIONAL AND ENVIRONMENTAL MEDICINE, 75(Suppl. 2), A456-A457 [10.1136/oemed-2018-ICOHabstracts.1298].

Availability:

This version is available <http://hdl.handle.net/11365/1051481> since 2018-05-17T11:24:18Z

Published:

DOI:10.1136/oemed-2018-ICOHabstracts.1298

Terms of use:

Open Access

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.

(Article begins on next page)

PLEURAL PLAQUES: MARKERS OF ASBESTOS EXPOSURE OR INDEPENDENT RISK FACTOR FOR PLEURAL MESOTHELIOMA? A CASE REPORT

Sartorelli P, d'Hauw G, Sisinni AG, Romeo R, Paolucci V, Spina D¹

Department of Medical Biotechnology, Unit of Occupational Medicine AOU Senese, University of Siena, Siena, Italy

¹ Department of Medical Biotechnology, Unit of Anatomic Pathology AOU Senese, University of Siena, Siena, Italy

Abstract

Pleural plaques (PPs) represent the most common non malignant asbestos-related disease. Currently PPs are widely referred as a markers of asbestos exposure and not as an independent risk factor for the development of asbestos-related neoplastic diseases (1). However, the association between PPs and pleural mesothelioma (PM) remains controversial, as some studies on cohorts of asbestos exposed workers have shown correlation between PPs and PM. Despite this, scientific literature is poor. Research has been focused on the relationship between PPs and lung cancer, while there is a lack of studies on the possible relationship between PPs and the development of PM.

Case presentation

A clinical case of a patient with a history of occupational asbestos exposure affected by asbestosis and PPs is presented. During the follow up managed by the Occupational Medicine of Siena a PM was diagnosed at a PP.

Conclusions

The widespread belief that there is no relationship between PPs and PM is not supported by a vast literature as it happens for the absence of association between PPs and lung cancer. In the patient described it is not possible to determine whether the localization of the PM at a PP is incidental or not and whether the PM can be attributed only to intense exposure or even to the presence of PPs. Similar cases may not be uncommon even if not reported in the literature.

References

1. Maxim LD, Niebo R, Utell MJ (2015) Are pleural plaques an appropriate endpoint for risk analyses? *Inhalation Toxicology*; 27: 321-334
2. Pairon JC, Laurent F, Rinaldo M, Clin B, Andujar P et al. (2013) Pleural Plaques and the Risk of Pleural Mesothelioma. *JNCI*; 105: 293-301; doi: 10.1093/jnci/djs513