



“How to detect the impact of marine debris on Mediterranean biodiversity? The three fold monitoring approach”

Maria Cristina Fossi, Department of Physical, Earth and Environmental Sciences, University of Siena, Siena, Italy

Matteo Baini, Department of Physical, Earth and Environmental Sciences, University of Siena, Siena, Italy

Cristina Panti, Department of Physical, Earth and Environmental Sciences, University of Siena, Siena, Italy

ABSTRACT

The Mediterranean Sea has been described as one of the most affected areas by marine debris, including microplastics, in the world. Recent studies in the different regions of the basin suggested that some areas are affected by important concentration of microplastics and plastic additives, representing a potential risk for endangered species (baleen whales, sea turtles, filter feeders sharks) and for the all Mediterranean biodiversity. To cover the current knowledge gaps on this issue a harmonised methodological approach for the assessment of the marine debris impact on Mediterranean biodiversity is needed. The quantification of marine debris/microplastics in the marine environment can depend on several environmental factors and change according to multiple oceanographic features, and therefore, cannot reflect the potential impact on organisms and ecosystems. The information obtained by biondicator species could better integrate the spatial and temporal presence of marine litter/microplastics in the marine environment. In addition, the use of biondicators can allow to measure not only the occurrence of marine litter in the species and its environment but also the threat posed to organisms by the evaluation of contaminants accumulation and any related biological effect. To assess the harm by marine debris ingestion a threefold approach, simultaneously measuring the presence and effects (accumulation of plastic associated contaminants and biomarker responses), can provide the harm and the sub-lethal effects to organisms related marine litter ingestion. The gaps pointed out by this research and the bioindicators species selected could represent a step forward for the risk assessment and the implementation of future mitigation measure for the Mediterranean area, habitat and species affected by marine litter ingestion.