

Decreased neutralization of the Eta SARS-CoV-2 variant by sera of previously infected and uninfected vaccinated individuals

This is the peer reviewed version of the following article: Original: Boccuto, A., Dragoni, F., Bergna, A., Ventura, C.D., Giammarino, F., Saladini, F., et al. (2021). Decreased neutralization of the Eta SARS-COV-2 variant by sera of previously infected and uninfected vaccinated individuals. JOURNAL OF INFECTION [10.1016/j.jinf.2021.08.005]. Availability: This version is availablehttp://hdl.handle.net/11365/1152828 since 2021-09-08T06:27:10Z Published: DOI:10.1016/j.jinf.2021.08.005 Terms of use: Open Access The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. Works made available under a Creative Commons license can be used according to the terms and conditions of said license.

For all terms of use and more information see the publisher's website.

(Article begins on next page)



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

JID: YJINF

ARTICLE IN PRESS

Journal of Infection xxx (xxxx) xxx



Contents lists available at ScienceDirect

Journal of Infection



journal homepage: www.elsevier.com/locate/jinf

Letter to the Editor

Decreased neutralization of the Eta SARS-CoV-2 variant by sera of previously infected and uninfected vaccinated individuals

Dear Editor,

In this Journal, the emergence of novel variants with potential to escape vaccine-induced immunity has received commentary [1].

The emergence of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) variants of concern (VOC) and variants of interest (VOI) are challenging the management of the evolving pandemic across countries. The VOI labelled as Eta (WHO Classification) [1], combines relevant spike mutations detected in several VOC, such as the same 3 deletions of the Alpha lineage (69del, 70del, 144del), the E484K mutation found in the Gamma and Beta lineages as well as in some Alpha isolates and the ubiquitous D614G. In addition, three mutations (A67V, Q677H and F888L) are unique to Eta variant and it is currently unknown whether they favor escape from natural or vaccine induced immunity to the wild type lineage (B.1), as shown for other variants [2]. To test this hypothesis, we measured the serum neutralizing antibody (NtAb) response to Eta variant, as well as to other viral variants, in a cohort of health care workers (HCWs) including both previously infected (n = 15) and uninfected individuals (n = 15) vaccinated with two doses of the BNT162b2 COVID-19 mRNA vaccine. The study was approved by the Ethics Committee of the University of Milan (protocol n. 23/21) and conducted in compliance with Good Clinical Practice guidelines and the Declaration of Helsinki. The previously infected group was tested at baseline (T0_{inf}) and 17\pm6 days after receiving the second vaccine dose (T2_{inf}); the uninfected HCWs were tested 18 \pm 4 days after the second dose vaccination (T2_{uninf}). The infected group had median age [IQR] of 38 (31-52) years, included 8 females and was infected during the first wave of the pandemic. The uninfected group had a median age of 38 (29-59) years with 11 females.

NtAb titers were determined by a microneutralization live virus assay performed in VERO E6 cells using the quantification of cell viability as readout system, as previously described [3]. NtAb titers were expressed as median (IQR) and were defined as the reciprocal value of the sample dilution that showed a 50% protection of virusinduced cytopathic effect (ID₅₀). Sera with ID₅₀ titres \geq 10 were defined as SARS-CoV-2 neutralizing, while sera with ID₅₀ <10 were defined as negative and scored as 5 for statistical analysis. Fifteen, 14 and 11 individuals at $T0_{inf}$, $T2_{inf}$ and $T2_{uninf}$, respectively, had also a quantitative anti-spike protein Ab determination, performed by the SARS-CoV-2 IgG II Quant assay (Abbott). The viral isolates used in the microneutralization live virus assay were sequenced by NGS and the full-lenght SARS-CoV-2 genome was submitted to GI-SAID (http://gisaid.org/) to assign the right variant (Accession numbers: EPI_ISL_2,472,896, EPI_ISL_1,085,167, EPI_ISL_2,472,918 and EPI_ISL_2,472,916 for the wild type, Alpha, Gamma and Eta vari-



Fig. 1. Neutralizing antibody (NtAb) titres to four SARS-CoV-2 variants (wild type, Gamma, Alpha and Eta) in 15 previously infected subjects at baseline (TO_{inf}) and after two doses of vaccine ($T2_{uninf}$) and in 15 uninfected subjects after two doses of vaccine ($T2_{uninf}$). Asterisks indicate significance levels: ***, p < 0.001. Median (IQR) titres of neutralizing antibody are reported below.

ants, respectively). Statistical analyses were performed using IBM SPSS Statistics, version 20. The non-parametric Friedman test and Wilcoxon Signed Rank Sum test was used to analyze changes in paired data. The non-parametric Mann-Whitney test was used to compare unpaired data. Spearman analysis was used to measure the correlation between NtAb titres against the different variants.

In previously infected HCWs, NtAb titres to all viral variants significantly increased after vaccination (mean T2_{inf}/T0_{inf} ratio 119 \pm 66; *p*<0.001). Notably, 2 to 12 subjects, depending on the reference virus, were negative at TO_{inf} but all of them seroconverted following vaccination. As expected, the NtAb titer after vaccination was higher in the previously infected compared with the uninfected group (mean T2_{inf}/T2_{uninf} ratio 6 ± 2 ; p < 0.001 (Fig. 1). Overall, median NtAb titres to the Eta variant (63 [7-323] ID₅₀) correlated well with those to the wild type (133 [9-456]), Gamma (148 [46-988]) and Alpha (87 [5-681]) (p<0.001 for all comparisons) and high correlation was indeed observed between NtAb titres to any pair of virus variants (Fig. 2). Of note, NtAb titres to Eta variant were significantly lower with respect to those obtained for each variant (p < 0.001). Anti-spike protein antibodies, as measured by enzyme immunoassay, were highly correlated with NtAb titres to B.1 (rho = 0.934), P.1 (rho = 0.914), B.1.1.7 (rho = 0.913) and A. Boccuto, F. Dragoni, A. Bergna et al.

ARTICLE IN PRESS



Fig. 2. Spearman correlation between NtAb titres to each pair of the SARS-CoV-2 variants used in the study. Data were cumulated for all sera tested at T0_{inf}, T2_{inf} and T2_{uninf}.

B.1.525 (rho = 0.918) viruses (p<0.001 for all comparisons). Also, a significant increase was observed when comparing the anti-spike Ab median titres at T2_{inf} and at T0_{inf} (27,763 [18,282–46,108] vs. 1.7 [0.5–4.4]; p = 0.001).

Overall, in our small cohort of previously infected or uninfected vaccinated-HCWs it appears that cross-neutralization among different viral variants remains substantial, following natural or artificial immunization with the wild type lineage. However, neutralization of Eta variant is significantly reduced with respect to other variants. Indeed, NtAb titres could be ranked with the definite order Gamma>wild type=Alpha>Eta. In vitro correlates of protection against the Eta variant has been investigated in uninfected vaccinated individuals only in two different works delivering inconsistent results. Indeed, Liu et al. [4] observed a modest reduction, while Zani et al. [5] reported an increase in Eta variant NtAb titres with respect to the wild type variant. Of note, NtAb studies published so far have used different combination of strategies (e.g., live virus vs. pseudoparticles), viral variants, cell lines and readouts, in the absence of standardized methods and reference viral strains and neutralizing sera [6-9]. For example, the full-length sequencing of the isolates used in the assay should be always reported and submitted to public repositories. Most importantly, while NtAb studies certainly provide a solid basis to infer crossprotection among vaccines and virus variants, the in vivo correlates of in vitro data remain to be established and must be defined through accurate and continuous monitoring of vaccine induced reduction of morbidity and mortality in the context of molecular surveillance of SARS-CoV-2 lineages.

References

- **1.** Pereira F, Tosta S, Lima MM, Reboredo de Oliveira da Silva L, Nardy VB, Gómez MKA, et al. Genomic surveillance activities unveil the introduction of the SARS-CoV-2 B1.525 variant of interest in Brazil: case report. *J Med Virol* 2021 Epub ahead of print.
- Janik E, Niemcewicz M, Podogrocki M, Majsterek I, Bijak M. The Emerging Concern and Interest SARS CoV-2 Variants. *Pathogens* 2021;10(6):633.
- Vicenti I, Gatti F, Scaggiante R, Boccuto A, Zago D, Basso M, et al. Single-dose BNT162b2 mRNA COVID-19 vaccine significantly boosts neutralizing antibody response in health care workers recovering from asymptomatic or mild natural SARS-CoV-2 infection. Int J Infect Dis 2021;108:176–8.
- Liu J, Liu Y, Xia H, Zou J, Weaver SC, Swanson KA, et al. BNT162b2-elicited neutralization of B.1.617 and other SARS-CoV-2 variants. *Nature* 2021(June).
- Zani A, Caccuri F, Messali S, Bonfanti C, Caruso A. Serosurvey in BNT162b2 vaccine-elicited neutralizing antibodies against authentic B.1, B.1.1.7, B.1.351, B.1.525 and P.1 SARS-CoV-2 variants. *Emerg Microbes Infect* 2021;7:1–6.
- Betton M, Livrozet M, Planas D, Fayol A, Monel B, Védie B, et al. Sera neutralizing activities against SARS-CoV-2 and multiple variants six month after hospitalization for COVID-19. *Clin Infect Dis* 2021 ciab308Epub ahead of printPMID: 33851216PMCID: PMC8083257. doi:10.1093/cid/ciab308.
- Planas D, Bruel T, Grzelak L, Guivel-Benhassine F, Staropoli I, Porrot F, Planchais C, et al. Sensitivity of infectious SARS-CoV-2 B1.1.7 and B.1.351 variants to neutralizing antibodies. *Nat Med* 2021;27(5):917–24.
- Wang P, Casner RG, Nair MS, Wang M, Yu J, Cerutti G, Liu L, et al. Increased resistance of SARS-CoV-2 variant P.1 to antibody neutralization. *Cell Host Microbe* 2021;29(5):747–51 e4.
- 9. Wang P, Nair MS, Liu L, Iketani S, Luo Y, Guo Y, et al. Antibody resistance of SARS-CoV-2 variants B.1.351 and B.1.1.7. *Nature* 2021;593(7857):130–5.

Adele Boccuto[#], Filippo Dragoni[#] Department of Medical Biotechnologies, University of Siena, Siena, Italy

ARTICLE IN PRESS

[m5G;August 21, 2021;12:25]

Journal of Infection xxx (xxxx) xxx

A. Boccuto, F. Dragoni, A. Bergna et al.

JID: YJINF

Maurizio Zazzi, Ilaria Vicenti Department of Medical Biotechnologies, University of Siena, Siena, Italy

Alessia Lai* Department of Biomedical and Clinical Sciences L. Sacco, University of Milan, Milan, Italy

*Corresponding author.

E-mail address: alessia.lai@unimi.it (A. Lai) # Adele Boccuto and Filippo Dragoni contributed equally to the manuscript

Annalisa Bergna, Carla Della Ventura Department of Biomedical and Clinical Sciences L. Sacco, University of Milan, Milan, Italy

Federica Giammarino, Francesco Saladini Department of Medical Biotechnologies, University of Siena, Siena, Italy

Laura Pezzati Department of Infectious Diseases, ASST Fatebenefratelli Sacco, Milan, Italy

Gianguglielmo Zehender Department of Biomedical and Clinical Sciences L. Sacco, University of Milan, Milan, Italy