

*This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.*



**ISSN: 2155-9570**

## **Journal of Clinical & Experimental Ophthalmology**

The International Open Access  
Journal of Clinical & Experimental Ophthalmology

### **Editor-in-Chief**

**Richard B. Rosen**

Professor of Ophthalmology, New York Medical College, USA

### **Executive Editors**

**Sayon Roy**

Boston University School of Medicine, USA

**Yoko Miura**

University of Luebeck, Germany

**Available online at:** OMICS Publishing Group ([www.omicsonline.org](http://www.omicsonline.org))

This article was originally published in a journal by OMICS Publishing Group, and the attached copy is provided by OMICS Publishing Group for the author's benefit and for the benefit of the author's institution, for commercial/research/educational use including without limitation use in instruction at your institution, sending it to specific colleagues that you know, and providing a copy to your institution's administrator.

All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are requested to cite properly.

Digital Object Identifier: <http://dx.doi.org/10.4172/2155-9570.1000313>

# Qualitative Investigation of Corneal Changes after Accelerated Corneal Collagen Cross-linking (A-CXL) by *In vivo* Confocal Microscopy and Corneal OCT

Cosimo Mazzotta<sup>1</sup>, Anna Lucia Paradiso<sup>1</sup>, Stefano Baiocchi<sup>1</sup>, Stefano Caragiuli<sup>1</sup> and Aldo Caporossi<sup>2</sup><sup>1</sup>Department of Medical, Surgical and Neurosciences, Ophthalmology Section, Siena University, Italy<sup>2</sup>Ophthalmology Institute, Rome Catholic University, Italy**Abstract**

**Purpose:** To assess qualitative micro-morphological corneal changes by confocal microscopy and corneal OCT after accelerated corneal crosslinking (A-CXL) in keratoconic patients.

**Study design:** Prospective non-randomized pilot study.

**Methods:** 20 eyes of 20 patients, aged between 13 and 26 years (mean 22.6 years) underwent A-CXL by the KXL UV-A source (Avedro Inc. Waltham MA, USA). Patients were divided into 4 groups according to different riboflavin solutions and UV A powers. 15 patients underwent epithelium-off A-CXL: 5 (Group 1) by riboflavin 0.1% plus dextran 20% at 12 mW/cm<sup>2</sup> for 10 min; 5 (Group 2) at 30 mW/cm<sup>2</sup> for 4 min; 5 (Group 3) by dextran-free riboflavin 0.1% plus HPMC at 30 mW/cm<sup>2</sup> for 4 min and 5 (Group 4) by riboflavin 0.25% plus EDTA, BAK, TRIS epithelium-on A-CXL for 2 min and 40 sec. Micro-morphological analysis was assessed by *in vivo* HRT II confocal microscopy and corneal OCT.

**Results:** Epithelium regenerated into 3 days. Sub-epithelial nerves disappeared after treatment regenerating into 6 months. Epithelium off A-CXL penetration, measured evaluating keratocytes loss at confocal microscopy and demarcation lines at corneal OCT, resulted at 180 μm on average in the Group 1, 160 μm in the Group 2, 150 μm in the Group 3. Epithelium-on A-CXL (Group 4) revealed a penetration at 80 μm on average. No endothelial damage was recorded in all groups.

**Conclusion:** A-CXL shortened conventional CXL procedure under 20 minutes, being well tolerated. Its clinical efficacy needs to be determined in the mid-long term follow-up and in a large cohort of patients.

**Keywords:** Accelerated cross-linking; A-CXL; Keratoconus; Confocal microscopy; Demarcation line

**Introduction**

Riboflavin UV-A induced corneal collagen crosslinking (CXL) represents a relatively new procedure available for the conservative treatment of progressive keratoconus [1,2] and secondary corneal ectasia [3] due to its capacity in increasing biomechanical corneal resistance [4,5] and intrinsic anti-collagenase activity [6].

The physicochemical basis of crosslinking lies in the photo-dynamic type I-II reactions [7] induced by the interaction between 0.1% riboflavin molecules absorbed in corneal tissue and UV-A rays delivered at 3 mW/cm<sup>2</sup> for 30 minutes (5.4 J/cm<sup>2</sup> energy dose) releasing reactive oxygen species (ROS) that mediated cross-links formation between and within collagen fibers [8,9].

The conventional epithelium-off cross-linking procedure (CXL) demonstrated its safety and long-term efficacy stabilizing progressive keratoconus and secondary ectasia in different clinical trials [10-14]. On the other hand the procedure is time consuming lasting from 40 minutes to 1 hour [15] with patient's discomfort.

The physical concept of photochemical reactions stated in the Bunsen-Roscoe's law of reciprocity [16-18] theoretically demonstrated that the photochemical process behind cross-linking depends on the absorbed UV-A energy and its biological effect is proportional to the total energy dose delivered in the tissue [16-18].

According to this physical theory it is theoretically possible to deliver the same energy dose ensuring a proportional biological effect by setting different UV-A powers and exposure times in order

to accelerate and shorten the crosslinking procedure in the so called Accelerated Cross-Linking (A-CXL) [17-19].

According to "equal-dose" principle 10 mW/cm<sup>2</sup> for 9 min, 30 mW/cm<sup>2</sup> for 3 min, 18 mW/cm<sup>2</sup> for 5 min, 45 mW/cm<sup>2</sup> for 2 min at constant energy dose of 5.4 J/cm<sup>2</sup> are the same as the standard 3 mW/cm<sup>2</sup> for 30 min, a basic concept leading to A-CXL [18,19].

An energy dose of 7.2 J/cm<sup>2</sup> was demonstrated to be effective both in terms of corneal strengthening and anti-enzyme activity compared with the standard dose of 5.4 J/cm<sup>2</sup>, respectively tested by biaxial corneal extensimetry and papain digestion (Avedro's laboratory unpublished data, presented by M. D. Friedman, Ph.D at 8<sup>th</sup> International CXL Congress, Geneva 8 December 2012).

In this pilot study we report a qualitative analysis of the cornea after A-CXL assessed by means of *in vivo* HRT II scanning laser confocal microscopy (Heidelberg, Germany) and by Visante time domain

**\*Corresponding author:** Cosimo Mazzotta MD, PhD, Dipartimento di Scienze Mediche, Chirurgiche e Neuroscienze, U.O.C Oculistica, Policlinico Santa Maria delle Scotte, Viale Bracci 8, 53100 Siena, Italy, Tel: +39 0577 356618; Fax: +39 0577 356618; E-mail: [cgmazzotta@libero.it](mailto:cgmazzotta@libero.it)

**Received** November 04, 2013; **Accepted** December 06, 2013; **Published** December 13, 2013

**Citation:** Mazzotta C, Paradiso AL, Baiocchi S, Caragiuli S, Caporossi A (2013) Qualitative Investigation of Corneal Changes after Accelerated Corneal Collagen Cross-linking (A-CXL) by *In vivo* Confocal Microscopy and Corneal OCT. J Clin Exp Ophthalmol 4: 313. doi: [10.4172/2155-9570.1000313](http://dx.doi.org/10.4172/2155-9570.1000313)

**Copyright:** © 2013 Mazzotta C, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

corneal OCT (Zeiss Meditec, Jena, Germany), in a series of 20 patients with progressive keratoconus investigating the induced corneal changes and the penetration of A-CXL.

## Methods

After unanimous approval of the local ethics committee under the principles of the Helsinki declaration and signing of specific informed consent 20 eyes of 20 patients affected from keratoconus, age between 13 and 26 years (mean 22.6 years), underwent Accelerated CXL by the KXL UV-A source (Avedro Inc. Waltham MS, USA) at the Ophthalmology Unit of Siena University Hospital. All patients included in the treatment protocol were affected by progressive keratoconus with a documented clinical and instrumental worsening at least in the last three months of observation.

## Inclusion criteria

The parameters considered to establish keratoconus progression and inclusion criteria for each group were: worsening of UCVA/BSCVA > 0.50 Snellen lines, increase of SPH/CYL > 0.50 D, increase of topographic symmetry index SAI/SI > 0.50 D, increase of maximum K reading > 1 D, reduction of the thinnest point at AC OCT optical pachymetry  $\geq 10 \mu\text{m}$ , clear cornea at bio-microscopic examination, absence of reticular dark striations at confocal laser microscopy *in vivo*. We considered "significant" for the inclusion in the study the variation of at least 3 of the parameters listed above (one clinical plus two instrumental). The following examinations were performed before and after the operation: *in vivo* scanning laser confocal microscopy (HRT II, Rostock Cornea Module, Heidelberg, Germany) and anterior segment OCT analysis (Visante OCT, Zeiss Meditec, Jena, Germany) to assess qualitative A-CXL induced corneal changes and treatment penetration.

Patients were divided into 4 groups matched according to age and keratoconus stage as different A-CXL protocols showed in Table 1.

**Group 1 epithelium-off A-CXL:** 5 eyes, age 16-23 y (mean age 19 years) Riboflavin 0.1% plus Dextran 20% (VibeX), 15 minutes of corneal soaking after mechanical epithelial debridement (blunt metal spatula), UV-A power at  $12 \text{ mW/cm}^2$  (Energy dose:  $7.2 \text{ J/cm}^2$ ), 10 min exposure time.

**Group 2 epithelium-off A-CXL:** 5 eyes, age 13-26 y (mean age 19.5 years) Riboflavin 0.1% plus Dextran 20% (VibeX), 20 minutes of corneal soaking after mechanical epithelial debridement (blunt metal spatula), UV-A power at  $30 \text{ mW/cm}^2$  (Energy dose:  $7.2 \text{ J/cm}^2$ ), 4 min exposure time.

**Group 3 epithelium-off A-CXL:** 5 eyes, age 14-24 y (mean age 20.5 years) Riboflavin 0.1% (dextran free) plus HPMC (VibeX Rapid), 10 minutes of corneal soaking after mechanical epithelial debridement

(blunt metal spatula), UV-A power at  $30 \text{ mW/cm}^2$  (Energy dose:  $7.2 \text{ J/cm}^2$ ), 4 min exposure time.

**Group 4 epithelium-on A-CXL:** 5 eyes, age 21-26 y (mean age 23.5 years) Riboflavin 0.25% plus EDTA, BAK, TRIS (Paracel) tapered every 90 seconds for 4 minutes of soaking, followed by corneal rinsing with Riboflavin 0.25% saline solution (Vibex X-tra) administered every 90 seconds for 6 min (total epithelium-on soaking time 10 minutes), UV-A power at  $45 \text{ mW/cm}^2$  (Energy dose:  $7.2 \text{ J/cm}^2$ ), 2 min. and 40 sec exposure time.

## Postoperative protocol

All patients underwent a postoperative soft contact lens bandage for 3 days, cyclopentolate eye drops twice for 3 days, ciprofloxacin eye drops four times/day for 3 days, diclofenac eye drops four times/day for 3 days and eye lubricants four times/day and on demand. After therapeutic contact lens removal all patients were medicated by dexamethasone eye drops and sodium hyaluronate 0.2% eye lubricants 4 times/day for 15 days and 2 times/day for 15 days.

Treatment penetration (keratocytes loss, cornea edema) was compared by a descriptive point of view with literature data, coming out from our research team on conventional [20-22] and trans-epithelial CXL [23].

## Results

A comprehensive review of treatment groups and results are summarized in Table 1.

### Group 1 epithelium-off A-CXL

Riboflavin 0.1% plus Dextran 20% (VibeX), 15 minutes of corneal soaking, UV-A power at  $12 \text{ mW/cm}^2$  (Energy dose:  $7.2 \text{ J/cm}^2$ ), 10 min exposure time.

All eyes re-epithelialized by 3 days of therapeutic soft contact lens bandage. Epithelial stratification improved in time, being complete at 3<sup>rd</sup> month. Sub-epithelial and anterior stromal nerves disappeared immediately after treatment. Nerves regeneration started one month after treatment being complete after 6 months. Anterior stromal tissue presented a high reflectivity after A-CXL with keratocytes loss (apoptosis hence photo-necrosis) until  $200 \mu\text{m}$  of depth and classical spongy or lacunar edema as previously demonstrated by us [20-22] in standard epithelium-off CXL was evident until 3<sup>rd</sup> month, gradually disappearing thereafter. Keratocytes repopulation started one month after treatment increasing at 3<sup>rd</sup> month and being complete at 6<sup>th</sup> post-operative month. An uneven demarcation line was determined at a mean depth of  $180 \mu\text{m}$  (range  $160\text{-}200 \mu\text{m}$ ) measured from epithelial surface (Figure 1). Confocal data of increased stromal reflectivity and

Group 1 Epithelium-Off A-CXL	Group 2 Epithelium-Off A-CXL	Group 3 Epithelium-Off A-CXL	Group 4 Epithelium-On A-CXL
5 eyes	5 eyes	5 eyes	5 eyes
16-23	13-26	14-24	21-26
Riboflavin 0.1% plus Dextran 20% (VibeX)	Riboflavin 0.1% plus Dextran 20% (VibeX)	Riboflavin 0.1% plus HPMC (VibeX Rapid)	Riboflavin 0.25% plus EDTA, BAK, TRIS (Paracel)
15 minutes	20 minutes	10 minutes	10 minutes
$12 \text{ mW/cm}^2$	$30 \text{ mW/cm}^2$	$30 \text{ mW/cm}^2$	$45 \text{ mW/cm}^2$
10 min	4 min	4 min	2 min. and 40 sec
$7.2 \text{ J/cm}^2$	$7.2 \text{ J/cm}^2$	$7.2 \text{ J/cm}^2$	$7.2 \text{ J/cm}^2$
$180 \mu\text{m}$ (range $160\text{-}200 \mu\text{m}$ )	$160 \mu\text{m}$ (range $150\text{-}180 \mu\text{m}$ )	$155 \mu\text{m}$ (range $140\text{-}180 \mu\text{m}$ )	$80 \mu\text{m}$ (range $50\text{-}120 \mu\text{m}$ )
$180 \mu\text{m}$ (range $150\text{-}200 \mu\text{m}$ )	$160 \mu\text{m}$ (range $150\text{-}180 \mu\text{m}$ )	$150 \mu\text{m}$ (range $140\text{-}180 \mu\text{m}$ )	noevident demarcation line

**Table 1:** Overview of treatment parameters and respective results according to demarcation lines measurements after *in vivo* scanning laser confocal microscopy and corneal OCT.

demarcation line (defined by corneal edema and keratocytes apoptosis with changes in stromal reflectivity) was established by anterior chamber OCT at a mean depth of 180  $\mu\text{m}$  (range 150-200  $\mu\text{m}$ ). The demarcation line was also clinically well evident after treatment at slit lamp examination (Figure 2). No endothelial damage was observed in terms of morphology and cell count after A-CXL.

### Group 2 epithelium-off A-CXL

Riboflavin 0.1% plus Dextran 20% (VibeX), 20 minutes of corneal soaking, UV-A power at 30  $\text{mW}/\text{cm}^2$  (Energy dose: 7.2  $\text{J}/\text{cm}^2$ ), 4 min exposure time.


The results in this group are the same of *Group 1* concerning epithelium and nerves regeneration. Some differences were found in the stromal analysis that showed a higher reflectivity of the anterior stromal tissue combined with keratocytes apoptosis and typical lacunar edema in the first three postoperative months followed by gradual cells repopulation. The demarcation line (defined by corneal edema and keratocytes apoptosis with changes in stromal reflectivity) at confocal scans was unevenly distributed in a mean depth of 160  $\mu\text{m}$  (range 150-180  $\mu\text{m}$ ) (Figure 3). No morphological changes in endothelial cells were observed. OCT imaging confirmed a mean depth of the demarcation line (defined by the higher reflectivity of cross-linked tissue) at 160  $\mu\text{m}$

(range 150-180  $\mu\text{m}$ ), moreover a demarcation line is clinically evident at bio-microscopy (Figure 4).


### Group 3 epithelium-off A-CXL

Riboflavin 0.1% (dextran free) plus HPMC (VibeX Rapid), 10 minutes of corneal soaking, UV-A power at 30  $\text{mW}/\text{cm}^2$  (Energy dose: 7.2  $\text{J}/\text{cm}^2$ ), 4 min exposure time.


The results in this group are superimposable with those observed in previous epithelium off treatment groups both concerning epithelial regrowth and nerves regeneration. While epithelium regenerates rapidly into 3 days, neural flocculation is detectable one month after treatment. The main differences were recorded in stromal healing where reflectivity was increased compared to preoperative scans but concentrated in the anterior 150  $\mu\text{m}$  of the stroma (Figure 5). In this case the riboflavin solution used for corneal soaking is dextran free, containing the hydroxyl-propyl-methyl-cellulose (HPMC) as riboflavin vehicle [24]. An uneven demarcation line is detectable at mean depth of 155  $\mu\text{m}$  (range 140-180  $\mu\text{m}$ ) and the reflectivity of extracellular matrix is relatively lower than those observed in group 1 and 2 patients. Demarcation line depth is confirmed by corneal OCT at a mean depth of 150  $\mu\text{m}$  (range 140-180  $\mu\text{m}$ ) (Figure 6). No endothelial damage is detectable in the postoperative period.




**Figure 1:** Accelerated corneal collagen crosslinking Group 1 (epithelium-off): Riboflavin 0.1% plus Dextran 20%, 15 min of corneal soaking, UV-A power at 12  $\text{mW}/\text{cm}^2$ , Energy dose at 7.2  $\text{J}/\text{cm}^2$ , 10 min exposure time. Left (a) white arrow: biomicroscopic picture of demarcation line after treatment. Right (b) white arrows: corneal OCT imaging revealing the demarcation line after A-CXL at about 180  $\mu\text{m}$  of depth (range 150-200  $\mu\text{m}$ ).




**Figure 3:** Accelerated corneal collagen crosslinking Group 2 (epithelium-off): Riboflavin 0.1% plus Dextran 20%, 20 min of corneal soaking, UV-A power at 30  $\text{mW}/\text{cm}^2$ , Energy dose at 7.2  $\text{J}/\text{cm}^2$ , 4 min exposure time. Left (a) white arrow: biomicroscopic picture of demarcation line after treatment. Right (b) white arrows: corneal OCT imaging revealing the demarcation line after A-CXL at about 160  $\mu\text{m}$  depth (range 150-180  $\mu\text{m}$ ).




**Figure 2:** Accelerated corneal collagen crosslinking Group 1 (epithelium-off): Riboflavin 0.1% plus Dextran 20%, 15 min of corneal soaking, UV-A power at 12  $\text{mW}/\text{cm}^2$ , Energy dose at 7.2  $\text{J}/\text{cm}^2$ , 10 min exposure time. Postoperative confocal scans revealed lacunar edema (upper left white arrow-a) with keratocytes apoptosis (upper right white arrow-b; bottom left white arrow-c) and increased density of extracellular matrix surrounding edema. A transition between hypo-cellular stroma and stroma unreached by the treatment (vertical transition area) is showed in bottom left - d scan with an estimated penetration depth of 180  $\mu\text{m}$  (range 160-200  $\mu\text{m}$ ).



**Figure 4:** Accelerated corneal collagen crosslinking Group 2 (epithelium-off): Riboflavin 0.1% plus Dextran 20%, 20 min of corneal soaking, UV-A power at 30  $\text{mW}/\text{cm}^2$ , Energy dose at 7.2  $\text{J}/\text{cm}^2$ , 4 min exposure time. Postoperative confocal scans revealed lacunar edema (upper left white arrow-a) with keratocytes apoptosis (upper right white arrow-b) and increased density of extracellular matrix (bottom left white arrow-c). A transition between hypo-cellular stroma and stroma unreached by the treatment (vertical transition area) is showed in bottom left - d scan with an estimated penetration depth of 160  $\mu\text{m}$  (range 150-180  $\mu\text{m}$ ).



**Figure 5:** Accelerated corneal collagen crosslinking Group 3 (epithelium-off): Riboflavin 0.1% (dextran free) plus HPMC, 10 minutes of corneal soaking, UV-A power at 30 mW/cm<sup>2</sup> (Energy dose: 7.2 J/cm<sup>2</sup>), 4 min exposure time. Left (a) light blue arrow: biomicroscopic picture of demarcation line after treatment. Right (b) light blue arrows: corneal OCT imaging revealing the demarcation line after A-CXL at about 150 μm depth (range 140-180 μm).



**Figure 6:** Accelerated corneal collagen crosslinking Group 3 (epithelium-off): Riboflavin 0.1% (dextran free) plus HPMC, 10 minutes of corneal soaking, UV-A power at 30 mW/cm<sup>2</sup> (Energy dose: 7.2 J/cm<sup>2</sup>), 4 min exposure time. Postoperative confocal scans revealed lacunar edema (upper left light blue arrow-a) with keratocytes apoptosis (upper right light blue arrow-b) and increased density of extracellular matrix (bottom left light blue arrow-c). A transition between the hypo-cellular stroma and stromal tissue unreached by the treatment (vertical transition area) is showed in the bottom left - d scan, with an estimated penetration of 155 μm (range 140-180 μm).


### Group 4 epithelium-on A-CXL

Riboflavin 0.25% plus EDTA, BAK,TRIS (Paracel) for 4 min, Riboflavin 0.25% saline solution (VibeX X-tra) for 6 min, UV-A power at 45 mW/cm<sup>2</sup> (Energy dose: 7.2 J/cm<sup>2</sup>), 2 min. 40 sec exposure time.


Epithelium-on treatment showed an acute actinic-like diffuse punctate epitheliopathy that was recovered following 3 days of soft contact lens bandage and sodium hyaluronate lubricants. Sub-epithelial nerves were damaged and partially disappeared after this high UV power setting, even if delivered with epithelium *in situ*. In any case stromal healing demonstrated poor apoptosis and sub-edema more diffuse than lacunar. A limited and uneven apoptotic affect is detectable after epithelium-on A-CXL in the anterior stroma at a mean depth of 80 μm (range 50-120 μm) (Figure 7). No endothelial damage was observed. OCT corneal scans confirmed a slightly increased reflectivity under the Bowman's lamina without an evident demarcation line. Demarcation line is not clinically visible after epithelium on A-CXL also at bio microscopic examination (Figure 8).

### Conventional epithelium-off CXL

Compared with epithelium-off ACXL groups, after epithelium-off standard CXL treatment (3 mW/cm<sup>2</sup> for 30 minutes of UV-A exposure),



**Figure 7:** Accelerated corneal collagen crosslinking Group 4 (epithelium-on): Riboflavin 0.25% plus EDTA, BAK,TRIS for 4 min, Riboflavin 0.25% saline solution for 6 min, UV-A power at 45 mW/cm<sup>2</sup> (Energy dose: 7.2 J/cm<sup>2</sup>), 2 min. 40 sec exposure time. Left (a) yellow arrow: biomicroscopic picture after epithelium-on accelerated cross-linking showed a slight increased density of sub-Bowman stroma without a visible demarcation line. Right (b) yellow arrows: OCT corneal scans confirmed a slightly increased reflectivity under of corneal stroma under the Bowman's lamina without evidence of demarcation line.



**Figure 8:** Accelerated corneal collagen crosslinking Group 4 (epithelium-on): Riboflavin 0.25% plus EDTA, BAK,TRIS for 4 min, Riboflavin 0.25% saline solution for 6 min, UV-A power at 45 mW/cm<sup>2</sup> (Energy dose: 7.2 J/cm<sup>2</sup>), 2 min. 40 sec exposure time. Postoperative confocal scans revealed a superficial diffuse edema under the Bowman's lamina with rarefaction and damage of sub-epithelial nerves fibers (upper left yellow arrows - a); an uneven keratocytes apoptosis is detectable in the anterior stroma (upper right and bottom right yellow arrows - b and d) with a vertical transition area (bottom left yellow arrow - c) at 80-90 μm of depth on average (range 50-120 μm).

epithelial healing and nerves regeneration were superimposable, being completed respectively after 3-4 days and 3-6 months. In A-CXL, keratocytes apoptosis reached the anterior-mid corneal stroma until 150-200 μm instead of 250-300 microns of the classic CXL treatment. Cell apoptosis after CXL was more evident after soft contact lens removal and along the first postoperative month. The apoptotic process after CXL treatment required at least 48-72 hours, becoming well evident at *in vivo* confocal scans (apoptotic bodies) along the first post-operative month. In the first week after treatment it was masked by the presence of marked honeycomb-like stromal edema. Reflectivity of extracellular matrix in the first 3 months was slightly higher in the epithelium-off A-CXL patients compared with standard epithelium-off CXL. No endothelial damage was observed in both treatments modalities.

### Standard epithelium-on (TE-CXL)

Compared with epithelium-on treatment performed by us with Riboflavin 0.1% plus Dextran 15%, EDTA and Trometamol solution, at 3 mW/cm<sup>2</sup> for 30 minutes, after A-CXL we recorded the same superficial,

diffuse and irregular epithelial photo-chemical damage. Sub-epithelial nerves plexus was present after classic epithelium on treatment, while nerve fibers disappearance was evident after A-CXL. The timing of nerves fibers regeneration after Epi-on ACXL was similar to standard epithelium-off CXL (3-6 months). Keratocytes apoptosis was uneven and confined under 80 microns of depth and stromal edema was unevenly distributed under the Bowman lamina in the anterior stroma. No endothelial damage was observed in both techniques.

## Discussion

*Epithelium-off* A-CXL demonstrated in the first 3 groups morphological changes and treatment penetration defined by stromal edema and keratocytes loss at *in vivo* confocal microscopy and by increased stromal reflectivity at AC OCT, comprised between 150 and 180  $\mu\text{m}$  on average (range 140-200  $\mu\text{m}$ ).

As reported in literature [25] *in vivo* UV-A induced oxidative damage (apoptotic effect and cell viability) depends on the energy, riboflavin concentration and mode of exposure. In this context, the exposure time together with riboflavin concentration become very important in cross-linking treatment (interactions between UV-A photons, riboflavin and collagen).

Even if UV-A intensity is increased while maintaining a constant energy dose (5.4 or 7.2  $\text{J}/\text{cm}^2$ ), a prolonged exposure time influenced a deeper penetration of oxidative damage [25], increasing treatment volume, like demonstrated in our first protocol at 12  $\text{mW}/\text{cm}^2$  for 10 minutes of exposure time (Group 1) that reported a mean penetration of 180  $\mu\text{m}$  (Figures 1 and 2). The A-CXL protocol at 30  $\text{mW}/\text{cm}^2$  for 4 minutes of exposure time (Group 2) revealed a mean penetration of 160  $\mu\text{m}$  both at confocal analysis and corneal OCT (Figures 3 and 4), slightly inferior to Group 1.

This result is slightly better with those recently reported in literature by Colin research group [26] probably due to high energy dose that we used in our treatments according to Avedro's laboratory data (7.2  $\text{J}/\text{cm}^2$  instead of 5.4  $\text{J}/\text{cm}^2$ ).

The clinical aspect of the corneas after A-CXL was good after therapeutic soft contact lens removal and in the first postoperative month without any complication such as persistent epithelial defects or haze. A demarcation line was clearly visible in all epithelium-off A-CXL treatments at slit lamp examination just after therapeutic soft contact lens removal (Figures 2,4 and 6).

Keratocytes apoptosis correlating with treatment penetration [23] was limited to the anterior-mid stroma until a maximum depth of 200  $\mu\text{m}$  if compared with conventional Dresden protocol that reached 300  $\mu\text{m}$  without epithelium as well demonstrated by our first confocal studies *in vivo* in humans [20-22]. On the other hand, the intensity of extracellular matrix after epithelium-off A-CXL resulted higher in the anterior 150  $\mu\text{m}$  of stroma suggesting a good collagen compaction and corneal stiffening with reduced corneal edema and less cell toxicity (Figure 3). The higher reflectivity recorded in group 1 and 2 may be explained by the higher tissue dehydration after A-CXL by using riboflavin 0.1% plus Dextran 20% (VibeX) solution.

As reported in literature the most important biomechanical effect related to crosslinking is concentrated in the anterior 200  $\mu\text{m}$  of the cornea [27], in the so called stiff cornea, so the impact of A-CXL may be sufficient in terms of biomechanical and biochemical effect.

A relatively low reflectivity of extracellular matrix was observed in the Group 3 protocol (Figures 5 and 6), compared with Group 1 and

2 patients (Figures 1 and 3), that may be explained by the different (dextran free VibeX Rapid) riboflavin solution used for corneal soaking, containing the hydroxyl-propyl-methyl-cellulose (HPMC) as riboflavin vehicle, reducing intraoperative corneal dehydration [28].

Epithelium on A-CXL demonstrated a powerful toxic effect on epithelium related to enhanced riboflavin solutions containing ethylene-diamine-tetra-acetic acid (EDTA), benzalkonium chloride (BAK), trometamol (TRIS) and also to high UV-A intensity at 45  $\text{mW}/\text{cm}^2$  delivered in a very short time (2 min and 40 sec) on epithelial cells producing an immediate, even short, postoperative patient discomfort. Moreover higher UV-A intensity, even if delivered with epithelium *in situ*, induced a slight damage of sub-epithelial plexus nerves probably due to altered condition of epithelial surface itself. In any case the stromal healing after epithelium-on A-CXL demonstrated poor cells apoptosis and sub-edema, more diffuse than lacunar. A limited and uneven apoptotic effect is detectable after epithelium-on A-CXL in the anterior stroma (Figure 7), and a demarcation line is not visible (Figure 8), confirming that epithelium leaved *in situ* and the high intraepithelial riboflavin concentration represent a barrier for the UV-A diffusion into the stroma that is essential for cross-linking penetration, inducing a superficial oxidative damage (surface CXL). Also in epithelium-on A-CXL, like in the previous trans-epithelial CXL procedures [29-31], the first analysis *in vivo* in humans by using confocal microscopy [23], demonstrated that the presence of corneal epithelium *in situ* constitutes a physical barrier to UV-A radiation reducing its penetration into the corneal stroma and the results of present qualitative analysis confirm this finding. The low penetration of the epithelium on Accelerated CXL could not stabilize biomechanically the keratoconic cornea in the mid-long term follow-up as demonstrated in literature [32] by us after standard trans-epithelial procedure (TE-CXL).

To date we don't know exactly the optimal interactions between UV-A energy, riboflavin concentration and exposure time in order to obtain the maximum cross-linking effect ensuring a long-lasting (possibly a long-life) keratoconus stability and the better functional outcome, even if the necessity to improve the procedure and shorten the CXL treatment time are highly desirable.

In any case, the conventional epithelium-off CXL procedure (Riboflavin 0.1% plus dextran 20%, UV-A 3  $\text{mW}/\text{cm}^2=5.4 \text{ J}/\text{cm}^2$  for 30 minutes) remains the gold standard in the conservative treatment of early stages progressive keratoconus. On the other hand the rule of Bunsen-Roscoe's law of reciprocity, established for photo-chemical reactions, cannot be directly transferred in terms of photo-biological effect to complex biological systems as the living cornea [25].

Accelerated cross-linking with epithelium removal demonstrated its safety for endothelium and posterior ocular structures. Treatment penetration achieve the anterior part of the stromal tissue stiffening the cornea in the first 160  $\mu\text{m}$  on average (range 140-200  $\mu\text{m}$ ) with relative differences between the different protocols we used.

In our experience, A-CXL shortened CXL procedure under 20 minutes being well tolerated by patients. However its clinical efficacy, both in terms of keratoconus stabilization and functional impact, must be determined in the mid-long term follow-up and in a large cohort of patients according to different patient's age, keratoconus stage and progression rate.

## Financial Disclosure

The authors declare that they have no financial interest in the manuscript.

## References

1. Wollensak G, Spoerl E, Seiler T (2003) Riboflavin/ultraviolet-a-induced collagen crosslinking for the treatment of keratoconus. *Am J Ophthalmol* 135: 620-627.
2. Caporossi A, Baiocchi S, Mazzotta C, Traversi C, Caporossi T (2006) Parasurgical therapy for keratoconus by riboflavin-ultraviolet type A-induced cross-linking of corneal collagen: preliminary refractive results in an Italian study. *J Cataract Refract Surg* 32: 837-845.
3. Hafezi F, Kanellopoulos J, Wiltfang R, Seiler T (2007) Corneal collagen crosslinking with riboflavin and ultraviolet A to treat induced keratectasia after laser *in situ* keratomileusis. *J Cataract Refract Surg* 33: 2035-2040.
4. Wollensak G, Spoerl E, Seiler T (2003) Stress-strain measurements of human and porcine corneas after riboflavin-ultraviolet-A-induced cross-linking. *J Cataract Refract Surg* 29: 1780-1785.
5. Wollensak G, Wilsch M, Spoerl E, Seiler T (2004) Collagen fiber diameter in the rabbit cornea after collagen crosslinking by riboflavin/UVA. *Cornea* 23: 503-507.
6. Spoerl E, Wollensak G, Seiler T (2004) Increased resistance of crosslinked cornea against enzymatic digestion. *Curr Eye Res* 29: 35-40.
7. Kamaev P, Friedman MD, Sherr E, Muller D (2012) Photochemical kinetics of corneal cross-linking with riboflavin. *Invest Ophthalmol Vis Sci* 53: 2360-2367.
8. Spoerl E, Huhle M, Seiler T (1998) Induction of cross-links in corneal tissue. *Exp Eye Res* 66: 97-103.
9. Spoerl E, Seiler T (1999) Techniques for stiffening the cornea. *J Refract Surg* 15: 711-713.
10. Raiskup-Wolf F, Hoyer A, Spoerl E, Pillunat LE (2008) Collagen crosslinking with riboflavin and ultraviolet-A light in keratoconus: long-term results. *J Cataract Refract Surg* 34: 796-801.
11. Caporossi A, Mazzotta C, Baiocchi S, Caporossi T (2010) Long-term results of riboflavin ultraviolet a corneal collagen cross-linking for keratoconus in Italy: the Siena eye cross study. *Am J Ophthalmol* 149: 585-593.
12. Wittig-Silva C, Whiting M, Lamoureux E, Lindsay RG, Sullivan LJ, et al. (2008) A randomized controlled trial of corneal collagen cross-linking in progressive keratoconus: preliminary results. *J Refract Surg* 24: S720-725.
13. Caporossi A, Mazzotta C, Baiocchi S, Caporossi T, Denaro R (2011) Age-Related Long-Term Functional Results after Riboflavin UV A Corneal Cross-Linking. *J Ophthalmol* 2011: 608041.
14. Caporossi A, Mazzotta C, Baiocchi S, Caporossi T, Denaro R, et al. (2012) Riboflavin-UVA-induced corneal collagen cross-linking in pediatric patients. *Cornea* 31: 227-231.
15. Spoerl E, Mrochen M, Sliney D, Trokel S, Seiler T (2007) Safety of UVA-riboflavin cross-linking of the cornea. *Cornea* 26: 385-389.
16. BRINDLEY GS (1952) The Bunsen-Roscoe law for the human eye at very short durations. *J Physiol* 118: 135-139.
17. Schumacher S, Oeftiger L, Mrochen M (2011) Equivalence of biomechanical changes induced by rapid and standard corneal cross-linking, using riboflavin and ultraviolet radiation. *Invest Ophthalmol Vis Sci* 52: 9048-9052.
18. Wernli J, Schumacher S, Spoerl E, Mrochen M (2013) The efficacy of corneal cross-linking shows a sudden decrease with very high intensity UV light and short treatment time. *Invest Ophthalmol Vis Sci* 54: 1176-1180.
19. Celik HU, Alagöz N, Yildirim Y, Agca A, Marshall J, et al. (2012) Accelerated corneal crosslinking concurrent with laser *in situ* keratomileusis. *J Cataract Refract Surg* 38: 1424-1431.
20. Mazzotta C, Traversi C, Baiocchi S, Sergio P, Caporossi T, et al. (2006) Conservative treatment of keratoconus by riboflavin-uva-induced cross-linking of corneal collagen: qualitative investigation. *Eur J Ophthalmol* 16: 530-535.
21. Mazzotta C, Balestrazzi A, Traversi C, Baiocchi S, Caporossi T, et al. (2007) Treatment of progressive keratoconus by riboflavin-UVA-induced cross-linking of corneal collagen: ultrastructural analysis by Heidelberg Retinal Tomograph II *in vivo* confocal microscopy in humans. *Cornea* 26: 390-397.
22. Mazzotta C, Traversi C, Baiocchi S, Caporossi O, Bovone C, et al. (2008) Corneal healing after riboflavin ultraviolet-A collagen cross-linking determined by confocal laser scanning microscopy *in vivo*: early and late modifications. *Am J Ophthalmol* 146: 527-533.
23. Caporossi A, Mazzotta C, Baiocchi S, Caporossi T, Paradiso AL (2013) Trans-epithelial corneal collagen crosslinking for keratoconus: qualitative investigation by *in vivo* HRT II confocal analysis. *Eur J Ophthalmol* 22: S81-88.
24. Mazzotta C, Baiocchi S, Caporossi T, Caragiuli S, Paradiso AL, et al. (2013) Riboflavin 0.1% (VibeX) for the treatment of keratoconus. *Expert Opinion on Orphan Drugs* 1: 235-240.
25. Merwald H, Klosner G, Kokesch C, Der-Petrossian M, Hönigsmann H, et al. (2005) UVA-induced oxidative damage and cytotoxicity depend on the mode of exposure. *J Photochem Photobiol B* 79: 197-207.
26. Touboul D, Efron N, Smadja D, Praud D, Malet F, et al. (2012) Corneal Confocal Microscopy Following Conventional, Transepithelial, and Accelerated Corneal Collagen Cross-linking Procedures for Keratoconus. *J Refract Surg* 28: 769-776.
27. Kohlhaas M, Spoerl E, Schilde T, Unger G, Wittig C, et al. (2006) Biomechanical evidence of the distribution of cross-links in corneas treated with riboflavin and ultraviolet A light. *J Cataract Refract Surg* 32: 279-283.
28. Kymionis GD, Kounis GA, Portaliou DM, Grentzelos MA, Karavitaki AE, et al. (2009) Intraoperative pachymetric measurements during corneal collagen cross-linking with riboflavin and ultraviolet A irradiation. *Ophthalmology* 116: 2336-2339.
29. Leccisotti A, Islam T (2010) Transepithelial corneal collagen cross-linking in keratoconus. *J Refract Surg* 26: 942-948.
30. Filippello M, Stagni E, O'Brart D (2012) Transepithelial corneal collagen crosslinking: bilateral study. *J Cataract Refract Surg* 38: 283-291.
31. Koppen C, Wouters K, Mathysen D, Rozema J, Tassignon MJ (2012) Refractive and topographic results of benzalkonium chloride-assisted transepithelial crosslinking. *J Cataract Refract Surg* 38: 1000-1005.
32. Caporossi A, Mazzotta C, Paradiso AL, Baiocchi S, Marigliani D, et al. (2013) Transepithelial corneal collagen crosslinking for progressive keratoconus: 24-month clinical results. *J Cataract Refract Surg* 39: 1157-1163.

**Citation:** Mazzotta C, Paradiso AL, Baiocchi S, Caragiuli S, Caporossi A (2013) Qualitative Investigation of Corneal Changes after Accelerated Corneal Collagen Cross-linking (A-CXL) by *In vivo* Confocal Microscopy and Corneal OCT. *J Clin Exp Ophthalmol* 4: 313. doi: [10.4172/2155-9570.1000313](https://doi.org/10.4172/2155-9570.1000313)

## Submit your next manuscript and get advantages of OMICS Group submissions

### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

### Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: [www.editorialmanager.com/clinicalgroup](http://www.editorialmanager.com/clinicalgroup)

