

Article

Generation of the Quaternary Normal Faults in the Messina Strait (Italy)

Enzo Mantovani, Marcello Viti * , Daniele Babbucci and Caterina Tamburelli

Department of Physical Sciences, Earth and Environment, University of Siena, 53100 Siena, Italy; enzo.mantovani@unisi.it (E.M.); babbucci@unisi.it (D.B.); tamburelli@unisi.it (C.T.)

* Correspondence: marcello.viti@unisi.it

Abstract: It is widely recognized that since the Early–Middle Pleistocene, the Messina zone, the site of strong earthquakes, has undergone extension, but the geodynamic context which determined this deformation is still a matter of debate. This work suggests that such a tectonic event was caused by the interaction of northern Calabria with the continental Adriatic domain. The suture of that consuming boundary produced major changes in the microplate mosaic and the related kinematic pattern in the Southern Italian zones, which was triggered by the activation of the Sibari and Vulcano faults. In the new context, the Peloritani belt sector, dragged by the Hyblean block, rotated clockwise and then moved northward, causing its divergence from southern Calabria. The normal faults which have accommodated that separation may be the main seismogenic source in the Messina Strait.

Keywords: Messina Strait; seismic source; Quaternary evolution; tectonic setting; central Mediterranean



Citation: Mantovani, E.; Viti, M.; Babbucci, D.; Tamburelli, C. Generation of the Quaternary Normal Faults in the Messina Strait (Italy). *Geosciences* **2023**, *13*, 248. <https://doi.org/10.3390/geosciences13080248>

Academic Editors: Riccardo Caputo and Jesus Martinez-Frias

Received: 9 June 2023

Revised: 3 August 2023

Accepted: 14 August 2023

Published: 17 August 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Since the Messina Strait could become the site of a long bridge between Sicily and Calabria, recognizing the location and geometry of the main seismic sources in that area may be particularly useful. Several attempts to get this information have been made by studying the effects of the last two main earthquakes which affected the area, one in the northern Messina trough ($M = 6.2$) on 6 February 1783 (e.g., [1]) and one in the Messina sphenocasm ($M = 7.1$) on 28 December 1908 [2–15]. Some of the fault models indicated by the above authors for the most studied event, the 1908 shock, are shown in Figure 1. The variety of the proposed solutions testifies to the poor constraining power of the evidence now available, mostly represented by coseismic effects, with particular regard to levelling data from 1907 to 1909 [16]. The most striking ambiguity is given by the fact that the set of proposed solutions involves both onshore and offshore fault locations. Such difference mostly derives from the choice of considering or not the capable faults identified by morphological and geological observations (Figure 2, e.g., [13,17–20]). To mitigate the above uncertainty, it would be necessary to identify further possible constraints. In this work, we approach this problem by trying to recognize the geodynamic context that generated the normal faults in the Messina area. Since several hypotheses have been advanced about that problem, involving the gravitational sinking of the Ionian slab [21–24], deep-seated asthenosphere processes [25,26], extrusion processes [27–29] or other tectonic mechanisms (e.g., [20,30]), we report some remarks on how the implications of the proposed geodynamic models can account for the observed deformation pattern in the study area. To increase the constraining power of this analysis we have imposed that the tectonic context that determined the Pleistocene evolution of the study area was compatible with the processes that controlled the Pliocene deformation pattern in the central Mediterranean area.




Figure 1. Fault locations (red lines) and cross sections, with related deep fault geometries, adopted in the previous literature ((a) [2]; (b) [3]; (c) [4]; (d) [5,7,10]; (e) [6,8,12]; (f) [11]; (g) [13]; (h) [14]) to model the geodetic levelling dataset [16] carried out before and after the 1908 earthquake ([13,14] and references therein). The location of the area here considered is shown in the last picture (yellow box).




Figure 2. Quaternary normal faults (red lines) in the Messina Strait and surroundings indicated by geological and morphological evidence (e.g., [20,31–33]). AF = Armo fault; CF = Cittanova fault; CP = Capo Peloro fault; MF = Messina fault; MTF = Messina–Taormina fault; MSF = Messina Strait fault; RCF = Reggio Calabria fault; SEF = Sant’Eufemia fault; SF = Scilla fault. The star indicates the epicentre of the 1908 earthquake.

2. Pliocene

The Neogene evolution of the Mediterranean region has been driven by two main kinematic boundary conditions: the Nubia–Eurasia convergence and the westward escape of Anatolia (induced by the Arabia indentation [34–37]). Until the Late Miocene–Early Pliocene, the evolution of the central Mediterranean area was scarcely influenced by the escape of Anatolia since the convergence between these two major systems was mainly accommodated by the consumption of the interposed thinned zone (Pindos, [28,29,38–40]). The tectonic context changed considerably since the Early Pliocene when the Aegean system (Hellenides) collided with the Adriatic foreland. After the consequent suture of this consuming boundary, the westward push of the Anatolian–Aegean–Pelagonian system (AAPS) caused the decoupling of a large part of the Adriatic promontory (Adria plate) from Nubia, by the activation of a major fracture, constituted by the Victor Hensen–Medina–Sicily Channel fault systems. Once decoupled, this plate underwent a clockwise rotation accompanied by a NNW ward motion (Figure 3B, [27–29,41]).




Figure 3. (A) Tectonic configuration of the study area in the Late Miocene–Early Pliocene, after the formation of the Northern Tyrrhenian basin. Ge = Gela basin; NA, SA = Northern and Southern Apennines. (B) Middle Pliocene. After the collision of the Anatolian–Aegean–Pelagonian system (AAPS) with the continental Adriatic domain, a large part of the Adriatic promontory (Adria plate) decoupled from Nubia by the activation of the Victor Hensen–Medina (VHM) and Sicily Channel (SCH)

fault system. Once decoupled, the Adria plate underwent clockwise rotation and minor NW to NNW ward motion. The consequent convergence between the Adria plate and Northwestern Nubia was accommodated by major E-W shortening in the Pelagian zone, as well as in the Alpine–Maghrebian belt. Aw = Adventure wedge; Eg = Egadi fault; GN = Gela nappe; Sc = Sciacca fault; SR = Scicli–Ragusa fault; SV = Schio–Vicenza fault; VB = Vavilov basin (Central Tyrrhenian). (C) Early Pleistocene. After the collision of the Southern Apennine extruding wedge with the Apulian swell (AS), accretionary activity along the outer Calabrian front formed the External Calabrian accretionary belt (ECA), while crustal extension developed in the wake of the migrating Arc, forming the Marsili basin (Ma). In this strong compressional context, the Calabria–Peloritani (CP) wedge underwent fast uplift and bowing. Pa = Palinuro fault, Ta = Taormina fault. (1) Continental (a) and thinned continental; (b) Eurasian domains. (2) Continental (a) and thinned continental; (b) African/Adriatic domains. (3) Tethyan belt (AAPS). (4) Other orogenic belts. (5) Old oceanic domains. (6) Zones affected by intense (a) or moderate (b) crustal thinning (7–9). Compressional, extensional and strike-slip tectonic features. Blue arrows indicate the kinematic pattern with respect to Europe [42,43].

This evolutionary phase involved a considerable change in the tectonic setting in the southern part of the central Mediterranean area, as shown in greater detail in Figure 4.

The roughly E-W convergence between the southernmost Adria plate and Northwestern Nubia (Tunisia) induced strong compression in the Pelagian–Hyblean domain, which was accommodated by some major shortening processes [27–29,44–49]. A major one was the roughly northward escape of the Adventure wedge, guided by the Sciacca and Egadi lateral faults (Figure 4B). The extension that developed in the wake of this extruding wedge formed the Pantelleria trough, while the compressional deformation that occurred along the front of the wedge was accommodated by thrusting and northward bending/displacement of the adjacent Alpine–Maghrebian belt (Figure 5). Such deformation caused the interaction of the Maghrebian belt with the Corsica–Sardinia block [28,29]. This hypothesis can explain why during that evolutionary phase (Lower-middle Pliocene to the Pleistocene) tectonic activity affected the southwestern edge of the Corsica–Sardinia block, with the formation of the Campidano trough (e.g., [50–52]). An alternative explanation of that event can hardly be found, since the Corsica–Sardinia block did not experience any other interaction with the surrounding structures after the Middle Miocene [53–55].

Another major effect of the convergence between the Adria plate and northern Nubia was the consumption of the inner thinned part of the Hyblean domain (Gela basin, Figure 3A), which was accommodated by two opposite processes. One was the southward oroclinal bending of the Sicilian Apennines (Gela nappe) and the other resulted from the northward displacement of the fragments (Empedocle, Girgenti, Linosa, Malta, Selinunte, Urialo in Figure 4A) generated by the breaking of a narrow Hyblean plateau. The extension that developed in the wake of the escaping fragments generated the Linosa and Malta troughs (Figure 4B, [28,29]).

The present shape of the Kabyliides–Calabrides and Maghrebides–Apennines belts, determined by the northward escape of the Adventure–Hyblean wedge, and other main tectonic features in the Pelagian–Hyblean domain are shown in Figure 5. The proposed interpretation, involving the lateral escape of the Adventure wedge and the consequent deformation of the Maghrebian belt, can account for a peculiar aspect of the Central Mediterranean tectonic setting, i.e., the fact that the Maghrebian belt lying in front of the Adventure wedge is the only sector of the peri-Tyrrhenian belts that moved towards the Corsica–Sardinia block, while all the other belt sectors (Calabrian Arc and Apennines) moved away from Corsica–Sardinia.




Figure 4. Evolutionary reconstruction of the Hyblean–Pelagian domain and the adjacent Alpine–Maghrebian belt, as effect of the convergence between the Adria plate and northwestern Nubia. (1) Continental domains. (2) Thinned continental and oceanic domains. (3) Orogenic belts. (4) Zones of intense (a) or moderate (b) crustal thinning. (A) Late Miocene. Initial configuration of

the study area. Cal = Calabrian wedge; Ad, Em, Gi, Hy, Li, Ma, Se, Ur = Empedocle–Girgenti–Linosa–Malta–Selinunte–Urialo original parts of the Hyblean plateau; Ge = Gela basin (thinned Hyblean domain); Hel = Hellenides; Pa = Palinuro fault, Sy = Syracuse escarpment, Ta = Taormina fault. (B) Middle-upper Pliocene. Aw = Adventure wedge, Ca = Campidano graben, Eg = Egadi fault, GN = oroclinal Gela Nappe; Lit, Mat, Plt = Linosa, Malta and Pantelleria troughs; SCH = Sicily channel fault, Sc = Sciacca fault; SR= Scicli–Ragusa fault; Va = Vavilov basin; VHM = Victor Hensen–Medina fault (C) Late Pliocene–Early Pleistocene. The suture of the Southern Apennines consuming boundary accelerated the lateral escape of the Calabrian wedge, triggering the formation of the Marsili basin (Mar) and the External Calabrian Arc (ECA), Ce = Cefalonia fault, (D) Middle Pleistocene–Present. The collision of the northern Calabrian wedge with the Adriatic foreland triggered the formation of the Sibari (Si) and Vulcano (Vu) faults. The geometries and locations of the Hyblean plateau fragments have been taken from [56]. PM = Peloritani microplate. Other symbols as in Figure 3. More details about the Pleistocene evolution are reported in Figures 6 and 7.




Figure 5. Present tectonic setting in the study area. Kabyrides–Calabrides (1) and Maghrebides–Apennines (2) belts (modified after [46,57,58]); (3) foreland domains, (4–6) strike-slip, compressional and tensional features. SA = Southern Apennines. The fact that the Kabyrides–Calabrides and Maghrebides–Apennines belts, deformed by the escape of the Adventure wedge have reached the Corsica–Sardinia block may explain why this last structure underwent tectonic activity in the Pliocene.

3. Early Pleistocene–Present

Another major reorganization of the tectonic pattern occurred around the Early–Middle Pleistocene when the lateral escape of the Calabrian wedge reached the Adriatic foreland (Figure 4d). After this continental collision, the extrusion trend of that wedge had to change, to favour the consumption of the remaining Ionian thinned domain (e.g., [28,29]). This deviation was allowed by the activation of two new lateral guides, the Sibari and Vulcano faults (Figure 6b, e.g., [18,20,59–64]), which replaced the previous guides (Taormina and Palinuro faults). The formation of the Vulcano fault also had major effects on the

kinematics of the Hyblean block, which accelerated and moved with a greater northward component. Initially, the dextral transpression at the Vulcano fault did not allow a complete decoupling between the Hyblean block and the Peloritani belt sector (PM = Peloritani microplate). As a consequence, this microplate, partially dragged by the Hyblean block, rotated clockwise with a minor northward component, so diverging from southern Calabria (Figures 4, 6a and 7). This reconstruction can account for the generation of faults in the Messina sphenocasm (MF, MTF, RCF and AF faults in Figure 2) and then of the northern trough (MSF and SF faults in Figure 2). The supposed northward motion of PM can also explain the occurrence, since the Early–Middle Pleistocene, of compressional deformation along the outer front of this microplate, as indicated by the development of the Capo Peloro transpressional fault (Figure 2, e.g., [20]). Moreover, in the same period, a NE-trending anticline grew up a few km north of the Messina strait [20]. This structural high is considered by some authors [65] as the submarine prolongation of an uplifting area (1.1 mm/y) in PM. It is noteworthy that this anticline developed in the southernmost sector of the internal Calabrian arc, which was previously characterized by extension. This evidence and the formation of ridges characterized by high uplift rates [65] clearly reveal that such a zone underwent an inversion of strain regime since the Middle Pleistocene. The Quaternary compressional deformation observed in the Maghrebides lying offshore northern Sicily (e.g., [46,66–71]) is compatible with the supposed roughly northward motion of the Hyblean block (Figure 6). The important decoupling role that is still played by the Syracuse fault is testified by the occurrence of strong historical earthquakes offshore eastern Sicily (e.g., the 1169 M = 6.5 and 1693 M = 7.3 shocks) [72].




Figure 6. Tectonic sketch of the Calabria–Peloritani (CP) and Hyblean (Hy) wedges before (A) and after (B) the collision of Northern Calabria with the Adriatic continental domain. (1) Nubian continental domain; (2) Adriatic continental domain; (3) Ionian oceanic domain; (4) Orogenic belts; (5) Peloritani microplate; (6) Outer fronts of the Alpine belt. (A) Early–Middle Pleistocene. The lateral escape of the CP wedge is guided by the Taormina (Ta) and Palinuro (Pa) faults. ECA = External Calabrian Arc; NH = Northern Hellenides; Sy = Syracuse fault; Pe = Peloritani. (B) Post Middle Pleistocene. The extrusion trend of the CP wedge changed significantly by the activation of two new lateral guides (Sibari (Si) and Vulcano (Vu) faults (e.g., [18,46,56])).




Figure 7. Tentative reconstruction of the microplate kinematics and tectonic setting which led to the formation of the Messina trough (MT). (a) Late Pliocene. The Calabria–Peloritani wedge (CP) extrudes laterally, guided by the Palino (Pa) and Taormina (Ta) faults at the expense of the Ionian domain. Hy = Hyblean domain. (b) Early–Middle Pleistocene. After the activation of the Vulcano (Vu) and Sibari (Si) faults, the extrusion trend of CP changes, while Hy accelerates, moving with a greater northward component. PM = Peloritani microplate. (c) Late Pleistocene–Present. Stressed by the transpressional interaction with Hy, PM undergoes clockwise rotation and then a minor northward motion. The consequent divergence of this microplate from Southern Calabria generates normal faults in the Messina zone (Figure 2). The present tectonic setting shown in (f) is taken from the Tectonic Map of Italy [73]. Using the same Map, the location of that tectonic setting is tentatively reconstructed for the previous evolutionary phases (d,e).

4. Discussion

This work describes the Plio–Quaternary tectonic evolution of the central Mediterranean region that in our opinion can best account for the Pleistocene occurrence of extension in the Messina Strait. Since various alternative geodynamic interpretations have been so far proposed for this region, it may be useful to explain why the solution proposed here can be preferred to the hypotheses previously advanced.

The interpretation most cited in the literature (under the name of the slab-pull model, e.g., [44–47]) suggests that a primary role in the Plio–Quaternary evolution of the central Mediterranean was played by the gravitational sinking of the Ionian slab, which deter-

mined the migration of the Calabrian Arc and the opening of the Tyrrhenian basin. The main difficulties in reconciling the implications of this driving force with the available evidence are discussed in various papers (e.g., [27–29,41,43,74]). Here, we report further considerations about some major tectonic features that can impose important constraints on the geodynamic context.

The fast uplift that since the middle Pleistocene, the Calabrian Arc has undergone (more than 1200 m, with an increasing rate, e.g., [31,75–79]) cannot easily be reconciled with the implications of the slab-pull model that are predicted by modellings, with particular regard to the occurrence of major subsidence in the migrating arc (e.g., [80–84]).

An important piece of evidence that can help us to identify the driving force of the Calabrian wedge is provided by the cross-section shown in Figure 8A, which shows a vertical bending of crustal layers in the Calabrian Arc. This peculiar deformation, indicated by upward and downward displacements, testifies to the very strong horizontal compression that such a structure must have undergone during the development of the Tyrrhenian–Calabrian arc-trench-back arc system. Such shortening is clearly incompatible with the horizontal extension of the migrating arc that is expected from the slab-pull driving force [80,81,83,84]. In our opinion, the above major features should discourage any attempt to invoke the slab-pull mechanism as the driving force of the Calabrian evolution.

It is widely recognized that the northern Tyrrhenian basin developed from the Tortonian to the late Miocene, the central Tyrrhenian (Vavilov basin) from the Early Pliocene to the Late Pliocene and the southernmost Tyrrhenian (Marsili basin) since Pleistocene (e.g., [56,85–89]). Explaining this peculiar space–time evolution of that basin as an effect of slab sinking must face many difficulties, as discussed in some papers [27–29,41,74]. In particular, one should explain why the gravitational sinking of the subducted lithosphere first developed in the northern Tyrrhenian sector, despite that during such period (upper Miocene) the underlying slab was more developed in the southernmost sector of the migrating arc.




Figure 8. Cross sections from the southernmost Tyrrhenian to the Adriatic foreland, through Northern Calabria (A) and the Southern Apennines (B), modified after [90,91]. (1) Oceanic crust; (2) upper continental crust; (3) lower continental crust; (4) lithospheric mantle; (5) asthenospheric mantle; (6) remnants of the Alpine Tethys; (7) Kabylo–Calabrides units; (8) Ionian Tethys oceanic slab; (9) Ionides; (10) pre-Pliocene sedimentary cover; (11) Pliocene–Quaternary sediments; (12–14) major thrust, normal and strike-slip faults. Dots indicate the Adriatic crustal units (from [29], modified).

The tectonic features cited above are instead compatible with the hypothesis that the trench-arc-back arc systems in the central Mediterranean region developed as effects of extrusion processes (e.g., [27–29]), induced by the convergence of the confining plates (Nubia, Eurasia and Arabia). In fact, both the considerable uplift of Calabria and the vertical bending of its crustal structure may be explained as a consequence of the strong constriction that stressed the Calabrian wedge after the suture of the Southern Apennines consuming boundary. In that context, the above wedge was forced to escape laterally through a narrow corridor, confined by lateral continental domains (Hyblean plateau on one side and Adriatic domain on the other side, Figure 3). It can be noted that the section crossing the southern Apennine wedge (Figure 8B) does not show the vertical bending of the crust given in Figure 8A, indicating that the compressional stress which drove the lateral escape of that Apennine sector was less intense than the one controlling the extrusion of the Calabrian wedge.

Some authors tentatively interpret the uplift and extension in the Messina zone as an effect of deep-seated asthenospheric processes (e.g., [92–94]), with particular reference to mantle upwelling (e.g., [25,95]). However, it is very difficult to believe that such kind of process, involving the mobility of large deep structures, may have created the observed upper crustal tectonic setting, characterized by coeval extensional and compressional strain regimes within some tens of km. In particular, mantle upwelling cannot easily explain why the Messina trough was generated in the Pleistocene, considering that the presumed retreat of the Ionian slab and the related mantle upwelling under the Calabrian Arc was already active in the Pliocene, as suggested by the timing of the Vavilov basin.

Another important aspect of the tectonic setting that can help recognize the driving force in the study area is the occurrence of compressional deformation along the outer border of the Peloritani sector in the Pleistocene [20,65]. Since this belt sector was affected by extensional deformation in the Pliocene, any proposed driving force must provide a plausible explanation of why that regime completely changed in the Pleistocene. The geodynamic interpretation adopted here (Figures 6 and 7) can account for this change, as discussed earlier, whereas mantle upwelling can hardly account for the timing of the above strain regimes.

Another hypothesis about the tectonic mechanism responsible for the generation of the Messina trough is advanced by [20], which suggests that such deformation may be related to a transfer zone located between the SE ward retreating Ionian slab and the advancing slab under the Hyblean-Sicilian domain. The tectonic scheme shown in the inset of Figure 1 of [20] indicates that the transition from the thrust front (along the Tyrrhenian offshore of Sicily) and the extensional zones (in Calabria) corresponds to the Messina trough. However, no shear faults are reported in that zone, where only normal faults are recognized (Figure 2). It seems to be more plausible supposing that the decoupling between the Sicilian and Ionian slabs is accommodated by the Vulcano fault, where a dextral transpressional regime is clearly recognized (e.g., [26,59]). However, this fault does not mark the transition from the northern Sicilian compressional front to the Calabrian extensional zones (Figure 2), as it cuts the long compressional front that runs from Ustica to the Messina zone, and thus it cannot support the geodynamic interpretation mentioned above.

The deformations cited above are compatible with the tectonic reconstruction here proposed (Figures 3, 4, 6 and 7) since the expected northward motion of the Peloritani microplate (stressed by the Hyblean block) implies thrusting along the northern boundary of this belt sector since the Early–Middle Pleistocene. In this view, the expected divergence between Peloritani and Calabria is supposed to have driven extension in the Messina zone (Figure 2). The angular shape of the Messina sphenocasm (Figure 7) is compatible with the genetic mechanism we propose (involving a clockwise rotation of the Peloritani sector) and with the fact that the most intense extension in such trough migrated from the southern branch, in the Early Pleistocene, to the northern one, in the Middle–Late Pleistocene (e.g., [18]).

5. Conclusions

The tectonic mechanism that generated the Messina trough was triggered by the Early Pleistocene interaction between the Calabrian wedge and the continental Adriatic domain. This event determined the activation of two major decoupling zones, the Sibari and Vulcano faults. By these new lateral guides, the Calabrian wedge changed its escape trend, at the expense of the remaining Ionian domain, while the Hyblean block accelerated its northward motion, dragging the Peloritani belt sector. As a consequence, such a sector ceased to be a part of the Calabrian wedge and became an almost independent microplate, undergoing northward motion and clockwise rotation. The proposed kinematic pattern of the Peloritani belt sector can explain why since the Early Pleistocene the outer border of this structure has undergone compressional deformation [20].

This microplate kinematics involved the divergence between the Peloritani and the Calabrian wedge, which was undergoing an outward migration. The above interpretation may help to identify the genetic mechanism of the normal faults recognized in the Messina area (Figure 2), most probably representing the main seismogenetic sources of the last two major earthquakes.

The longest normal fault recognized by geological evidence in the Strait zone (Messina–Taormina fault), lying offshore the Sicilian coast, may be the source of the 1908 earthquake. Assuming this east dipping fault location, with a 5 m dip-slip at depth, it is possible to reproduce the coseismic vertical movements caused by the 1908 earthquake [13]. The same fault can explain the uplift of the shorelines that outcrop between the Messina and Taormina towns. Seismic soundings show that the above fault offsets the pre-Pleistocene basement in the Messina sphenocasm. The location of the above seismic source can explain why no ruptures have been observed onshore [96]. Moreover, the uplifted marine terraces on the footwall of the Messina–Taormina fault [97] imply a marked throw-rate change at about 50 ka (acceleration from 0.96 to 2.34 mm/y), while the slip rate decreased in the other late Quaternary faults, as the Armo and Reggio Calabria ones. This implies a recent re-organization of strain in the Messina Straits and, therefore, provides relevant information about seismic hazards in this area. In this regard, it is useful to point out that the macroseismic field of the 1783 (6 February) earthquake suggests an epicentre located in the northern part of the Messina trough, perhaps related to the Scilla fault [9].

Author Contributions: Conceptualization and methodology, E.M. and M.V.; investigation and data curation, E.M., M.V., D.B. and C.T.; writing, E.M.; funding acquisition, E.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Regione Toscana (Italy), Department of Seismic Prevention, grant number: B65F19003190002.

Data Availability Statement: This information was reported by the cited references.

Acknowledgments: We are grateful to two reviewers for their suggestions, which have allowed a considerable improvement of the work.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Peruzza, L.; Pantosti, D.; Slejko, D.; Valensise, G. Testing a new hybrid approach to seismic hazard assessment: An application to the Calabrian Arc (Southern Italy). *Nat. Hazards* **1997**, *14*, 113–126. [[CrossRef](#)]
2. Schick, R. *Eine Seismotektonische Bearbeitung des Erdbebens von Messina-im Jahre 1908*; Geologisches Jahrbuch Reihe E, Band E 11; Schweizerbart Science Publishers: Stuttgart, Germany, 1977; 74p, ISBN 978-3-510-96074-3.
3. Mulargia, F.; Boschi, E. The 1908 Messina earthquake and related seismicity. In *Earthquakes: Observation, Theory and Interpretation, Proceedings of the E. Fermi International School in Physics, Course 85, Varenna*; Kanamori, H., Boschi, E., Eds.; Springer: Amsterdam, The Netherlands; New York, NY, USA, 1983; pp. 493–518.
4. Bottari, A.; Carapezza, E.; Carapezza, M.; Carveni, P.; Cefali, F.; Lo Giudice, E.; Pandolfo, C. The 1908 Messina Strait earthquake in the regional geostructural framework. *J. Geodyn.* **1986**, *5*, 275–302.
5. Capuano, B.Y.P.; Natale, G.D.E.; Gasparini, P.; Pingue, F.; Scarpa, R. A model for the 1908 Messina Straits (Italy) earthquake by inversion of levelling data. *Bull. Seismol. Soc. Am.* **1988**, *78*, 1930–1947.

6. Boschi, E.; Pantosti, D.; Valensise, G. Modello di sorgente per il terremoto di Messina del 1908 ed evoluzione recente dell'area dello Stretto. In Proceedings of the Atti 89th GNDT Workshop, Roma, Italy, 7–9 November 1989; pp. 245–258.
7. De Natale, G.; Pingue, F. A Variable Slip Fault Model For the 1908 Messina Straits (Italy) Earthquake, By Inversion of Levelling Data. *Geophys. J. Int.* **1991**, *104*, 73–84. [[CrossRef](#)]
8. Valensise, G.; Pantosti, D. A 125 Kyr-long geological record of seismic source repeatability: The Messina Straits (Southern Italy) and the 1908 earthquake (Ms 7^{1/2}). *Terra Nova* **1992**, *4*, 472–483. [[CrossRef](#)]
9. Cucci, L.; D'Addezio, G.; Valensise, G.; Burrato, P. Investigating seismogenic faults in Central and Southern Apennines (Italy): Modeling of fault-related landscape features. *Ann. Geofis.* **1996**, *39*, 603–618.
10. Amoroso, A.; Crescentini, L.; Scarpa, R. Source parameters of the 1908 Messina Straits, Italy, earthquake from geodetic and seismic data. *J. Geophys. Res. Solid Earth* **2002**, *107*, ESE 4-1–ESE 4-11. [[CrossRef](#)]
11. Aloisi, M.; Bruno, V.; Cannavò, F.; Ferranti, L.; Mattia, M.; Monaco, C.; Palano, M. Are the source models of the M 7.1 1908 Messina Straits earthquake reliable? Insights from a novel inversion and sensitivity analysis of levelling data. *Geophys. J. Int.* **2013**, *192*, 1025–1041. [[CrossRef](#)]
12. Convertito, V.; Pino, N.A. Discriminating among distinct source models of the 1908 Messina Straits earthquake by modelling intensity data through full waveform seismograms. *Geophys. J. Int.* **2014**, *198*, 164–173. [[CrossRef](#)]
13. Meschis, M.; Roberts, G.P.; Mildon, Z.K.; Robertson, J.; Michetti, A.M.; Faure Walker, J.P. Slip on a Mapped normal Fault for the 28th December 1908 Messina Earthquake (Mw 7.1) in Italy. *Sci. Rep.* **2019**, *9*, 6481. [[CrossRef](#)]
14. Barreca, G.; Gross, F.; Scarfi, L.; Aloisi, M.; Monaco, C.; Krastel, S. The Strait of Messina: Seismotectonics and the Source of the 1908 Earthquake. *Earth-Sci. Rev.* **2021**, *218*, 103685. [[CrossRef](#)]
15. Andrenacci, C.; Bello, S.; Barbano, M.S.; de Nardis, R.; Pirrotta, C.; Pietrolungo, F.; Lavecchia, G. Reappraisal and Analysis of Macroseismic Data for Seismotectonic Purposes: The Strong Earthquakes of Southern Calabria, Italy. *Geosciences* **2023**, *13*, 212. [[CrossRef](#)]
16. Loperfido, A. Livellazione geometrica di precisione eseguita dall'IGM sulla costa orientale della Sicilia, da Messina a Catania, a Gesso ed a Faro Peloro e sulla costa occidentale della Calabria da Gioia Tauro a Melito di Porto Salvo. In *Relazione della Commissione Reale Incaricata di Designare le Zone più Adatte per la Ricostruzione Degli Abitanti Colpiti dal Terremoto del 28 Dicembre 1908 o da Altri Precedenti (1909)*; C.R. Accademia Naz. Lincei: Roma, Italy, 1909; Volume 35, pp. 131–156.
17. Del Ben, A.; Gargano, C.; Lentini, R. Ricostruzione strutturale e stratigrafica dell'area dello Stretto di Messina mediante analisi comparata dei dati geologici e sismici. *Mem. Soc. Geol. It.* **1996**, *51*, 703–717.
18. Del Ben, A.; Barnaba, C.; Toboga, A. Strike-slip systems as the main tectonic features in the Plio-Quaternary kinematics of the Calabrian Arc. *Mar. Geophys. Res.* **2008**, *29*, 1–12. [[CrossRef](#)]
19. Scarfi, L.; Langer, H.; Scaltrito, A. Seismicity, seismotectonics and crustal velocity structure of the Messina Strait (Italy). *Phys. Earth Planet. Int.* **2009**, *177*, 65–78. [[CrossRef](#)]
20. Doglioni, C.; Ligi, M.; Scrocca, D.; Bigi, S.; Bortoluzzi, G.; Carminati, E.; Cuffaro, M.; D'Oriano, F.; Forleo, V.; Muccini, F.; et al. The tectonic puzzle of the Messina area (Southern Italy): Insights from new seismic reflection data. *Sci. Rep.* **2012**, *2*, 970. [[CrossRef](#)] [[PubMed](#)]
21. Malinverno, A.; Ryan, W.B.F. Extension on the Tyrrhenian Sea and shortening in the Apennines as results of arc migration driven by sinking of the lithosphere. *Tectonics* **1986**, *5*, 227–245. [[CrossRef](#)]
22. Jolivet, L.; Faccenna, C. Mediterranean extension and the Africa-Eurasia collision. *Tectonics* **2000**, *19*, 1095–1106. [[CrossRef](#)]
23. Faccenna, C.; Funiciello, F.; Civetta, L.; D'Antonio, M.; Moroni, M.; Piromallo, C. Slab disruption, mantle circulation, and the opening of the Tyrrhenian basins. In *Cenozoic Volcanism in the Mediterranean Area*; Beccaluva, L., Bianchini, G., Wilson, M., Eds.; Geological Society of America Special Papers; AGU: Washington, DC, USA, 2007; Volume 418, pp. 153–169. [[CrossRef](#)]
24. Faccenna, C.; Becker, T.W.; Auer, L.; Billi, A.; Boschi, L.; Brun, J.P.; Capitanio, F.A.; Funiciello, F.; Horvath, F.; Jolivet, L.; et al. Mantle dynamics in the Mediterranean. *Rev. Geophys.* **2014**, *52*, 283–332. [[CrossRef](#)]
25. Gvirtzman, Z.; Nur, A. Plate detachment, asthenosphere upwelling, and topography across subduction zones. *Geology* **1999**, *27*, 563–566. [[CrossRef](#)]
26. Palano, M.; Ferranti, L.; Monaco, C.; Mattia, M.; Aloisi, M.; Bruno, V.; Cannavò, F.; Siligato, G. GPS velocity and strain fields in Sicily and Southern Calabria, Italy: Updated geodetic constraints on tectonic block interaction in the central Mediterranean. *J. Geophys. Res.* **2012**, *117*, B07401. [[CrossRef](#)]
27. Mantovani, E.; Viti, M.; Babbucci, D.; Tamburelli, C.; Cenni, N. How and Why the Present Tectonic Setting in the Apennine Belt Has Developed. *J. Geol. Soc. Lond.* **2019**, *176*, 1291. [[CrossRef](#)]
28. Mantovani, E.; Viti, M.; Babbucci, D.; Tamburelli, C.; Cenni, N. Geodynamics of the Central Western Mediterranean Region: Plausible and Non-Plausible Driving Forces. *Mar. Pet. Geol.* **2020**, *113*, 104121. [[CrossRef](#)]
29. Viti, M.; Mantovani, E.; Babbucci, D.; Tamburelli, C.; Caggiati, M.; Riva, A. Basic Role of Extrusion Processes in the Late Cenozoic Evolution of the Western and Central Mediterranean Belts. *Geosciences* **2021**, *11*, 499. [[CrossRef](#)]
30. SgROI, T.; Polonia, A.; Barberi, G.; Billi, A.; Gasperini, L. New seismological data from the Calabrian arc reveal arc-orthogonal extension across the subduction zone. *Sci. Rep.* **2021**, *11*, 473. [[CrossRef](#)]
31. Monaco, C.; Tortorici, L. Active faulting in the Calabrian arc and eastern Sicily. *J. Geodyn.* **2000**, *29*, 407–424. [[CrossRef](#)]
32. Catalano, S.; De Guidi, G.; Monaco, C.; Tortorici, G.; Tortorici, L. Active faulting and seismicity along the Siculo-Calabrian Rif Zone (Southern Italy). *Tectonophysics* **2008**, *453*, 177–192. [[CrossRef](#)]

33. Faure Walker, J.P.; Roberts, G.P.; Cowie, P.A.; Papanikolaou, I.; Michetti, A.M.; Sammonds, P.; Wilkinson, M.; McCaffrey, K.J.W.; Phillips, R.J. Relationship between topography, rates of extension and mantle dynamics in the actively-extending Italian Apennines. *Earth Planet. Sci. Lett.* **2012**, *325–326*, 76–84. [[CrossRef](#)]
34. Koçyiğit, A.; Yılmaz, A.; Adamia, S.; Kuloshvili, S. Neotectonics of East Anatolian Plateau (Turkey) and Lesser Caucasus: Implication for transition from thrusting to strike-slip faulting. *Geodin. Acta* **2001**, *14*, 177–195. [[CrossRef](#)]
35. Hubert-Ferrari, A.; King, G.; Manighetti, I.; Armijo, R.; Meyer, B.; Tapponnier, P. Long-term elasticity in the continental lithosphere; modelling the Aden Ridge propagation and the Anatolian extrusion process. *Geophys. J. Int.* **2003**, *153*, 111–132. [[CrossRef](#)]
36. Hippolyte, J.C.; Espurt, N.; Kaymakci, N.; Sangu, E.; Müller, C. Cross-sectional anatomy and geodynamic evolution of the Central Pontide orogenic belt (Northern Turkey). *Int. J. Earth. Sci. (Geol. Rundsch.)* **2016**, *105*, 81–106. [[CrossRef](#)]
37. Mantovani, E.; Babbucci, D.; Tamburelli, C.; Viti, M. Late Cenozoic Evolution and Present Tectonic Setting of the Aegean–Hellenic Arc. *Geosciences* **2022**, *12*, 104. [[CrossRef](#)]
38. Robertson, A.; Shallo, M. Mesozoic-Tertiary Tectonic Evolution of Albania in Its Regional Eastern Mediterranean Context. *Tectonophysics* **2000**, *316*, 197–254. [[CrossRef](#)]
39. Royden, L.H.; Papanikolaou, D.J. Slab Segmentation and Late Cenozoic Disruption of the Hellenic Arc. *Geochem. Geophys. Geosyst.* **2011**, *12*, Q03010. [[CrossRef](#)]
40. Burchfiel, B.C.; Royden, L.H.; Papanikolaou, D.; Pearce, F.D. Crustal Development within a Retreating Subduction System: The Hellenides. *Geosphere* **2018**, *14*, 1119–1130. [[CrossRef](#)]
41. Mantovani, E.; Babbucci, D.; Tamburelli, C.; Viti, M. A review on the driving mechanism of the Tyrrhenian–Apennines system: Implications for the present seismotectonic setting in the Central-Northern Apennines. *Tectonophysics* **2009**, *476*, 22–40. [[CrossRef](#)]
42. Mantovani, E.; Viti, M.; Babbucci, D.; Albarello, D. Nubia-Eurasia Kinematics: An Alternative Interpretation from Mediterranean and North Atlantic Evidence. *Ann. Geophys.* **2007**, *50*, 311–336. [[CrossRef](#)]
43. Viti, M.; Mantovani, E.; Babbucci, D.; Tamburelli, C. Plate Kinematics and Geodynamics in the Central Mediterranean. *J. Geodyn.* **2011**, *51*, 190–204. [[CrossRef](#)]
44. Hieke, W.; Hirschleber, H.B.; Deghani, G.A. The Ionian Abyssal Plain (central Mediterranean Sea): Morphology, sub-bottom structures and geodynamic history—An inventory. *Mar. Geophys. Res.* **2003**, *24*, 279–310. [[CrossRef](#)]
45. Hieke, W.; Cita, M.B.; Forcella, F.; Muller, C. Geology of the Victor Hensen Seahill (Ionian Sea, eastern Mediterranean): Insights from the study of cored sediment sequences. *Boll. Della Soc. Geol. Ital.* **2006**, *125*, 245–257.
46. Finetti, I.R.; Lentini, F.; Carbone, S.; Del Ben, A.; Di Stefano, A.; Forlin, E.; Guarneri, P.; Pipan, M.; Prizzon, A. Geological Outline of Sicily and Lithospheric Tectono-Dynamics of its Tyrrhenian Margin from New CROP Seismic Data. In *CROP PROJECT, Deep Seismic Exploration of the Central Mediterranean and Italy*; Finetti, I.R., Ed.; Elsevier: Amsterdam, The Netherlands, 2005; Chapter 15; pp. 319–376.
47. Pepe, F.; Sulli, A.; Bertotti, G.; Catalano, R. Structural highs formation and their relationship to sedimentary basins in the North Sicily continental margin (Southern Tyrrhenian Sea): Implications for the Drepano Thrust. *Front. Tectonophysics* **2005**, *409*, 1–18. [[CrossRef](#)]
48. Civile, D.; Lodolo, E.; Accettella, D.; Geletti, R.; Ben-Avraham, Z.; Deponete, M.; Facchin, L.; Ramella, R.; Romeo, R. The Pantelleria graben (Sicily Channel, Central Mediterranean): An example of intraplate ‘passive’ rift. *Tectonophysics* **2010**, *490*, 173–183. [[CrossRef](#)]
49. Civile, D.; Lodolo, E.; Accaino, F.; Geletti, R.; Schiattarella, M.; Giustiniani, M.; Fedorik, J.; Zecchin, M.; Zampa, L. Capo granitola-Sciaccia fault zone (Sicilian channel, central Mediterranean): Structure vs. magmatism. *Mar. Pet. Geol.* **2018**, *96*, 627–644. [[CrossRef](#)]
50. Carmignani, L.; Decandia, F.A.; Fantozzi, P.L.; Lazzarotto, A.; Liotta, D.; Meccheri, M. Tertiary Extensional Tectonics in Tuscany (Northern Apennines, Italy). *Tectonophysics* **1994**, *238*, 295–315. [[CrossRef](#)]
51. Casula, G.; Cherchi, A.; Montadert, L.; Murru, M.; Sarria, E. The Cenozoic Graben System of Sardinia (Italy): Geodynamic Evolution from New Seismic and Field Data. *Mar. Pet. Geol.* **2001**, *18*, 863–888. [[CrossRef](#)]
52. Cocco, F.; Funedda, A.; Patacca, E.; Scandone, P. Plio-Pleistocene Extensional Tectonics in the Campidano Graben (SW Sardinia, Italy): Preliminary Note. *Rend. Online Della Soc. Geol. Ital.* **2013**, *29*, 31–34.
53. Finetti, I.R.; Boccaletti, M.; Bonini, M.; Del Ben, A.; Pipan, M.; Prizzon, A.; Sani, F. Lithospheric Tectono-Stratigraphic Setting of the Ligurian Sea-Northern Apennines-Adriatic Foreland from Integrated CROP Seismic Data. In *CROP PROJECT, Deep Seismic Exploration of the Central Mediterranean and Italy*; Finetti, I.R., Ed.; Elsevier: Amsterdam, The Netherlands, 2005; Chapter 8; pp. 119–158.
54. Gattacceca, J.; Deino, A.; Rizzo, R.; Jones, D.S.; Henry, B.; Beaudoin, B.; Vadeboin, F. Miocene rotation of Sardinia: New paleomagnetic and geochronological constraints and geodynamic implications. *Earth Planet. Sci. Lett.* **2007**, *258*, 359–377. [[CrossRef](#)]
55. Molli, G. Northern Apennine-Corsica Orogenic System: An Updated Overview. In *Tectonic Aspects of the Alpine-Dinarides-Carpathians System*; Siegesmund, S., Fugenschu, B., Froitzheim, N., Eds.; The Geological Society: London, UK, 2008; Volume 298, pp. 413–442.
56. Finetti, I.R.; Del Ben, A. Geophysical study of the Tyrrhenian opening. *Boll. Di Geofis. Teor. E Appl.* **1986**, *110*, 75–156.
57. Ben Avraham, Z.; Boccaletti, M.; Cello, G.; Grasso, M.; Lentini, F.; Torelli, L.; Tortorici, L. Principali domini strutturali originatisi dalla collisione neogenico-quadernaria nel mediterraneo centrale. *Mem. Soc. Geol. Ital.* **1990**, *45*, 453–462.

58. Lentini, F.; Carbone, S.; Catalano, S. Main structural domains of the central Mediterranean region and their tectonic evolution. *Boll. Geofis. Teor. Appl.* **1994**, *36*, 103–125.
59. Argnani, A.; Serpelloni, E.; Bonazzi, C. Pattern of deformation around the central Aeolian Islands: Evidence from multichannel seismics and GPS data. *Terra Nova* **2007**, *19*, 317–323. [[CrossRef](#)]
60. Zecchin, M.; Praeg, D.; Ceramicola, S.; Muto, F. Onshore to offshore correlation of regional unconformities in the Plio-Pleistocene sedimentary successions of the Calabrian Arc (central Mediterranean). *Earth-Sci. Rev.* **2015**, *142*, 60–78. [[CrossRef](#)]
61. Sulli, A.; Lo Presti, V.; Morticelli, M.G.; Antonioli, F. Vertical movements in NE Sicily and its offshore: Outcome of tectonic uplift during the last 125 ky. *Quat. Int.* **2013**, *288*, 168–182. [[CrossRef](#)]
62. Ferranti, L.; Burrato, P.; Pepe, F.; Santoro, E.; Mazzella, M.E.; Morelli, D.; Passaro, S.; Vannucci, G. An active oblique-contractional belt at the transition between the Southern Apennines and Calabrian Arc: The Amendolara Ridge, Ionian Sea, Italy. *Tectonics* **2014**, *33*, 2169–2194. [[CrossRef](#)]
63. Volpi, V.; Del Ben, A.; Civile, D.; Zgur, F. Neogene tectono-sedimentary interaction between the Calabrian Accretionary Wedge and the Apulian foreland in the northern Ionian Sea. *Mar. Pet. Geol.* **2017**, *83*, 246–260. [[CrossRef](#)]
64. Teofilo, G.; Antoncicchi, I.; Caputo, R. Neogene-Quaternary evolution of the offshore sector of the Southern Apennines accretionary wedge, Gulf of Taranto, Italy. *Tectonophysics* **2018**, *738–739*, 16–32. [[CrossRef](#)]
65. Scacchia, E.; Tinterri, R.; Gamberi, F. The Influence of Channel Planform and Slope Topography on Turbidity Current Overbank Processes: The Example of the Acquarone Fan (Southeastern Tyrrhenian Sea). *Front. Earth Sci.* **2022**, *9*, 785164. [[CrossRef](#)]
66. Sulli, A. Structural framework and crustal characteristics of the Sardinia Channel Alpine transect in the central Mediterranean. *Tectonophysics* **2000**, *324*, 321–336. [[CrossRef](#)]
67. Sulli, A.; Morticelli, M.G.; Agate, M.; Zizzo, E. Active north-vergent thrusting in the northern Sicily continental margin in the frame of the Quaternary evolution of the Sicilian collisional system. *Tectonophysics* **2021**, *802*, 228717. [[CrossRef](#)]
68. Guarnieri, P. Structural evidence for deformation by block rotation in the context of transpressive tectonics, northwestern Sicily (Italy). *J. Struct. Geol.* **2004**, *26*, 207–219. [[CrossRef](#)]
69. Pondrelli, S.; Salimbeni, S.; Ekström, G.; Morelli, A.; Gasperini, P.; Vannucci, G. The Italian CMT dataset from 1977 to the present. *Phys. Earth Planet. Inter.* **2006**, *159*, 286–303. [[CrossRef](#)]
70. Grevemeyer, I.; Ranero, C.; Zitellini, N.; Sallares, V.; Prada, M. Seismic wide-angle constrains on the structure of the northern Sicily margin and Vavilov Basin: Implications for the opening of the Tyrrhenian back-arc basin. In Proceedings of the EGU General Assembly 2020, Online, 4–8 May 2020. EGU2020-5130. [[CrossRef](#)]
71. Loreto, M.F.; Palmiotto, C.; Muccini, F.; Ferrante, V.; Zitellini, N. Inverted Basins by Africa–Eurasia Convergence at the Southern Back-Arc Tyrrhenian Basin. *Geosciences* **2021**, *11*, 117. [[CrossRef](#)]
72. Rovida, A.; Locati, M.; Camassi, R.; Lolli, B.; Gasperini, P.; Antonucci, A. (Eds.) *Italian Parametric Earthquake Catalogue (CPTI15); Version 4.0*; Istituto Nazionale di Geofisica e Vulcanologia (INGV): Roma, Italy, 2022. [[CrossRef](#)]
73. Funicello, R.; Parotto, M.; Praturlon, A.; Coord. *Carta Tettonica d'Italia alla scala 1:1500000*; CNR Progetto Finalizzato Geodinamica, Pubbl. 269; Grafica Editoriale Cartografica: Roma, Italy, 1981.
74. Mantovani, E.; Viti, M.; Babbucci, D.; Tamburelli, C.; Cenni, N.; Baglione, M.; D'Intinosante, V. Generation of back-Arc Basins as Side Effect of Shortening Processes: Examples from the Central Mediterranean. *Int. J. Geosci.* **2014**, *5*, 1062–1079. [[CrossRef](#)]
75. Cucci, L. Raised marine terraces in the Northern Calabrian Arc (Southern Italy): A 600 kyr long geological record of regional uplift. *Ann. Geophys.* **2004**, *47*, 1391–1406. [[CrossRef](#)]
76. Zecchin, M.; Nalin, R.; Roda, C. Raised Pleistocene marine terraces of the Crotona peninsula (Calabria, Southern Italy): Facies analysis and organization of their deposits. *Sediment. Geol.* **2004**, *172*, 165–185. [[CrossRef](#)]
77. Antonioli, F.; Ferranti, L.; Lambeck, K.; Kershaw, S.; Verrubbi, V.; Dai Pra, G. Late Pleistocene to Holocene record of changing uplift rates in Southern Calabria and northeastern Sicily (Southern Italy, Central Mediterranean Sea). *Tectonophysics* **2006**, *422*, 23–40. [[CrossRef](#)]
78. Ferranti, L.; Monaco, C.; Antonioli, F.; Maschio, L.; Kershaw, S.; Verrubbi, V. The contribution of regional uplift and coseismic slip to the vertical crustal motion in the Messina Straits, Southern Italy: Evidence from raised Late Holocene shorelines. *J. Geophys. Res.* **2007**, *112*, B06401. [[CrossRef](#)]
79. Ferranti, L.; Santoro, E.; Mazzella, M.E.; Monaco, C.; Morelli, D. Active transpression in the northern Calabria Apennines, Southern Italy. *Tectonophysics* **2009**, *476*, 226–251. [[CrossRef](#)]
80. Shemenda, A.I. Subduction of the lithosphere and back arc dynamics: Insights from physical modelling. *J. Geophys. Res.* **1993**, *98*, 16167–16185. [[CrossRef](#)]
81. Hassani, R.; Jongmans, D.; Chery, J. Study of plate deformation and stress in subduction processes using two dimensional numerical models. *J. Geophys. Res.* **1997**, *102*, 17951–17965. [[CrossRef](#)]
82. Buiter, S.J.H.; Govers, R.; Wortel, M.J.R. A modelling study of vertical surface displacements at convergent plate margins. *Geophys. J. Int.* **2001**, *147*, 415–427. [[CrossRef](#)]
83. Hampel, A.; Pfiffner, A. Relative importance of trenchward upper plate motion and friction along the plate interface for the topographic evolution of subduction related mountain belts. In *Analogue and Numerical Modelling of Crustal-Scale Processes*; Buiter, S.J.H., Schreurs, G., Eds.; Geological Society: London, UK, 2006; Volume 253, pp. 105–115. [[CrossRef](#)]
84. Husson, L. Dynamic topography above retreating subduction zones. *Geology* **2006**, *34*, 741–744. [[CrossRef](#)]

85. Sartori, R. The Main Results of ODP Leg 107 in the Frame of Neogene to Recent Geology of Perityrrhenian Areas. *Proc. ODP Sci. Results Coll. Stn. Tx.* **1990**, *107*, 715–730.
86. Finetti, I.R.; Del Ben, A. Crustal Tectono-Stratigraphy of the Ionian Sea from New Integrated CROP Seismic Data. In *CROP PROJECT, Deep Seismic Exploration of the Central Mediterranean and Italy*; Finetti, I.R., Ed.; Elsevier: Amsterdam, The Netherlands, 2005; Chapter 19; pp. 447–470.
87. Savelli, C. Time–Space Distribution of Magmatic Activity in the Western Mediterranean and Peripheral Orogens during the Past 30 Ma (A Stimulus to Geodynamic Considerations). *J. Geodyn.* **2002**, *34*, 99–126. [[CrossRef](#)]
88. Sartori, R.; Torelli, L.; Zitellini, N.; Carrara, G.; Magaldi, M.; Mussoni, P. Crustal features along a W-E Tyrrhenian transect from Sardinia to Campania margins (Central Mediterranean). *Tectonophysics* **2004**, *383*, 171–192. [[CrossRef](#)]
89. Corradino, M.; Balazs, A.; Faccenna, C.; Pepe, F. Arc and forearc rifting in the Tyrrhenian subduction system. *Sci. Rep.* **2022**, *12*, 4728. [[CrossRef](#)]
90. Finetti, I.R. The Calabrian Arc and Subducting Ionian Slab from new CROP Seismic Data. In *CROP PROJECT, Deep Seismic Exploration of the Central Mediterranean and Italy*; Finetti, I.R., Ed.; Elsevier: Amsterdam, The Netherlands, 2005; Chapter 17; pp. 393–412.
91. Finetti, I.R.; Lentini, F.; Carbone, S.; Del Ben, A.; Di Stefano, A.; Guarnieri, P.; Pipan, M.; Prizzon, A. Crusta1 Tectono-Stratigraphy and Geodynamics of the Southern Apennines from CROP and Other Integrated Geophysical-Geological Data. In *CROP PROJECT, Deep Seismic Exploration of the Central Mediterranean and Italy*; Finetti, I.R., Ed.; Elsevier: Amsterdam, The Netherlands, 2005; Chapter 12; pp. 225–262.
92. Wortel, M.J.R.; Spakman, W. Subduction and slab detachment in the Mediterranean-Carpathian region. *Science* **2000**, *290*, 1910–1917. [[CrossRef](#)]
93. Doglioni, C.; Innocenti, F.; Mariotti, S. Why Mt. Etna? *Terra Nova* **2001**, *13*, 25–31. [[CrossRef](#)]
94. Goes, S.; Giardini, D.; Jenny, S.; Hollenstein, C.; Kahle, H.G.; Geiger, A. A recent tectonic reorganization in the south-central Mediterranean. *Earth Planet. Sci. Lett.* **2004**, *226*, 335–345. [[CrossRef](#)]
95. Gvirtzman, Z.; Nur, A. The formation of Mount Etna as the consequence of slab rollback. *Nature* **1999**, *401*, 782–785. [[CrossRef](#)]
96. Comerci, V.; Vittori, E.; Blumetti, A.M.; Brustia, E.; Di Manna, P.; Guerrieri, L.; Lucarini, M.; Serva, L. Environmental effects of the December 28, 1908, Southern Calabria–Messina (Southern Italy) earthquake. *Nat. Hazards* **2015**, *76*, 1849–1891. [[CrossRef](#)]
97. Meschis, M.; Roberts, G.P.; Robertson, J.; Mildon, Z.K.; Sahy, D.; Goswami, R.; Sgambato, C.; Walker, J.F.; Michetti, A.M.; Iezzi, F. Out of phase Quaternary uplift-rate changes reveal normal fault interaction, implied by deformed marine palaeoshorelines. *Geomorphology* **2022**, *416*, 108432. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.